



Correction to: Llama peripheral B-cell populations producing conventional and heavy chain-only IgG subtypes are phenotypically indistinguishable but immunogenetically distinct

Kevin A. Henry¹ · Henk van Faassen¹ · Doreen Harcus² · Anne Marcil² · Jennifer J. Hill¹ · Serge Muyldermans³ · C. Roger MacKenzie^{1,4}

Published online: 13 February 2023
© Crown 2023

Correction to: Immunogenetics (2019) 71:307–320
<https://doi.org/10.1007/s00251-018-01102-9>

At the time of publication approval could not be obtained to disclose the amino acid sequences of the anti-isotypic mAbs 1C10 and 5E4. These are now provided in Supplementary Table SV.

Supplementary Information The online version contains supplementary material available at <https://doi.org/10.1007/s00251-023-01299-4>.

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

The original article can be found online at <https://doi.org/10.1007/s00251-018-01102-9>.

✉ Kevin A. Henry
kevin.henry@nrc-cnrc.gc.ca

¹ Human Health Therapeutics Research Centre, National Research Council Canada, 100 Sussex Drive, Ottawa, ON K1A 0R6, Canada

² Human Health Therapeutics Research Centre, National Research Council Canada, 6100 Royalmount Avenue, Montréal, QC H4P 2R2, Canada

³ Laboratory of Cellular and Molecular Immunology, Vrije Universiteit Brussel, Brussels, Belgium

⁴ School of Environmental Sciences, University of Guelph, 50 Stone Road East, Guelph, ON N1G 2W1, Canada