



Correction to: Regulation of human brown adipose tissue by adenosine and A_{2A} receptors – studies with [¹⁵O]H₂O and [¹¹C]TMSX PET/CT

Minna Lahesmaa^{1,2} · Vesa Oikonen¹ · Semi Helin¹ · Pauliina Luoto¹ · Mueez U Din^{1,2} · Alexander Pfeifer³ · Pirjo Nuutila^{1,4} · Kirsi A. Virtanen^{1,2}

Published online: 8 September 2018

© Springer-Verlag GmbH Germany, part of Springer Nature 2018

Correction to: European Journal of Nuclear Medicine and Molecular Imaging

<https://doi.org/10.1007/s00259-018-4120-2>

The original version of this article contained a mistake in the first sentence of the Results section of the Abstract. The line “Adenosine increased the perfusion of BAT even more than cold exposure (baseline 8.3 ± 4.5 , cold 19.6 ± 9.3 , adenosine 28.6 ± 7.9 $\mu\text{mol}/100$ g/min, $p < 0.01$).” should be “Adenosine increased the perfusion of BAT even more than cold exposure (baseline 8.3 ± 4.5 , cold 19.6 ± 9.3 , adenosine 28.6 ± 7.9 ml/100 g/min, $p < 0.01$).”

The original article has been corrected.

The online version of the original article can be found at <https://doi.org/10.1007/s00259-018-4120-2>

✉ Kirsi A. Virtanen
kirsi.virtanen@utu.fi

¹ Turku PET Centre, University of Turku, P.O. Box 52, FI-20520 Turku, Finland

² Turku PET Centre, Turku University Hospital, Turku, Finland

³ Institute of Pharmacology and Toxicology, University of Bonn, Bonn, Germany

⁴ Department of Endocrinology, Turku University Hospital, Turku, Finland