

ERRATUM


# Erratum to: Leaf area lost, rather than herbivore type, determines the induction of extrafloral nectar secretion in a tropical plant (*Clerodendrum philippinum*)

Fang Fang Xu<sup>1</sup> · Jin Chen<sup>1</sup> · John Husson<sup>2</sup>

Published online: 30 April 2015  
© Springer Science+Business Media Dordrecht 2015

**Erratum to: Arthropod-Plant Interactions (2014)  
8:513–518  
DOI 10.1007/s11829-014-9331-2**

In the original publication of the article, the y axis of the Fig. 1 should be “ $S_{\text{After herbivore}} - S_{\text{Before herbivore}}$ ”, not “ $S_{\text{Before herbivore}} - S_{\text{After herbivore}}$ ”. The corrected Fig. 1 is given below.



**Fig. 1** Regression of the change of EFN secretion rate on the leaf area eaten by herbivores ( $R_{\text{adj}}^2 = 0.12$ ,  $F_{1,23} = 4.38$ ,  $P = 0.0475$ ). The change of the EFN secretion was calculated as  $S_{\text{After herbivore}} - S_{\text{Before herbivore}}$ , where  $S_{\text{After herbivore}}$  = the EFN secretion rate after herbivore treatment,  $S_{\text{Before herbivore}}$  = the EFN secretion rate before herbivore treatment, measured 1 day before herbivore treatment

The online version of the original article can be found under doi:10.1007/s11829-014-9331-2.

✉ Jin Chen  
cj@xtbg.org.cn

<sup>1</sup> Key Laboratory of Tropical Forest Ecology, Xishuangbanna Tropical Botanical Garden, Chinese Academy of Sciences, Mengla 666303, Yunnan, China

<sup>2</sup> Undergraduate Program, Middlebury College, Middlebury, VT 05753, USA