

Erratum to: Different expression profiles of jasmonic acid and salicylic acid inducible genes in the tomato plant against herbivores with various feeding modes

Kei Kawazu¹ · Atsushi Mochizuki¹ · Yukie Sato¹ ·
Wataru Sugeno¹ · Mika Murata¹ · Shigemi Seo² ·
Ichiro Mitsuhashi²

Published online: 6 January 2016
© Springer Science+Business Media Dordrecht 2016

Erratum to: Arthropod-Plant Interactions (2012)
6:221–230
DOI 10.1007/s11829-011-9174-z

The authors would like to apologize for this error.
The correct version of Table 1 is given below.

Unfortunately, there is an error in the Table 1 in the original publication of the article.

Table 1 Genes and primers used for real-time PCR amplification

Gene	Forward primer: sequence (5′–3′)	Reverse primer: sequence (5′–3′)
<i>Pin2</i>	CATCTTCTGGATTGCCCA	ACACACAACCTTGATGCCAC
<i>LapA1</i>	GGGACTAATGATGTTTGGAA	GTGGCAATTTTATTTAGGCA
<i>PRb-1b</i>	TTTCCCTTTTGATGTTGCT	TGGAAACAAGAAGATGCAGT
<i>GluB</i>	CCTTTTGTGTTGCATTCTGG	GCAGTTACCAGATCAGAGTGT
<i>Actin</i>	CCAGGTATTGCTGATAGAATGAG	GAGCCTCCAATCCAGACAC

The online version of the original article can be found under doi:[10.1007/s11829-011-9174-z](https://doi.org/10.1007/s11829-011-9174-z).

✉ Atsushi Mochizuki
lepi@affrc.go.jp

¹ National Institute for Agro-Environmental Sciences,
3-1-3 Kannondai, Tsukuba, Ibaraki 305-8604, Japan

² National Institute of Agrobiological Sciences,
2-1-2 Kannondai, Tsukuba, Ibaraki 305-8602, Japan