

Erratum to: Altered NR4A Subfamily Gene Expression Level in Peripheral Blood of Parkinson's and Alzheimer's Disease Patients

Francesca Montarolo¹ · Simona Perga¹ · Serena Martire¹ · De'sire'e Nicole Navone¹ · Alberto Marchet² · Daniela Leotta² · Antonio Bertolotto¹

Published online: 2 December 2016
© Springer Science+Business Media New York 2016

Erratum to: Neurotox Res (2016) 30:338–344
DOI 10.1007/s12640-016-9626-4

The original version of this article unfortunately contained a mistake. Some of the values in Table 2 are not correctly presented. The corrected version of Table 2 is given below.

Table 2 Clinical characteristics of enrolled PD and AD populations

	PD		AD			
N	20		22			
Disease onset (y) median (range)	61.5 (36–75)		65.0 (52–81)			
Familiarity N (%)	yes	4 (20)	3 (9.2)			
	no	16 (80)	19 (81.8)			
Patients treated with specific therapies	L-DOPA	L-DOPA + Dopamine agonists	Donepezil	Memantine	Rivastigmine	Combination of two of them
N (%)	19 (95)	16 (80)	11 (50)	6 (27.3)	4 (18.2)	4 (18.2)
UPDRS at least part III median (range)	33.5 (78–16)		-			
H&Y scale median (range)	2.7 (4–1.5)		-			

HC healthy controls, *PD* Parkinson's disease, *AD*, Alzheimer's disease, *y* years, *N* number of subjects, *UPDRS* unified Parkinson's disease rating scale, *H&Y* Hoehn and Yahr scale

The online version of the original article can be found at <http://dx.doi.org/10.1007/s12640-016-9626-4>

✉ Francesca Montarolo
francesca.montarolo@unito.it

¹ Neurobiology Unit, Neurologia 2 – CReSM (Regional Referring Center of Multiple Sclerosis), Neuroscience Institute Cavalieri Ottolenghi (NICO), University of Turin & AOU San Luigi Gonzaga, Regione Gonzole 10, 10043 Orbassano, TO, Italy

² Division of Neurology, Martini Hospital, Via Tofane 71, 10141 Turin, TO, Italy