



## Correction to: Analysis of bronopol (2-bromo-2-nitropropan-1,3-diol) residues in rice (*Oryza sativa* L.) by SPE using Bond Elut Plexa and liquid chromatography-tandem mass spectrometry

Shuang-shuang Chai<sup>1</sup> · Mei-ling Qin<sup>1</sup> · You-ning Ma<sup>1</sup> · Huan-huan Gao<sup>1</sup> · Qiao He<sup>1</sup> · Han-tong Zhang<sup>1</sup>

Received: 12 January 2018 / Accepted: 12 January 2018 / Published online: 13 February 2018  
© Springer-Verlag GmbH Germany, part of Springer Nature 2018

### Correction to: Anal Bioanal Chem

<https://doi.org/10.1007/s00216-017-0730-y>

The authors would like to call the reader's attention to the fact that unfortunately during a recent cross-check of the experimental record, they found that the positions of intercept and slope were reversed in Table 1 in the original manuscript. The authors apologize for the mistake.

Please find below the corrected version of Table 1.

**Table 1** Validation of the proposed method

Compound	LOD <sup>a</sup> ( $\mu\text{g kg}^{-1}$ )	LOQ <sup>b</sup> ( $\mu\text{g kg}^{-1}$ )	Linear range ( $\mu\text{g L}^{-1}$ )	Regression line			Lack-of-fit test		Spiked level ( $\mu\text{g kg}^{-1}$ )	Recovery (%)	RSD (%)	Matrix effect <sup>c</sup>
				Intercept ( $\times 10^2$ )	Slope ( $\times 10^2$ )	$R^2$	F value	$p$ value				
Bronopol	1.0	3.3	5.0–625.0	−7.1	31.4	0.9992	1.13	0.383	25.0	73.3	2.4	1.01
									125.0	84.9	7.9	
									625.0	96.7	1.2	

<sup>a</sup> Method limit of detection

<sup>b</sup> Method limit of quantification

<sup>c</sup> Matrix effect =  $\text{slope}_{\text{matrix-matched standard}} / \text{slope}_{\text{standards in solvent}}$

The online version of the original article can be found at <https://doi.org/10.1007/s00216-017-0730-y>

✉ You-ning Ma  
11116088@zju.edu.cn

<sup>1</sup> China National Rice Research Institute, No. 28, Rice Road, Hangzhou, Zhejiang 310006, China