




Correction to: Weight changes associated with antiepileptic mood stabilizers in the treatment of bipolar disorder

Koen P. Grootens^{1,2}  · Anna Meijer^{3,4} · Erwin G. Hartong⁵ · Bennard Doornbos⁶ · P. Roberto Bakker^{7,8} · Asmar Al Hadithy⁹ · Kirsten N. Hoogerheide¹⁰ · Frans Overmeire¹² · Radboud M. Marijnissen^{4,11} · Henricus G. Ruhe^{2,4}

Published online: 14 September 2018

© Springer-Verlag GmbH Germany, part of Springer Nature 2018

Correction to: European Journal of Clinical Pharmacology
<https://doi.org/10.1007/s00228-018-2517-2>

In the original version of this article unfortunately two tables have been missing. By mistake they have been published as Supplementary Material. We apologize for any inconvenience caused.

The original article has been corrected.

The online version of the original article can be found at <https://doi.org/10.1007/s00228-018-2517-2>

✉ Koen P. Grootens
k.grootens@reiniervanarkel.nl

¹ Reinier van Arkel, Bethaniestraat 2, 5211 LJ 's-Hertogenbosch, The Netherlands

² Radboud University Medical Centre, Nijmegen, The Netherlands

³ Department of Teacher Education, Faculty of Behavioural and Social Sciences, University of Groningen, Groningen, The Netherlands

⁴ University Medical Centre Groningen, Groningen, The Netherlands

⁵ Department of Psychiatry, Canisius Wilhelmina Hospital Nijmegen, Nijmegen, The Netherlands

⁶ GGZ Drenthe, Assen, The Netherlands

⁷ GGZ Centraal, Amersfoort, The Netherlands

⁸ Maastricht University, Maastricht, The Netherlands

⁹ Pamassia Group, The Hague, The Netherlands

¹⁰ De Jutters, The Hague, The Netherlands

¹¹ Pro Persona, Wolfheze, The Netherlands

¹² VMDB (Dutch association of patients with bipolar disorder and their relatives), Amersfoort, Netherlands