



Remarks on input-to-state stability of collocated systems with saturated feedback

Birgit Jacob¹ · Felix L. Schwenninger^{2,3} · Lukas A. Vorberg¹

Received: 3 January 2020 / Accepted: 24 August 2020 / Published online: 7 September 2020
© The Author(s) 2020

Abstract

We investigate input-to-state stability (ISS) of infinite-dimensional collocated control systems subject to saturated feedback. Here, the unsaturated closed loop is dissipative and uniformly globally asymptotically stable. Under an additional assumption on the linear system, we show ISS for the saturated one. We discuss the sharpness of the conditions in light of existing results in the literature.

Keywords Input-to-state stability · Saturation · Collocated system · Semilinear system · Infinite-dimensional system

1 Introduction

In this note we continue the study of the stability of systems of the form

$$\begin{cases} \dot{x}(t) = Ax(t) - B\sigma(B^*x(t) + d(t)), \\ x(0) = x_0, \end{cases} \quad (\Sigma_{SLD})$$

✉ Birgit Jacob
bjacob@uni-wuppertal.de

Felix L. Schwenninger
f.l.schwenninger@utwente.nl

Lukas A. Vorberg
vorberg@uni-wuppertal.de

¹ School of Mathematics and Natural Sciences, University of Wuppertal, IMACM, Gaußstraße 20, 42119 Wuppertal, Germany

² Department of Applied Mathematics, University of Twente, P.O. Box 217, 7500,AE Enschede, The Netherlands

³ Department of Mathematics, University of Hamburg, Bundesstr. 55, 20146 Hamburg, Germany

derived from the linear collocated open-loop system

$$\begin{aligned}\dot{x}(t) &= Ax(t) + Bu(t), \\ y(t) &= B^*x(t).\end{aligned}$$

by the nonlinear feedback law $u(t) = -\sigma(y(t) + d(t))$. Here X and U are Hilbert spaces, $A : D(A) \subset X \rightarrow X$ is the generator of a strongly continuous contraction semigroup, and B is a bounded linear operator from U to X , i.e. $B \in \mathcal{L}(U, X)$. The function $\sigma : U \rightarrow U$ is locally Lipschitz continuous and maximal monotone with $\sigma(0) = 0$. Of particular interest is the case in which σ is even linear in a neighbourhood of 0. The open-loop system is called collocated as the output operator B^* equals the adjoint of the input operator B . In the following, we are interested in stability with respect to both the initial value x_0 , that is, *internal stability*, and the disturbance d , *external stability*. This is combined in the notion of *input-to-state stability* (ISS), which has recently been studied for infinite-dimensional systems e.g. in [7,9,19,20] and particularly for semilinear systems in [5,6,23], see also [18] for a survey. The effect of feedback laws acting (approximately) linearly only locally is known in the literature as *saturation* and first appeared in [24,25] in the context of stabilization of infinite-dimensional linear systems, see also [10]. There, internal stability of the closed-loop system was studied using nonlinear semigroup theory, a natural tool to establish existence and uniqueness of solutions for equations of the above type, see also the more recent works [11,15,16]. The simultaneous study of internal stability and the robustness with respect to additive disturbances in the saturation seems to be rather recent. This notion clearly includes uniform global (internal) stability, which is far from being trivial for such nonlinear systems. In [22], this was studied for a wave equation, and in [14] Korteweg–de Vries type equation was rigorously discussed, building on preliminary works in [12,13], see also [11].

The combination of saturation and ISS was initiated in [15] and, as for internal stability, complemented in [16]. For the rich finite-dimensional theory on ISS for related semilinear systems, we refer e.g. to [5,6] and the references therein. For (infinite-dimensional) nonlinear systems, ISS is typically assessed by Lyapunov functions, see e.g. [3,8,17,20,23]. These are often constructed by energy-based L^2 norms, but also Banach space methods exist [20], which are much easier to handle in the sense of L^∞ -estimates as present in ISS. We will use some of these constructions here.

In this note, we investigate the question whether internal stability of the linear undisturbed system, that is, (Σ_{SLD}) with $\sigma(u) = u$ and $d \equiv 0$, implies input-to-state stability of (Σ_{SLD}) . In doing so, we try to shed light on limitations of existing results. Because the linear system has a bounded input operator, the above question is equivalent to asking whether ISS of the linear system yields that (Σ_{SLD}) is ISS, see e.g. [9]. For nonlinear systems, uniform global asymptotic (internal) stability is only a necessary condition for ISS, which, however, may fail in the presence of saturation. Indeed, the following saturated transport equation will serve as a model for a counterexample which we shall discuss in this note in detail, see Theorem 7,

$$\begin{cases} \dot{x}(t, \xi) = \frac{d}{d\xi}x(t, \xi) - \text{sat}_{\mathbb{R}}(x(t, \xi)), & (t, \xi) \in (0, \infty) \times [0, 1], \\ x(t, 0) = x(t, 1), \\ x(0, \xi) = f(\xi), \end{cases} \quad (\Sigma_{\text{sat}})$$

where

$$\text{sat}_{\mathbb{R}}(z) := \begin{cases} \frac{z}{|z|}, & |z| \geq 1 \\ z, & z \in (-1, 1). \end{cases} \quad (1)$$

2 ISS for saturated systems

Definition 1 We call $\sigma : U \rightarrow U$ an *admissible feedback function* if

- (i) $\sigma(0) = 0$,
- (ii) σ is *locally Lipschitz continuous*, i.e. for every $r > 0$ there exists a $k_r > 0$ such that

$$\|\sigma(u) - \sigma(v)\|_U \leq k_r \|u - v\|_U \quad \forall u, v \in U \text{ with } \|u\|_U, \|v\|_U \leq r,$$

- (iii) σ is *maximal monotone*, i.e. $\Re\langle \sigma(u) - \sigma(v), u - v \rangle_U \geq 0 \quad \forall u, v \in U$.

If additionally a Banach space S is continuously, densely embedded in U with dual space S' such that

- (iv) $\|\sigma(u) - u\|_{S'} \leq \Re\langle \sigma(u), u \rangle_U \quad \forall u \in U$, and
- (v) there exists $C_0 > 0$ such that

$$\Re\langle u, \sigma(u + v) - \sigma(u) \rangle_U \leq C_0 \|v\|_U \quad \forall u, v \in U,$$

then we call σ a *saturation function*. Here $U \subset S'$ is understood in the sense of rigged Hilbert spaces, i.e. an element u in U is identified with the functional $s \mapsto \langle s, u \rangle_U$ in S' .

It seems that the notion of a saturation function appeared first in the context of infinite-dimensional systems in [24,25]. Note that the precise definition—in particular which properties it should include—has varied in the literature since then. Our definition here matches the one in [15], except for the fact that, in addition, it is required that $\|\sigma(u)\|_S \leq 1$. We distinguish between “admissible feedback functions” and “saturation functions” in order to point out which (minimal) assumptions are needed in the following results.

Example 2 Let $\text{sat}_{\mathbb{R}}$ be the function from (1). It is easy to see that the function

$$\text{sat} : L^2(0, 1) \rightarrow L^2(0, 1), \quad u \mapsto \text{sat}_{\mathbb{R}}(u(\cdot))$$

is an admissible feedback function. Moreover, for $S = L^\infty(0, 1)$ we have that

$$\begin{aligned} \|\text{sat}(u) - u\|_{L^1(0,1)} &= \int_0^1 |\text{sat}(u)(\xi) - u(\xi)| \, d\xi \\ &\leq \int_{\{u \geq 1\}} u(\xi) \, d\xi + \int_{\{-1 \leq u \leq 1\}} u^2(\xi) \, d\xi + \int_{\{u \leq -1\}} -u(\xi) \, d\xi \\ &= \langle \text{sat}(u), u \rangle_U \quad \forall u \in U. \end{aligned}$$

As Property (v) from Definition 1 follows similarly, sat is a saturation function. Note that this example is well known in the literature, see [15, 16] and the references therein.

Let σ be an admissible feedback function. In the rest of the paper, we will be interested in the following two types of systems: The *unsaturated system*,

$$\begin{cases} \dot{x}(t) = Ax(t) - BB^*x(t), \\ x(0) = x_0, \end{cases} \tag{\Sigma_L}$$

and the *disturbed saturated system*

$$\begin{cases} \dot{x}(t) = Ax(t) - B\sigma(B^*x(t) + d(t)), \\ x(0) = x_0. \end{cases} \tag{\Sigma_{SLD}}$$

with $d \in L^\infty(0, \infty; U)$. We abbreviate

$$\tilde{A} : D(\tilde{A}) \subset X \rightarrow X, \quad \tilde{A}x := Ax - BB^*x.$$

By the Lumer–Phillips theorem, \tilde{A} generates a strongly continuous semigroup of contractions $(\tilde{T}(t))_{t \geq 0}$ as $-BB^* \in L(X)$ is dissipative. Moreover, the nonlinear operator $A - B\sigma(B^*\cdot)$ generates a nonlinear semigroup of contractions [26, Thm. 1] since, obviously, $B\sigma(B^*\cdot) : X \rightarrow X$ is continuous and monotone, i.e.

$$\langle B\sigma(B^*x) - B\sigma(B^*y), x - y \rangle \geq 0, \quad \forall x, y \in X.$$

Clearly, (Σ_L) is a special case of (Σ_{SLD}) with $d = 0$, as $\sigma(u) = u$ is an admissible feedback function.

Definition 3 Let $x_0 \in X$, $d \in L^\infty_{\text{loc}}(0, \infty; U)$ and $t_1 > 0$. A continuous function $x : [0, t_1] \rightarrow X$ satisfying

$$x(t) = T(t)x_0 - \int_0^t T(t-s)B\sigma(B^*x(s) + d(s)) \, ds, \quad t \in [0, t_1],$$

is called a *mild solution* of (Σ_{SLD}) on $[0, t_1]$, and we may omit the reference to the interval. If $x : [0, \infty) \rightarrow X$ is such that the restriction $x|_{[0,t_1]}$ is a mild solution for every $t_1 > 0$, then x is called a *global mild solution*.

By our assumptions, (Σ_{SLD}) has a unique mild solution (on some maximal interval) for any $x_0 \in X$ and $d \in L^\infty(0, \infty; U)$, [21, Thm. 6.1.4]¹. In order to introduce the external stability notions, the following well-known comparison functions are needed,

$$\begin{aligned} \mathcal{K} &:= \{\alpha \in C(\mathbb{R}_+, \mathbb{R}_+) \mid \alpha \text{ is strictly increasing, } \alpha(0) = 0\}, \\ \mathcal{K}_\infty &:= \{\alpha \in \mathcal{K} \mid \alpha \text{ is unbounded}\}, \\ \mathcal{L} &:= \{\alpha \in C(\mathbb{R}_+, \mathbb{R}_+) \mid \alpha \text{ is strictly decreasing with } \lim_{t \rightarrow \infty} \alpha(t) = 0\}, \\ \mathcal{KL} &:= \{\beta \in C(\mathbb{R}_+ \times \mathbb{R}_+, \mathbb{R}_+) \mid \beta(\cdot, t) \in \mathcal{K} \ \forall t > 0, \beta(r, \cdot) \in \mathcal{L} \ \forall r > 0\}, \end{aligned}$$

where $C(\mathbb{R}_+, \mathbb{R}_+)$ refers to the continuous functions from \mathbb{R}_+ to \mathbb{R}_+ .

- Definition 4** (i) (Σ_{SLD}) is called *globally asymptotically stable (GAS)* if every mild solution x for $d = 0$ is global and the following two properties hold; $\lim_{t \rightarrow \infty} \|x(t)\|_X = 0$ for every initial condition $x_0 \in X$ and there exist $\sigma \in \mathcal{K}_\infty$ and $r > 0$ such that $\|x(t)\| \leq \sigma(\|x_0\|)$ for every $x_0 \in X$ with $\|x_0\| \leq r$, $d = 0$ and $t \geq 0$.
- (ii) (Σ_{SLD}) is called *semi-globally exponentially stable in $D(A)$* if for $d = 0$ and any $r > 0$ there exist $\mu(r) > 0$ and $K(r) > 0$ such that any mild solution x with initial value $x_0 \in D(A)$ is global and satisfies

$$\|x(t)\|_X \leq K(r)e^{-\mu(r)t} \|x_0\|_X \quad \forall t \geq 0$$

for $\|x_0\|_{D(A)} := \|x_0\|_X + \|Ax_0\|_X \leq r$.

- (iii) (Σ_{SLD}) is called *locally input-to-state stable (LISS)* if there exist $r > 0$, $\beta \in \mathcal{KL}$ and $\rho \in \mathcal{K}_\infty$ such that every mild solution x with initial value satisfying $\|x_0\|_X \leq r$ and disturbance d with $\|d\|_{L^\infty(0, \infty; U)} \leq r$ is global and for all $t \geq 0$ we have that

$$\|x(t)\|_X \leq \beta(\|x_0\|_X, t) + \rho(\|d\|_{L^\infty(0, t; U)}). \tag{2}$$

(Σ_{SLD}) is called *input-to-state stable (ISS)* if $r = \infty$.

System (Σ_{SLD}) is called LISS with respect to $C(0, \infty; U)$ if the above holds for continuous disturbances only. If (2) holds for (Σ_{SLD}) with $d \equiv 0$ and $r = \infty$, the system is called *uniformly globally asymptotically stable (UGAS)*, where the uniformity is with respect to the initial values.

Note that in our notation ‘‘UGAS’’ refers to ‘‘0-UGAS’’ and ‘‘GAS’’ refers to ‘‘0-GAS’’ more commonly used in the literature. The System (Σ_{SLD}) is globally asymptotically stable if and only if for every mild solution x for $d = 0$ we have $\lim_{t \rightarrow \infty} \|x(t)\|_X = 0$. This directly follows from the fact that the mild solutions of (Σ_{SLD}) with $d = 0$ can be represented by a (nonlinear) contraction semigroup, which implies that $\|x(t)\| \leq \|x_0\|$ for all $t \geq 0$, $x_0 \in X$. Compared to the other notions, semi-global exponential stability in $D(A)$ seems to be less common in the

¹ A careful look at the proof reveals that the continuity of the nonlinearity in t required in [21, Thm. 6.1.2] can be dropped in our setting.

literature, but appeared already in the context of saturated systems in [16]. The notion of semi-global exponential stability in X was studied in [14]. Note that for the linear System (Σ_L) UGAS is equivalent to the existence of constants $M, \omega > 0$ such that $\|\tilde{T}(t)\|_X \leq Me^{-\omega t}$ for all $t \geq 0$, see [4, Proposition V.1.2]. Clearly, if (Σ_{SLD}) is UGAS, then it is globally asymptotically stable. We note that semi-global exponential stability in $D(A)$ implies global asymptotical stability since $D(A)$ is dense in X and by the above-mentioned fact that the mild solutions are described by a nonlinear contraction semigroup. Moreover, using again the denseness of $D(A)$ in X , the System (Σ_L) is UGAS if and only if it is semi-globally exponentially stable in $D(A)$.

Next we investigate the question whether (semi-)global exponential stability in $D(A)$ or UGAS of System (Σ_L) implies (semi-)global exponential stability in $D(A)$ or UGAS of System (Σ_{SLD}) .

In [11, Theorem 2], it is shown that global asymptotic stability of (Σ_L) implies global asymptotic stability of (Σ_{SLD}) if

- $D(A)$ equipped with the norm $\|\cdot\|_{D(A)} = \|\cdot\|_X + \|A \cdot\|_X$ is a Banach space compactly embedded in X and
- σ is an admissible feedback function with the additional properties that for all $u \in U, \Re\langle u, \sigma(u) \rangle = 0$ implies $u = 0$.

Note that the other assumptions of [11, Theorem 2] are satisfied in our situation if σ is globally Lipschitz; this follows again by the fact that the mild solutions are represented by a nonlinear semigroup. In [19, Section V], it is shown that under these conditions and in finite dimensions, i.e. $X = \mathbb{R}^n$ and $U = \mathbb{R}^m$, (Σ_{SLD}) is UGAS.

Here we are interested in results for general admissible feedback functions and saturation functions. The following result was proved in [16] and [15].

Proposition 5 [[15, Theorem 1], [16, Theorem 2]] *Let (Σ_L) be UGAS and $\sigma : U \rightarrow U$ be a globally Lipschitz saturation function.*

- (i) *If $S = U$, then (Σ_{SLD}) is ISS.*
- (ii) *If there exists a bounded self-adjoint operator P which maps $D(A)$ to $D(A)$ and solves*

$$\langle \tilde{A}x, Px \rangle + \langle Px, \tilde{A}x \rangle \leq -\langle x, x \rangle, \quad \forall x \in D(\tilde{A}) = D(A), \tag{3}$$

and if

$$\exists c > 0 \forall x \in D(A) : \|B^*x\|_S \leq c\|x\|_{D(A)}, \tag{4}$$

then (Σ_{SLD}) is semi-globally exponentially stable in $D(A)$.

Note that in the second part of Proposition 5, the existence of a bounded, self-adjoint operator P satisfying (3) always follows from the assumption that (Σ_L) is UGAS. However, the property that such P leaves $D(\tilde{A})$ invariant does not hold in general. For instance, this is satisfied if there exists $\alpha > 0$ such that $\Re\langle Ax, x \rangle \leq -\alpha\|x\|^2$ all $x \in D(A)$, which follows directly from dissipativity. On the other hand, it is not hard to construct examples where this invariance is not satisfied. We will comment on this

condition also in Remark 10(ii). We will show next that Proposition 5(ii) does not hold without assuming (4) and, moreover, that (4) does neither imply UGAS nor ISS for (Σ_{SLD}) .

Proposition 6 *Let $X = U = L^2(0, 1)$, $S = L^\infty(0, 1)$, $A = 0$, $B = I$ and $\sigma = \text{sat}$. Then, System (Σ_L) is UGAS and System (Σ_{SLD}) is neither semi-globally exponentially stable in $D(A)$, nor UGAS nor ISS.*

Proof As System (Σ_L) is given by $\dot{x}(t) = -x(t)$, it is UGAS. System (Σ_{SLD}) is given by

$$\begin{cases} \dot{x}(t, \xi) = -\text{sat}_{\mathbb{R}}(x(t, \xi)), & t \geq 0, \xi \in (0, 1), \\ x(0, \xi) = f(\xi), \end{cases} \tag{5}$$

with the unique mild solution $x \in C([0, \infty); L^2(0, 1))$

$$x(t, \xi) = \begin{cases} f(\xi) - t, & \text{if } f(\xi) \geq 1 + t, \\ e^{-t} f(\xi), & \text{if } f(\xi) \in (-1, 1), \\ f(\xi) + t, & \text{if } f(\xi) \leq -1 - t, \\ e^{f(\xi)-1-t}, & \text{if } f(\xi) \in [1, 1 + t), \\ -e^{1-t-f(\xi)}, & \text{if } f(\xi) \in (-1 - t, -1], \end{cases} \tag{6}$$

which can be derived by solving (5) for fixed ξ as simple ODE. We will show that there exists a sequence $(f_n)_n \in L^2(0, 1)$ with $\|f_n\|_{D(A)} = \|f_n\|_{L^2(0,1)} = 1$ such that for all $t > 0$ there exists an $n \in \mathbb{N}$ such that $\|x_n(t)\|_{L^2(0,1)} > \frac{1}{2}$ where x_n denotes the corresponding solution of (5) with initial function f_n . For this purpose, we will only consider the restriction of x_n to $\{\xi \in [0, 1] \mid f(\xi) \geq 1 + t\}$ and define

$$f_n(\xi) := \frac{1}{\sqrt{n}} \xi^{-\alpha_n}$$

with $\alpha_n := \frac{1}{2} \left(1 - \frac{1}{n}\right)$. Clearly, $f_n \in L^2(0, 1)$, $\|f_n\|_{L^2} = 1$ and f_n is decreasing. Note that the equation $f_n(\xi) = 1 + t$ has a unique solution ξ for fixed n and t which is given by

$$\xi = \xi_{t,n} := \frac{1}{(\sqrt{n}(1+t))^{\frac{1}{\alpha_n}}}.$$

Therefore, $\{\xi \in [0, 1] \mid f_n(\xi) \geq 1 + t\} = \{\xi \in [0, 1] \mid \xi \leq \xi_{t,n}\}$. Hence,

$$\begin{aligned} \|x_n(t)\|_{L^2(0,1)}^2 &\geq \int_0^{\xi_{t,n}} x_n(t, \xi)^2 d\xi \\ &= \int_0^{\xi_{t,n}} (f_n(\xi) - t)^2 d\xi \end{aligned}$$

$$\begin{aligned}
 &= \int_0^{\xi_{t,n}} \left(\frac{1}{\sqrt{n}} \xi^{-\alpha_n} - t \right)^2 d\xi \\
 &= \frac{1}{n} \int_0^{\xi_{t,n}} \xi^{-2\alpha_n} d\xi - \frac{2t}{\sqrt{n}} \int_0^{\xi_{t,n}} \xi^{-\alpha_n} d\xi + \int_0^{\xi_{t,n}} t^2 d\xi \\
 &= \frac{1}{n} \frac{1}{1-2\alpha_n} \xi_{t,n}^{1-2\alpha_n} - \frac{2t}{\sqrt{n}} \frac{1}{1-\alpha_n} \xi_{t,n}^{1-\alpha_n} + t^2 \xi_{t,n} \\
 &= n^{\frac{1}{1-n}} (1+t)^{\frac{2}{1-n}} - \frac{1}{n+1} 2n^{\frac{1}{1-n}} 2t(1+t)^{\frac{1+n}{1-n}} + n^{\frac{n}{1-n}} t^2 (1+t)^{\frac{2n}{1-n}}.
 \end{aligned}$$

Taking the limit $n \rightarrow \infty$, we conclude

$$\lim_{n \rightarrow \infty} \|x_n(t)\|_{L^2(0,1)}^2 \geq 1.$$

Thus, the solution of System (5) does not converge uniformly to 0 with respect to the norm or graph norm of the initial value, so the system is neither semi-globally exponentially stable in $D(A)$ nor UGAS. \square

Note that System (Σ_{SLD}) from Proposition 6 is (GAS) for $d = 0$ by [11, Theorem 2]. After we have seen that (4) is necessary to conclude semi-global exponential stability in $D(A)$ in Proposition 5(ii), one may ask whether “more stability” can in fact be expected. The following theorem shows that UGAS of System (Σ_L) together with the hypotheses in Proposition 5(ii) is not sufficient to guarantee UGAS of System (Σ_{SLD}) .

Theorem 7 *Let $X = U = L^2(0, 1)$, $B = I$, $S = L^\infty(0, 1)$, $\sigma = \text{sat}$ and*

$$A = \frac{d}{d\xi}, \quad D(A) = \{y \in H^1(0, 1) \mid y(0) = y(1)\}.$$

Then, the following assertions hold.

- (i) *System (Σ_L) is UGAS and the hypothesis of Proposition 5(ii) holds,*
- (ii) *System (Σ_{SLD}) is semi-globally exponentially stable in $D(A)$,*
- (iii) *System (Σ_{SLD}) is neither UGAS nor ISS.*

We note that System (Σ_{SLD}) of Theorem 7 equals (Σ_{sat}) . Further, in [15, Thm. 1] it has been wrongly stated that the saturated system is UGAS.

Proof It is easy to see that System (Σ_L) is UGAS. Since A is dissipative, it follows that $P = I$ solves (3) for $\tilde{A} = A - BB^* = A - I$. Trivially, P maps $D(A)$ to $D(A)$. Condition (4) is satisfied because $H^1(0, 1)$ is continuously embedded in $L^\infty(0, 1)$. Hence, (Σ_{SLD}) is semi-globally exponentially stable in $D(A)$ by Proposition 5 and the fact that σ is globally Lipschitz continuous. This shows Assertions (i) and (ii). To see (iii), note that A generates the periodic shift semigroup on $L^2(0, 1)$. By extending the initial function f periodically to \mathbb{R}_+ , the unique mild solution $y \in C([0, \infty); L^2(0, 1))$ of (Σ_{SLD}) is given by

$$y(t, \xi) = x(t, \xi + t),$$

where x is defined in (6). By the particular form of (6), this implies that

$$\|x(t)\|_{L^2(0,1)} = \|y(t)\|_{L^2(0,1)}$$

holds for all $t \geq 0$. We can therefore choose the same sequence $(f_n)_n \in L^2(0, 1)$ with $\|f_n\|_{L^2(0,1)} = 1$ as in the proof of Proposition 6 in order to conclude

$$\lim_{n \rightarrow \infty} \|y_n(t)\|_{L^2(0,1)}^2 \geq 1.$$

This shows that System (Σ_{SLD}) is not UGAS and thus not ISS. □

An important tool for the verification of ISS of System (Σ_{SLD}) are ISS Lyapunov functions.

Definition 8 Let $U_r = \{x \in X : \|x\| \leq r\}$ and $r \in (0, \infty]$. Let \mathcal{U} be either $C(0, \infty; U)$ or $L^\infty_{loc}(0, \infty; U)$. A continuous function $V : U_r \rightarrow \mathbb{R}_{\geq 0}$ is called an *LISS Lyapunov function* for (Σ_{SLD}) with respect to \mathcal{U} , if there exists $\psi_1, \psi_2, \alpha, \rho \in \mathcal{K}_\infty$, such that for all $x_0 \in U_r, d \in \mathcal{U}, \|d\|_{L^\infty(0,\infty;U)} \leq r$,

$$\psi_1(\|x_0\|_X) \leq V(x_0) \leq \psi_2(\|x_0\|_X)$$

and

$$\dot{V}_d(x_0) := \limsup_{t \searrow 0} \frac{1}{t} (V(x(t)) - V(x_0)) \leq -\alpha(\|x_0\|_X) + \rho(\|d\|_{L^\infty(0,\infty;U)}), \quad (7)$$

where x is the mild solution of (Σ_{SLD}) with initial value x_0 and disturbance d . If $r = \infty$, then V is called an *ISS Lyapunov function*.

Note that our definition of an ISS Lyapunov function corresponds to the one of a “coercive ISS Lyapunov function in dissipative form” in the literature, [18]. By [3, Thm. 1], see also [18, Thm. 2.18], the existence of an (L)ISS Lyapunov implies (L)ISS for a large class of control systems which, in particular, have to satisfy the “boundedness-implies-continuation” property (BIC). System (Σ_{SLD}) with an admissible feedback function and continuous, or, more generally, piecewise continuous disturbances d belongs to this class, which allows to infer (L)ISS from the existence of a Lyapunov function. To see this, note in particular that the (BIC) property is satisfied by classical results on semilinear equations, [1, Prop. 4.3.3] or [21, Thm. 6.1.4]. In the following, we will infer ISS by constructing Lyapunov functions.

Theorem 9 *Suppose that there exists $\alpha > 0$ such that $\|T(t)\| \leq e^{-\alpha t}$ for all $t > 0$ and let σ be an admissible feedback function. Then, the function*

$$V(x) = \|x\|_X^2, \quad x \in X,$$

is an ISS Lyapunov function for (Σ_{SLD}) with respect to $C(0, \infty; U)$ and System (Σ_{SLD}) is ISS with respect to $C(0, \infty; U)$.

Proof Let $x \in C(0, t_1; X)$ be the mild solution of (Σ_{SLD}) with initial value $x_0 \in D(A)$ and disturbance $d \in C(0, \infty; U)$. Let $y \in C(0, t_2; X)$ be the mild solution of the system

$$\begin{cases} \dot{y}(t) = Ay(t) - B\sigma(B^*y(t) + \tilde{d}(t)) \\ y(0) = y_0 \end{cases}$$

with $\tilde{d} \in C(0, \infty; U)$ and $y_0 \in X$. Then there exists an $r > 0$ such that

$$\max\{\|B^*x(s) + d(s)\|_U, \|B^*y(s) + \tilde{d}(s)\|_U, \|B^*x(s)\|_U \mid s \in [0, \min\{t_1, t_2\}]\} < r$$

because x, y, d and \tilde{d} are continuous. Thus, we have for $t \in [0, \min\{t_1, t_2\})$

$$\|x(t) - y(t)\| \leq \|x_0 - y_0\| + \int_0^t \|B\|k_r(\|B\|\|x(s) - y(s)\| + \|d(s) - \tilde{d}(s)\|) ds.$$

Applying Gronwall’s inequality yields

$$\|x(t) - y(t)\| \leq \left(\|x_0 - y_0\| + \int_0^t \|B\|k_r\|d(s) - \tilde{d}(s)\| ds \right) e^{t\|B\|^2k_r}. \tag{8}$$

Let us for a moment assume that d is Lipschitz continuous with Lipschitz constant L . We will prove that x is right differentiable. For $0 < h < t_1 - t$, we can write $x(t + h)$ in the form

$$\begin{aligned} x(t + h) &= T(t + h)x_0 - \int_0^{t+h} T(t + h - s)B\sigma(B^*x(s) + d(s)) ds \\ &= T(t)x(h) - \int_0^t T(t - s)B\sigma(B^*x(s + h) + d(s + h)) ds. \end{aligned}$$

Thus, x at time $t + h$ equals the mild solution y of

$$\begin{cases} \dot{y}(t) = Ay(t) - B\sigma(B^*y(t) + d(t + h)) \\ y(0) = x(h) \end{cases} \tag{9}$$

at time t . Hence, by (8) we obtain

$$\|x(t + h) - x(t)\| \leq (\|x(h) - x_0\| + \|B\|k_rLht)e^{t\|B\|^2k_r}. \tag{10}$$

Note that

$$\frac{x(h) - x_0}{h} = \frac{T(h)x_0 - x_0}{h} - \frac{1}{h} \int_0^h T(h - s)B\sigma(B^*x(s) + d(s)) ds$$

converges to $Ax_0 - B\sigma(B^*x_0 + d(0))$ as $h \searrow 0$ since $x_0 \in D(A)$ and σ, x and d are continuous. Therefore, by (10), we deduce

$$\limsup_{h \searrow 0} \frac{\|x(t+h) - x(t)\|}{h} < \infty. \tag{11}$$

By the definition of the mild solution, we have that

$$\frac{T(h) - I}{h}x(t) = \frac{x(t+h) - x(t)}{h} + \frac{1}{h} \int_t^{t+h} T(t+h-s)B\sigma(B^*x(s) + d(s))ds.$$

Again by continuity of σ, x and d , we have that

$$\lim_{h \searrow 0} \frac{1}{h} \int_t^{t+h} T(t+h-s)B\sigma(B^*x(s) + d(s)) ds = B\sigma(B^*x(t) + d(t)).$$

Combining this with (11) shows that

$$x(t) \in \{z \in X \mid \limsup_{h \searrow 0} \frac{1}{h} \|T(h)x - x\| < \infty\},$$

which means that $x(t)$ is an element of the Favard space of the semigroup, and because X is reflexive, we can conclude that $x(t) \in D(A)$, [4, Cor. II.5.21]. This implies that x is right differentiable at t with

$$\lim_{h \searrow 0} \frac{x(t+h) - x(t)}{h} = Ax(t) - B\sigma(B^*x(t) + d(t)).$$

As $V(x) = \|x\|^2$, we hence obtain for the Dini derivative

$$D^+V(x(\cdot))(t) = \limsup_{h \searrow 0} \frac{1}{t} (V(x(t+h)) - V(x(t)))$$

that

$$\begin{aligned} D^+V(x(\cdot))(t) &= 2\Re(\langle Ax(t), x(t) \rangle_X - \langle B\sigma(B^*x(t) + d(t)), x(t) \rangle_X) \\ &\leq -2\alpha\|x(t)\|^2 - \Re(\langle \sigma(B^*x(t) + d(t)) - \sigma(B^*x(t)), B^*x(t) \rangle_X) \\ &\leq -2\alpha\|x(t)\|^2 + \|\sigma(B^*x(t) + d(t)) - \sigma(B^*x(t))\| \|B^*x(t)\| \\ &\leq -2\alpha\|x(t)\|^2 + k_r\|d(t)\| \|B\| \|x(t)\|, \end{aligned} \tag{12}$$

where we used that $-\Re(\langle \sigma(B^*x), B^*x \rangle) \leq 0$ by Property (i) and (ii) of admissible feedback functions and the local Lipschitz condition for σ . By [2, Cor. A.5.45], we obtain

$$V(x(t+h)) - V(x(t)) \leq \int_t^{t+h} -2\alpha\|x(s)\|^2 + k_r\|d(s)\| \|B\| \|x(s)\| ds. \tag{13}$$

From (8), we derive

$$\|x(t) - y(t)\| \leq (\|x_0 - y_0\| + t\|B\|k_r\|d - \tilde{d}\|_{L^\infty(0,t;U)})e^{t\|B\|^2k_r},$$

and therefore, the mild solution of (Σ_{SLD}) depends continuously on the initial data and the disturbance. Hence, by understanding $x(t + h)$ again as the solution of (9) at time t , (13) holds for all $x_0 \in X$ and $d \in C(0, \infty; U)$ which leads to

$$\begin{aligned} \dot{V}_d(x_0) &\leq -2\alpha\|x_0\|^2 + k_r\|d(0)\| \|B\| \|x_0\| \\ &\leq (\varepsilon - 2\alpha)\|x_0\|^2 + \frac{(k_r \|B\| \|d(0)\|)^2}{\varepsilon} \end{aligned}$$

for all $x_0 \in X$, $d \in C(0, \infty; U)$ and $\varepsilon > 0$. Choosing $\varepsilon < 2\alpha$, this shows that V is an ISS-Lyapunov function for (Σ_{SLD}) which implies that (Σ_{SLD}) is ISS by [18, Thm. 2.18]. □

Remark 10

- (i) Recall that the semigroup generated by A in Theorem 7 was not exponentially stable. Theorem 9 shows that this is not accidental.
- (ii) Note that the assumption on the semigroup made in Theorem 9 is strictly stronger than the condition that $(T(t))_{t \geq 0}$ is an exponentially stable contraction semigroup as can be seen e.g. for a nilpotent shift semigroup on $X = L^2(0, 1)$. It is a simple consequence of the Lumer–Phillips theorem that the following assertions are equivalent for a semigroup $(T(t))_{t \geq 0}$ generated by A and some constant $\omega > 0$.
 - (a) $\Re\langle Ax, x \rangle \leq -\omega\|x\|^2$ all $x \in D(A)$.
 - (b) $\sup_{t > 0} \|e^{\omega t} T(t)\| \leq 1$.
 - (c) $P = \frac{1}{\omega} I$ solves $\Re\langle Ax, Px \rangle \leq -\langle x, x \rangle$, for all $x \in D(A)$.

However, we also remark that the above condition is satisfied for a large class of examples, such as in the case when A is a normal operator.

- (iii) It is natural to ask whether Theorem 9 holds when A is merely assumed to generate an exponentially stable semigroup. However, it is unclear how to use the structural assumptions on σ in the general case. On the other hand, the assumption on the semigroup in Theorem 9 implies that $P = I$ satisfies (3) in Proposition 5(ii).
- (iv) An inspection of the proof shows that Theorem 9 can be generalized to piecewise continuous or regulated functions $d : [0, \infty) \rightarrow U$.

Locally linear admissible feedback functions yield LISS Lyapunov functions.

Theorem 11 *Let (Σ_L) be UGAS with $M, \omega > 0$ such that $\|\tilde{T}(t)\| \leq Me^{-\omega t}$ for all $t \geq 0$ and let σ be an admissible feedback function with $\sigma(u) = u$ for all $\|u\|_U \leq \delta$ and some $\delta > 0$. Then, (Σ_{SLD}) is LISS with Lipschitz continuous LISS Lyapunov function $V(x) := \max_{s \geq 0} \|e^{\frac{\omega}{2}s} \tilde{T}(s)x\|_X$.*

Proof Let $\|x_0\|_X \leq \|B\|^{-1}\delta$ and $r := \max\{\|B^*x(s)\|_U, \|B^*x(s) + d(s)\|_U \mid s \in [0, t]\}$ for some $t > 0$. We can rewrite (Σ_{SLD}) in the form

$$\begin{aligned} \dot{x}(t) &= \tilde{A}x(t) + B(B^*x(t) - \sigma(B^*x(t) + d(t))), \\ x(0) &= x_0. \end{aligned}$$

Hence, the mild solution satisfies

$$x(h) = \tilde{T}(h)x_0 + \int_0^h \tilde{T}(h-s)B(B^*x(s) - \sigma(B^*x(s) + d(s)))ds.$$

Denoting the integral by I_h , we have

$$\begin{aligned} \limsup_{h \searrow 0} \frac{1}{h} \|I_h\|_X &\leq \limsup_{h \searrow 0} \frac{1}{h} \left(\int_0^h M \|B\| \|B^*x(s) - \sigma(B^*x(s))\|_U ds \right. \\ &\quad \left. + \int_0^h M \|B\| \|\sigma(B^*x(s)) - \sigma(B^*x(s) + d(s))\|_U ds \right) \\ &\leq M \|B\| \|B^*x_0 - \sigma(B^*x_0)\|_U + M \|B\| k_r \|d\|_{L^\infty(0,\varepsilon;U)} \\ &= M \|B\| k_r \|d\|_{L^\infty(0,\varepsilon;U)}, \end{aligned}$$

where the continuity of x , the Lipschitz continuity of σ as well as the condition $\sigma(u) = u$ if $\|u\| \leq \delta$ have been used.

With $\|x\| \leq V(x) \leq M\|x\|$ and $V(\tilde{T}(t)x) \leq e^{-\frac{\omega}{2}t}V(x)$ for all $x \in X$ we obtain

$$\begin{aligned} \dot{V}_d(x_0) &= \limsup_{h \searrow 0} \frac{1}{h} (V(\tilde{T}(h)x_0 + I_h) - V(x_0)) \\ &\leq \limsup_{h \searrow 0} \frac{1}{h} (e^{-\frac{\omega}{2}h} - 1) V(x_0) + M \limsup_{h \searrow 0} \frac{1}{h} \|I_h\|_X \\ &\leq -\frac{\omega}{2} \|x_0\|_X + M^2 \|B\| k_r \|d\|_{L^\infty(0,\varepsilon;U)} \end{aligned}$$

for every $\varepsilon > 0$. The Lipschitz continuity of V follows from

$$\begin{aligned} |V(x) - V(y)| &\leq \left| \max_{s \geq 0} \|e^{\frac{\omega}{2}s} \tilde{T}(s)x\| - \max_{s \geq 0} \|e^{\frac{\omega}{2}s} \tilde{T}(s)y\| \right| \\ &\leq \max_{s \geq 0} \|e^{\frac{\omega}{2}s} \tilde{T}(s)(x - y)\| \\ &\leq M \|x - y\|, \end{aligned}$$

for all $x, y \in X$. Applying [17, Theorem 4] yields local input-to-state stability of (Σ_{SLD}) . □

Note that Property (iii) of Definition 1 has not been used in the proof of Theorem 11.

3 Conclusion

In this note we have continued the study of ISS for saturated feedback connections of linear systems. Theorem 7 states that ISS cannot be concluded from uniform exponential stability of the unsaturated closed-loop and stability of the (undisturbed) open-loop linear system

$$\dot{x}(t) = Ax(t)$$

(i.e. the semigroup generated by A is bounded). However, the conclusion does hold under more assumptions on A ; namely, that $\Re \langle Ax, x \rangle \leq -\alpha \|x\|^2$ for some $\alpha > 0$ and all $x \in D(A)$, see Theorem 9. The latter property can be seen as some kind of quasi-contractivity of the semigroup combined with exponential stability. This condition seems to be crucial for the proof, see Remark 10. The question remains whether the result could be generalized to more general semigroups, e.g. such as contractive semigroups which are exponentially stable, but do not satisfy the above mentioned quasi-contractivity. Note, however, that the assumption that A generates a contraction semigroup seems to be essential to employ dissipativity of the nonlinear system.

Another task for future research is the step towards unbounded operators B , prominently appearing in boundary control systems. As our techniques and also the ones used in existing results for ISS on saturated systems seem to heavily rely on the boundedness of B , this may require a different approach or more structural assumptions on A .

Acknowledgements The authors thank Hans Zwart for fruitful discussions on the proof of Proposition 6 during a visit of the third author at the University of Twente.

Funding Open Access funding provided by Projekt DEAL.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

References

1. Cazenave T, Haraux A (1998) An introduction to semilinear evolution equations, Oxford Lecture Series in Mathematics and its Applications, vol 13. The Clarendon Press, Oxford University Press, New York
2. Curtain R, Zwart H (2020) Introduction to infinite-dimensional systems theory: a state-space approach, Texts in applied mathematics, vol 71. Springer, New York, NY
3. Dashkovskiy S, Mironchenko A (2013) Input-to-state stability of infinite-dimensional control systems. *Math Control Signals Syst* 25(1):1–35
4. Engel KJ, Nagel R (2000) One-parameter semigroups for linear evolution equations, Graduate Texts in Mathematics, vol 194. Springer, New York. With contributions by S. Brendle, M. Campiti, T. Hahn, G. Metafuno, G. Nickel, D. Pallara, C. Perazzoli, A. Rhandi, S. Romanelli and R. Schnaubelt

5. Grüne L (1999) Input-to-state stability of exponentially stabilized semilinear control systems with inhomogeneous perturbations. *Syst Control Lett* 38(1):27–35
6. Guiver C, Logemann H (2020) A circle criterion for strong integral input-to-state stability. *Automatica J IFAC* 111:108641
7. Guiver C, Logemann H, Opmeer MR (2019) Infinite-dimensional Lur'e systems: input-to-state stability and convergence properties. *SIAM J Control Optim* 57(1):334–365
8. Jacob B, Mironchenko A, Partington JR, Wirth F (2019) Non-coercive Lyapunov functions for input-to-state stability of infinite-dimensional systems. [arXiv: 1911.01327](https://arxiv.org/abs/1911.01327)
9. Jacob B, Nabiullin R, Partington JR, Schwenninger F (2018) Infinite-dimensional input-to-state stability and Orlicz spaces. *SIAM J Control Optim* 56(2):868–889
10. Lasiecka I, Seidman T (2003) Strong stability of elastic control systems with dissipative saturating feedback. *Syst Control Lett* 48:243–252
11. Marx S, Andrieu V, Prieur C (2017) Cone-bounded feedback laws for m -dissipative operators on Hilbert spaces. *Math Control Signals Syst* 29(4), Art. 18, 32
12. Marx S, Cerpa E, Prieur C, Andrieu V (2015) Stabilization of a linear Korteweg-de Vries equation with a saturated internal control. In: 14th annual European Control Conference (ECC15). Linz, Austria
13. Marx S, Cerpa E, Prieur C, Andrieu V (2016) Global stabilization of a Korteweg-de Vries equation with a distributed control saturated in L_2 -norm. In: 10th IFAC symposium on nonlinear control systems (NOLCOS 2016). Monterey, CA, USA
14. Marx S, Cerpa E, Prieur C, Andrieu V (2017) Global stabilization of a Korteweg-de Vries equation with saturating distributed control. *SIAM J Control Optim* 55(3):1452–1480
15. Marx S, Chitour Y, Prieur C (2018) Stability results for infinite-dimensional linear control systems subject to saturations. In: 16th European control conference (ECC 2018). Limassol, Cyprus
16. Marx S, Chitour Y, Prieur C (2020) Stability analysis of dissipative systems subject to nonlinear damping via Lyapunov techniques. *IEEE Trans Autom Control* 65(5):2139–2146
17. Mironchenko A (2016) Local input-to-state stability: characterizations and counterexamples. *Syst Control Lett* 87:23–28
18. Mironchenko A, Prieur C (2019) Input-to-state stability of infinite-dimensional systems: recent results and open questions. [arXiv: 1910.01714](https://arxiv.org/abs/1910.01714)
19. Mironchenko A, Wirth F (2018) Characterizations of input-to-state stability for infinite-dimensional systems. *IEEE Trans Automat Control* 63(6):1602–1617
20. Mironchenko A, Wirth F (2018) Lyapunov characterization of input-to-state stability for semilinear control systems over Banach spaces. *Syst Control Lett* 119:64–70
21. Pazy A (1983) Semigroups of linear operators and applications to partial differential equations, *Applied Mathematical Sciences*, vol 44. Springer, New York
22. Prieur C, Tarbouriech S, Gomes da Silva JM (2016) Wave equation with cone-bounded control laws. *IEEE Trans Automat Control* 61(11):3452–3463
23. Schwenninger FL (2020) Input-to-state stability for parabolic boundary control: Linear and semi-linear systems. In: Kerner J, Laasri H, Mugnolo D (eds) *Control theory of infinite-dimensional systems*. Birkhäuser, 83–116
24. Seidman TI, Li H (2001) A note on stabilization with saturating feedback. *Discrete Contin Dyn Syst* 7(2):319–328
25. Slemrod M (1989) Feedback stabilization of a linear control system in Hilbert space with an a priori bounded control. *Math Control Signals Syst* 2(3):265–285
26. Webb GF (1972) Continuous nonlinear perturbations of linear accretive operators in Banach spaces. *J Funct Anal* 10:191–203