



# The choice of climate metric is of limited importance when ranking options for abatement of near-term climate forcers

Stefan Åström<sup>1,2</sup>  · Daniel J. A. Johansson<sup>1</sup>

Received: 18 October 2018 / Accepted: 26 March 2019 / Published online: 2 May 2019  
© The Author(s) 2019

## Abstract

The practice of using climate metrics to estimate carbon dioxide equivalent emissions has long been subject to scientific discussion. One strand of this literature has analysed whether the choice of metric affects the relative cost-effectiveness of options for climate change abatement, but there has been little discussion on the effect of metric choices on cost-effective abatement of near-term climate forcers (NTCFs). These NTCFs are air pollutants primarily regulated by policies outside the climate policy arena and their estimated carbon dioxide equivalent emissions are not typically considered in the evaluation of cost-effective abatement. However, the attention to NTCFs as climate forcers has increased during the last decade. The objective of this paper is to identify whether the relative cost-effectiveness of different NTCF abatement options is robust to climate metric choices. We assess nine plausible NTCF abatement options available in Sweden (with negligible effect on long-lived GHG emissions) and evaluate the robustness of the ranking of these, according to their estimated cost-effectiveness. Different metric designs are considered as well as climate impact uncertainty of NTCFs, with corresponding uncertainty in metric values. The results indicate that the choice of metric has little effect on the ranking of the options according to their cost-effectiveness, with options affecting  $\text{NO}_x$  indicated as an exception. This suggests that the choice of metric utilised when calculating cost-effectiveness of NTCF abatement options is likely to have minor effect on which NTCF abatement options should be targeted for policy interventions (if  $\text{NO}_x$  emissions are not significantly affected).

---

**Electronic supplementary material** The online version of this article (<https://doi.org/10.1007/s10584-019-02427-4>) contains supplementary material, which is available to authorized users.

✉ Stefan Åström  
astrom@chalmers.se

<sup>1</sup> Department of Space, Earth, and Environment, Division of Physical Resource Theory, Chalmers University of Technology, SE-412 93 Gothenburg, Sweden

<sup>2</sup> IVL Swedish Environmental Research Institute Ltd., P.O. Box 530 21, SE-400 14 Gothenburg, Sweden

## 1 Introduction

It is well established that global anthropogenic emissions of long-lived greenhouse gases (LLGHG) such as carbon dioxide (CO<sub>2</sub>) and nitrous oxide (N<sub>2</sub>O) are causing global climate change. Therefore, these emissions are governed in the United Nations Framework Convention on Climate Change (UNFCCC) since 1992. More recently, the fact that also emissions of certain air pollutants have significant climate impact has been gaining widespread international policy recognition for example through the publication of the UNEP synthesis report on near-term climate protection (Kuylenstierna et al. 2011; Shindell et al. 2012) and creation of the Climate and Clean Air Coalition (CCAC) in 2012. These air pollutants are generally referred to as near-term climate forcers (NTCFs) and include sulphur dioxide (SO<sub>2</sub>), black carbon (BC), organic carbon (OC), non-methane volatile organic compounds (NMVOC), nitrogen oxides (NO<sub>x</sub>), and carbon monoxide (CO). Also, the greenhouse gas methane (CH<sub>4</sub>) is usually included in the NTCF group due to its relatively short atmospheric perturbation time of about a decade (Aamaas et al. 2016), but also in the LLGHG group due to being long-lived enough to be well-mixed in the atmosphere. Increased control of NTCF can reduce the rate of global temperature increase, and the peak temperature, contingent that LLGHGs are stringently and perpetually controlled (Bowerman et al. 2013; Shoemaker et al. 2013).

It is complicated to estimate the climate impact of a given NTCF abatement option or regulation. To use large scale climate models costs time and money and requires an effort that is beyond the capacity of most climate policy studies, including NTCF abatement studies (Aamaas et al. 2013). So to estimate climate impact of NTCF emission policies, researchers in several fields and policy makers typically utilise climate metrics<sup>1</sup> (such as global warming potential (GWP), global temperature potential (GTP) (Myhre et al. 2013b), regional temperature potential (RTP) (Aamaas et al. 2016)) normalised against the climate impact of CO<sub>2</sub>. This practice allows for quick estimates of the climate impact—or rather carbon dioxide equivalent (CO<sub>2eq</sub>) emissions—that LLGHG and NTCF emissions give rise to (Schmale et al. 2014).

Today, dozens of climate metrics exist, including economic (Johansson 2011) as well as physical metrics (Aamaas et al. 2013). Given the numerous metrics available—representing different aspects of climate change and/or time horizons—and that researchers and policy makers are using metrics, it is only natural that there has been a discussion about which metric to use for climate and NTCF policy.

The discussion on which metric to use when placing different climate forcers on a common scale has mainly focused on metrics for comparing CO<sub>2</sub> equivalence of emissions that are included in the national reporting to UNFCCC (Manne and Richels 2001; Manning and Reisinger 2011; Peters et al. 2011), as well as combined LLGHG and NTCF abatement (Fesenfeld et al. 2018; Lelvasseur et al. 2016; Tanaka et al. 2010). Generally, these discussions indicate that the choice of metric design depends on the type of climate change impact considered, on time horizon, and on whether future climate impact is discounted, all normative choices (Tanaka et al. 2010). A recent addition to the discussion includes a consistency check regarding time horizon and discount rate (Sarofim and Giordano 2018). But such a discussion is also relevant for the comparison of cost-effective abatement of only NTCF emissions.

<sup>1</sup> Climate metrics present numerical indicator values of one indicator of the climate system from a given amount of emissions, commonly in relation to the same indicator of the climate system of a corresponding amount of a reference gas (usually CO<sub>2</sub>).

However, while NTCF abatement cost studies utilise different climate metrics when estimating cost-effectiveness, we have not identified any studies that analyse the impact of metric choice on the estimated cost-effectiveness of different NTCF abatement options. For example, the Norwegian Environment Agency (2014) used Global Temperature Potential with a 10-year time horizon ( $GTP_{10}$ ) when analysing a Norwegian action plan for abatement of NTCF, while UNEP (Kuylenstierna et al. 2011) used  $GWP_{100}$  when analysing NTCF abatement on a global scale, but none of the studies analyse the sensitivity of their findings with respect to metric choice.

The situation is made further complicated for several reasons. Not only is there a large plethora of metrics that could be applied, but the metric value for a given metric choice is uncertain since the climate impacts of NTCFs are short-lived, causing the climate effect to be heterogeneous in space and time. Due to the heterogeneity, the natural scientific oriented literature on metrics contains recommendations to adapt metrics to regional and seasonal origin of emissions when estimating  $CO_2$  equivalence of NTCF emissions (Aamaas et al. 2016; Henze et al. 2012; Shindell and Faluvegi 2009), and it has been recognised that climate metrics need to be ‘robust enough’ to allow for a ranking in terms of cost-effectiveness of abatement options when the analysis is to be used for policy purposes (Aamaas et al. 2016).

Studies on the effect of metric choice for NTCF abatement would add to the current knowledge on NTCF policy analysis since some aspects of NTCFs are unique in relation to LLGHG, primarily the current policy context. NTCF emissions are often regulated as a group separate from LLGHG emissions in international regulations such as the UNECE Air Convention and the EU National Emission Ceilings Directive.

All in all, the current situation is that (1) several governments and international bodies are interested in controlling NTCF emissions; (2) cost-effective regulations and abatement options are preferred; (3) climate metrics are often used to estimate the impact of options; (4) the existing literature have not analysed the effect of metric choice on cost-effectiveness of NTCF-only abatement options; and (5) most regulations and policies that regulate NTCF emissions do not regulate LLGHGs. Therefore, in this paper, we analyse whether the relative cost-effectiveness of NTCF abatement options (i.e. ranking) is affected by the choice of climate metric. We apply the assessment approach to NTCF abatement options estimated to be available in Sweden.

## 2 Data and method

Section 2 presents the data collected and the method used during the analysis, as well as a description of the sensitivity analysis. The analysis is done in Python by using Sublime and the code is available in the supplementary material 1.

### 2.1 Data selection

#### 2.1.1 Abatement options

In the literature, there is some agreement that small scale wood combustion, mobile machinery, road transport, solvent use, and agriculture are important sources of emissions to and precursors to NTCFs in economically developed countries, including Sweden (ACAP 2014; AMAP et al. 2008; Arctic Council 2011; Kindbom et al. 2015; Kuylenstierna et al. 2011; Stohl et al.

2015; UNEP and WMO 2011; Zaelke et al. 2012). Data on emission abatement options in Sweden is selected based on the criteria that the literature contains estimates on both abatement costs as well as effects on emissions and includes the emission sources mentioned above. Given the limited literature on NTCF abatement options that includes the necessary information, we allow the estimates to vary with respect to target year for the options (between 2020 and 2030). Based on reports on Swedish abatement options for these emitting sectors, we investigate the cost-effectiveness of reducing NTCF emissions through shifting from solid wood to wood pellets in small scale wood combustion (option *Pellets*) (CLEO 2014) or through investment in newer stoves, boilers, and fire places (*Mod. boiler*) (Amann 2014; CLEO 2014). We also include estimates on reducing methane emissions from the agricultural sector through increased use of fermentation of manure (*Man. ferm*) (Hellstedt et al. 2014; Swedish Board of Agriculture 2012), through covering of liquid slurry tanks (*Sl. cover*) (Swedish Board of Agriculture 2012) or through acidification of liquid slurry (*Sl. acid*) (Swedish Board of Agriculture 2012). Furthermore, we include options to reduce NTCF emissions from non-road mobile machinery through a rejuvenation of the machine stock (*Mod. NRMM*) through a shift from two-stroke to four-stroke engines in snow mobiles (*4S S-mob*) (Fridell and Åström 2009), or through a shift to four-stroke engines and electric engines for smaller machinery (*4S&EI NRMM*) (Fridell and Åström 2009). Finally, we include the option of increasing the shift to water-based solvents in products to reduce NMVOC emissions (*W. solv*) Amann et al. (2014). The reported average annual abatement cost per option implemented at its full scale in Sweden is converted from the reported costs and currencies to €<sub>2010</sub> values (Table 1).

These options are reported to have an effect on up to six pollutants with climate impact: three types of fine particulate matter with an aerodynamic diameter of < 2.5 μm (BC, OC, and

**Table 1** Average estimated annual costs of emission abatement for the options considered in this paper

Abatement option	Abbreviation	Average abatement cost per option [M€ <sub>2010</sub> /year]	Reference
More pellets in household wood combustion than expected	Pellets	16.1	CLEO (2014)
More new wood stoves/boilers/fireplaces in households	Mod. boiler	4.3	Amann (2014); CLEO (2014)
Fermentation of animal manure	Man. ferm.	32.1	Hellstedt et al. (2014); Swedish Board of Agriculture (2012)
Covering of liquid slurry storage tank	Sl. cover	3.2	Swedish Board of Agriculture (2012)
Acidification of liquid slurry	Sl. acid.	8.3	Swedish Board of Agriculture (2012)
Younger non-road mobile machinery stock	Mod. NRMM	145.0	Åström et al. (2013)
Shift from two-stroke to four-stroke engines in snow mobiles	4S S-mob.	0.1	Fridell and Åström (2009)
Shift from two-stroke engines to four-stroke engines and electric household small non-road mobile machinery	4S&EI NRMM	3.9	Fridell and Åström (2009)
Product modification of products projected to contain solvents in 2030	W. solv.	180.0	Amann et al. (2014)

remaining sub-fractions ( $PM_{res}$ );  $NO_x$ ;  $CH_4$ ; and NMVOC (Table 2). The options are in the source material not considered to significantly affect fossil fuel demand, which implies insignificant impact on  $CO_2$  and  $SO_2$  emissions. In Sweden, the allowed sulphur content in diesel fuel is < 10 ppm, (< 0.001%), and given the negligible impact on fuel demand, it is reasonable to assume an equally negligible impact on  $SO_2$  emissions. In this paper, we adhere to reported numbers, but follow up on them in the discussion.

### 2.1.2 Metric choice and climate impact uncertainty

We use climate metrics presented by the Intergovernmental Panel on Climate Change (IPCC) in their 5th Assessment Report on climate change (Myhre et al. 2013a, b) as the primary source of information on metric value, and complemented with IPCC sources when necessary (Collins et al. 2013). The IPCC and Collins et al. (2013) present values for the climate metrics  $GWP_{100}$ ,  $GWP_{20}$ ,  $GTP_{100}$ , and  $GTP_{20}$  for region-specific as well as for global average (or 4-region aggregates) emissions of BC, OC, NMVOC,  $CH_4$ , and  $NO_x$  (given in tables 8.A.1.3,4,5, & 6 in Myhre et al. (2013b); table 8.SM.14 in Myhre et al. (2013a); and tables 1 and 2 in Collins et al. (2013)). Out of the available regions, ‘Europe’ is considered most representative of Sweden.

The estimates of the radiative forcing (RF), climate impacts, and consequently the  $CO_2$  equivalent emissions of the pollutants covered in this paper are uncertain. We accommodate for this uncertainty by assigning low, mid, high numerical values of  $CO_{2eq}$  emissions for each pollutant and each climate metric: the low and high numerical values correspond to  $\pm$  one standard deviation, respectively.

The uncertainty ranges presented in the literature are not presented using similar uncertainty intervals, so some adaptations are necessary. For BC, we use low, mid, high values from table 2 in Collins et al. (2013) for European GTP metrics; calculate European GWP metrics based on low, mid, high values on absolute GWP (AGWP) from table 1 in Collins et al. (2013) divided with  $AGWP_{20}$  and  $AGWP_{100}$  for  $CO_2$  in Collins et al. (2013); and use low, mid, high values from Myhre et al. (2013b: table 8.A.6 row ‘BC total, global’) for global GTP and GWP metrics. For OC, we use the same approach as for BC, but since the presentation of global metric values is incomplete in the

**Table 2** Reported average national emission reduction of air pollutants per option

Abatement option	Pollutant						Reference
	$PM_{res}^a$ [tonne]	BC [tonne]	OC [tonne]	NMVOC [tonne]	$CH_4$ [tonne]	$NO_x$ [tonne]	
Pellets	468	166	308	1571	1882	44	CLEO (2014)
Mod. boiler	80	168	128	1617	2062		Amann (2014); CLEO (2014)
Man. ferm.					7750		Hellstedt et al. (2014); Swedish Board of Agriculture (2012)
Sl. cover					500		Swedish Board of Agriculture (2012)
Sl. acid.					500		Swedish Board of Agriculture (2012)
Mod. NRMM	13	38				1017	Åström et al. (2013)
4S S-mob.				80			Fridell and Åström (2009)
4S&EI NRMM				2100			Fridell and Åström (2009)
W. solv.				30,000			Amann et al. (2014)

<sup>a</sup> Emissions of  $PM_{res}$  is our own estimate calculated as  $PM_{res} = PM_{2.5} - BC - OC$

source material, we use average values for emissions from ‘East Asia’, ‘EU + North Africa’, ‘North America’, and ‘South Asia’ as proxies for global average emissions (Myhre et al. 2013b: table 8.A.6 row ‘4 regions’). For NMVOC, we use values from Myhre et al. (2013b: table 8.A.5) for all metrics. For European metrics, we use ‘EU + North Africa’ and for global metrics, we use ‘four regions above’. For NO<sub>x</sub>, we use values from Myhre et al. (2013b: table 8.A.3): ‘EU + North Africa’ values for European metrics, and ‘four above regions’ values for global metrics. For CH<sub>4</sub>, we do not separate European and global metrics. We use values from Myhre et al. (2013b: table 8.A.1) for GTP and GWP mid values. GTP low and high values are estimated by calculating the percentage deviation from mid values in Collins et al. (2013) table 2, and applying these percentages on the mid value from Myhre et al. (2013b: table 8.A.1). GWP low and high values are taken from Myhre et al. (2013a: table 8.SM.14) and deflated from a 90% confidence interval to one standard deviation (assuming a normal distribution of CH<sub>4</sub> uncertainty).

For PM<sub>res</sub>, we assume that the CO<sub>2</sub> equivalence of emissions is equal to the CO<sub>2</sub> equivalence of OC emissions, given that most PM<sub>res</sub> emissions from biomass burning are inorganic (mineral) salts with mainly light scattering properties (Chen et al. 2017). Due to the crudeness of this assumption, we test the robustness of the results by setting the CO<sub>2</sub> equivalent emissions of PM<sub>res</sub> emissions to zero in the sensitivity analysis. Finally, following the update recommendations in Myhre et al. (2013b), we update the 100-year perspective values with a factor 0.94 for GWP<sub>100</sub> and 0.92 for GTP<sub>100</sub>, an update not necessary for the 20-year time horizon due to negligible effect on that short-time horizon. This update is made to accommodate for updated values of the absolute metric value of CO<sub>2</sub>. The resulting climate metric values used in this paper are found in Table 3.

There exist up to 729 (3<sup>6</sup>) possible combinations of high, medium, and low CO<sub>2</sub> equivalence for the six pollutants. For each of the eight metrics, each of these 729 combinations leads to an ordered ranking of the nine abatement options. All in all, each option thereby has 5832 potential CO<sub>2</sub> equivalent emissions impacts.

## 2.2 Calculation method

We calculate cost-effectiveness of the nine options using these 729 CO<sub>2</sub> equivalent abatement levels for each option by dividing the cost of the option (Table 1) with the sum over all pollutants of the product of emission reduction (Table 2) and the metric-specific CO<sub>2eq</sub> values (Table 3) to give abatement cost per tonne CO<sub>2eq</sub> (Eq. 1). This calculation is done separately for all eight metrics. We do not consider uncertainty in costs and emission levels since the focus of this paper is on the effect of climate metrics on cost-effectiveness.

$$\text{Unit cost}_{o,m,r} = \frac{c_o}{\sum_p (\text{red}_{o,p} * \text{CI}_{p,m,r})} \quad (1)$$

where

<i>o</i>	abatement option
<i>p</i>	pollutant
<i>m</i>	climate metric
<i>r</i>	climate metric value range
Unit cost	cost per tonne CO <sub>2eq</sub> abated [M€ <sub>2010</sub> /tonne CO <sub>2eq</sub> ]
<i>c</i>	option cost [M€ <sub>2010</sub> ]
red	emission abatement [tonne]
CI	metric value [tonne CO <sub>2eq</sub> /tonne emission]

**Table 3** Values of CO<sub>2</sub> equivalent emissions per pollutant and climate metric considered in the analysis (Collins et al. 2013; Myhre et al. 2013a, b), including our assumption of PM<sub>res</sub> having the same climate impact as OC

Metric values from European emissions						
	GWP <sub>20-E</sub> [tonne CO <sub>2eq</sub> /tonne emission]			GWP <sub>100-E</sub> [tonne CO <sub>2eq</sub> /tonne emission]		
	Low (-1 std.)	Mid	High (+1 std.)	Low (-1 std.)	Mid	High (+1 std.)
PM <sub>res</sub>	-104.0 <sup>a</sup>	-172.0 <sup>a</sup>	-240.0 <sup>a</sup>	-28.1 <sup>a</sup>	-46.5 <sup>a</sup>	-64.8 <sup>a</sup>
BC	720.0 <sup>b</sup>	1480.0 <sup>b</sup>	2240.0 <sup>b</sup>	194.5 <sup>b</sup>	399.8 <sup>b</sup>	605.1 <sup>b</sup>
OC	-104.0 <sup>b</sup>	-172.0 <sup>b</sup>	-240.0 <sup>b</sup>	-28.1 <sup>b</sup>	-46.5 <sup>b</sup>	-64.8 <sup>b</sup>
NMVOC	9.5 <sup>c</sup>	18.0 <sup>c</sup>	26.5 <sup>c</sup>	2.6 <sup>c</sup>	5.3 <sup>c</sup>	7.9 <sup>c</sup>
CH <sub>4</sub>	68.8 <sup>d</sup>	84.0 <sup>c</sup>	99.2 <sup>d</sup>	21.4 <sup>d</sup>	28.0 <sup>c</sup>	34.6 <sup>d</sup>
NO <sub>x</sub>	-21.9 <sup>c</sup>	-39.4 <sup>c</sup>	-56.9 <sup>c</sup>	-9.2 <sup>c</sup>	-14.7 <sup>c</sup>	-20.1 <sup>c</sup>
	GTP <sub>20-E</sub> [tonne CO <sub>2eq</sub> /tonne emission]			GTP <sub>100-E</sub> [tonne CO <sub>2eq</sub> /tonne emission]		
	Low (-1 std.)	Mid	High (+1 std.)	Low (-1 std.)	Mid	High (+1 std.)
PM <sub>res</sub>	-40.0 <sup>a</sup>	-58.0 <sup>a</sup>	-76.0 <sup>a</sup>	-4.9 <sup>a</sup>	-7.1 <sup>a</sup>	-9.3 <sup>a</sup>
BC	340.0 <sup>b</sup>	530.0 <sup>b</sup>	720.0 <sup>b</sup>	42.3 <sup>b</sup>	65.3 <sup>b</sup>	88.3 <sup>b</sup>
OC	-40.0 <sup>b</sup>	-58.0 <sup>b</sup>	-76.0 <sup>b</sup>	-4.9 <sup>b</sup>	-7.1 <sup>b</sup>	-9.3 <sup>b</sup>
NMVOC	3.0 <sup>c</sup>	9.5 <sup>c</sup>	16.0 <sup>c</sup>	0.3 <sup>c</sup>	0.7 <sup>c</sup>	1.2 <sup>c</sup>
CH <sub>4</sub>	56.3 <sup>b</sup>	67.0 <sup>c</sup>	77.7 <sup>b</sup>	2.7 <sup>b</sup>	4.0 <sup>c</sup>	5.3 <sup>b</sup>
NO <sub>x</sub>	-33.1 <sup>c</sup>	-48.0 <sup>c</sup>	-62.9 <sup>c</sup>	-1.1 <sup>c</sup>	-2.3 <sup>c</sup>	-3.5 <sup>c</sup>
Metric values from global average emissions						
	GWP <sub>20-G</sub> [tonne CO <sub>2eq</sub> /tonne emission]			GWP <sub>100-G</sub> [tonne CO <sub>2eq</sub> /tonne emission]		
	Low (-1 std.)	Mid	High (+1 std.)	Low (-1 std.)	Mid	High (+1 std.)
PM <sub>res</sub>	-92.0 <sup>a</sup>	-160.0 <sup>a</sup>	-228.0 <sup>a</sup>	-24.4 <sup>a</sup>	-43.2 <sup>a</sup>	-62.0 <sup>a</sup>
BC	270.0 <sup>c</sup>	3200.0 <sup>c</sup>	6200.0 <sup>c</sup>	94.0 <sup>c</sup>	846.0 <sup>c</sup>	1598.0 <sup>c</sup>
OC	-92.0 <sup>c</sup>	-160.0 <sup>c</sup>	-228.0 <sup>c</sup>	-24.4 <sup>c</sup>	-43.2 <sup>c</sup>	-62.0 <sup>c</sup>
NMVOC	11.2 <sup>c</sup>	18.7 <sup>c</sup>	26.2 <sup>c</sup>	3.1 <sup>c</sup>	5.5 <sup>c</sup>	7.8 <sup>c</sup>
CH <sub>4</sub>	68.8 <sup>d</sup>	84.0 <sup>c</sup>	99.2 <sup>d</sup>	21.4 <sup>d</sup>	28.0 <sup>c</sup>	34.6 <sup>d</sup>
NO <sub>x</sub>	16.8 <sup>c</sup>	-15.9 <sup>c</sup>	-48.6 <sup>c</sup>	-0.8 <sup>c</sup>	-10.9 <sup>c</sup>	-21.0 <sup>c</sup>
	GTP <sub>20-G</sub> [tonne CO <sub>2eq</sub> /tonne emission]			GTP <sub>100-G</sub> [tonne CO <sub>2eq</sub> /tonne emission]		
	Low (-1 std.)	Mid	High (+1 std.)	Low (-1 std.)	Mid	High (+1 std.)
PM <sub>res</sub>	-39.0 <sup>a</sup>	-55.0 <sup>a</sup>	-71.0 <sup>a</sup>	-4.8 <sup>a</sup>	-6.7 <sup>a</sup>	-8.6 <sup>a</sup>
BC	95.0 <sup>c</sup>	920.0 <sup>c</sup>	2400.0 <sup>c</sup>	4.6 <sup>c</sup>	119.6 <sup>c</sup>	312.8 <sup>c</sup>
OC	-39.0 <sup>c</sup>	-55.0 <sup>c</sup>	-71.0 <sup>c</sup>	-4.8 <sup>c</sup>	-6.7 <sup>c</sup>	-8.6 <sup>c</sup>
NMVOC	4.3 <sup>c</sup>	10.0 <sup>c</sup>	15.7 <sup>c</sup>	0.4 <sup>c</sup>	0.8 <sup>c</sup>	1.3 <sup>c</sup>
CH <sub>4</sub>	56.3 <sup>b</sup>	67.0 <sup>c</sup>	77.7 <sup>b</sup>	2.7 <sup>b</sup>	4.0 <sup>c</sup>	5.3 <sup>b</sup>
NO <sub>x</sub>	-35.9 <sup>c</sup>	-62.1 <sup>c</sup>	-88.3 <sup>c</sup>	-0.1 <sup>c</sup>	-2.0 <sup>c</sup>	-4.0 <sup>c</sup>

<sup>a</sup> Own assumption, <sup>b</sup> Collins et al. (2013), <sup>c</sup> Myhre et al. (2013b), <sup>d</sup> Myhre et al. (2013a)

The resulting abatement cost per ton CO<sub>2eq</sub> for each option for mid values of the eight metrics is shown in Table 4. For certain metrics, some combinations of metric values can result in negative CO<sub>2eq</sub> emission abatement i.e. implying increasing CO<sub>2eq</sub> emissions. For example, this may happen if some emissions decrease and others increase due to the abatement options, or in our case where it mainly depends on time horizon for NO<sub>x</sub> (which is short-term warming whilst long-term cooling). These cases are indicated with N.A. in Table 4.

As can be seen, some abatement options are several orders of magnitude more expensive per unit CO<sub>2eq</sub> abatement than the cheapest option for all metrics.

For each unique combination of climate metric and metric value, we rank the cost-effectiveness of the nine options from most cost-effective (rank #1) to least cost-effective (rank #9). As mentioned, some options can increase CO<sub>2eq</sub> emission levels for specific metric values. We therefore first control whether the option would increase or decrease CO<sub>2eq</sub> emission levels. If one or several control option leads to an increase in the CO<sub>2eq</sub> emissions, those are considered least cost-effective, while the other options are simply ranked according to their relative cost-effectiveness.

**Table 4** Abatement cost in €<sub>2010</sub>/tonne CO<sub>2eq</sub> for the studied options, across the mid CO<sub>2eq</sub> values of all the studied climate metrics

Abatement option	Metric							
	GWP <sub>20-E</sub>	GWP <sub>100-E</sub>	GTP <sub>20-E</sub>	GTP <sub>100-E</sub>	GWP <sub>20-G</sub>	GWP <sub>100-G</sub>	GTP <sub>20-G</sub>	GTP <sub>100-G</sub>
Pellets	62	177	97	1160	31	104	65	688
Mod. boiler	12	35	20	228	6	21	14	153
Man. ferm.	49	148	62	1040	49	148	62	1040
Sl. cover	76	229	96	1600	76	229	96	1600
Sl. acid.	198	593	248	4150	198	593	248	4150
Mod. NRMM	42,000	N.A.	N.A.	2,900,000	1000	7000	N.A.	60,000
4S S-mob.	69	236	132	1790	89	278	125	1560
4S&EI NRMM	103	350	195	2650	133	413	186	2320
W. solv.	333	1130	632	8570	429	1330	600	7500

N.A. implies that a metric indicates increase in CO<sub>2eq</sub> emissions

### 2.3 Sensitivity analysis

We make five different sensitivity analyses. In the first sensitivity analysis, we assume that the radiative forcing strength (in absolute terms) of the particulate sub-fractions (PM<sub>res</sub>, BC, OC) would be perfectly correlated, as discussed in Aamaas et al. (2016). This implies that we make the calculations in three separate groups. In one group, all PM sub-fractions are assigned low-metric values (from Table 3), the next mid, and the third high (which gives 81 possible combinations of CO<sub>2</sub> equivalent emissions for each of the eight metrics). In the second sensitivity analysis, we disregard all climate impact from NO<sub>x</sub> emissions in the calculations (243 combinations for each metric). This sensitivity analysis is made since NO<sub>x</sub> is the only pollutant with climate metric values ranging from positive to negative depending metric choice and time horizon, which can have a large impact on ranking. In the third sensitivity analysis, we assume that the CO<sub>2eq</sub> of PM<sub>res</sub> is zero (243 combinations), in contrast to the main analysis where we assume that the CO<sub>2eq</sub> of PM<sub>res</sub> is equal to OC.

The fourth sensitivity analysis is made in order to verify that any potential robustness of the ranking found in the main analysis is not primarily a result of very uneven costs of the different abatement options (Table 4). In this sensitivity analysis, we construct a hypothetical marginal abatement cost function where the cost of the least cost option starts at 20 €<sub>2010</sub>/tonne CO<sub>2eq</sub> when using mid values of GWP<sub>100-G</sub>, and where the marginal abatement cost increases with an increment of 10 €<sub>2010</sub>/tonne CO<sub>2eq</sub> for the subsequent options. Hence, in this sensitivity analysis, the most cost-effective option has a cost of 20 €<sub>2010</sub>/tonne CO<sub>2eq</sub>, the second most cost-effective 30 €<sub>2010</sub>/tonne CO<sub>2eq</sub> and so on up until 100 €<sub>2010</sub>/tonne CO<sub>2eq</sub> for the least cost-effective option. This corresponds to the unit costs in Table 1 being recalibrated into 6.2 M€/year (*Pellets*); 4.0 M€/year (*Mod. boiler*); 6.5 M€/year (*Man. ferm*); 0.84 M€/year (*Sl. cover*); 1.2 M€/year (*Sl. acid*); 2.2 M€/year (*Mod. NRMM*); 0.018 M€/year (*4S S-mob*); 0.66 M€/year (*4S&EL NRMM*); 11 M€/year (*W. solv*). In the fifth sensitivity analysis, we combine sensitivity analysis #2 and 4.

## 3 Results

We present cost-effectiveness ranking results for all the possible rankings of the nine options. We first show the results of the main analysis (aggregated for all metrics), followed by



disaggregated comparisons of option ranking per metric. We also present the effects of region, metric choice, and time horizon, as well as results from the sensitivity analyses on PM correlation and the  $PM_{res}$  assumption. Finally, in brief, we present the disaggregated analysis of the effects of region, metric type, and time horizon, as well as the sensitivity analyses on  $NO_x$  effects and abatement costs, but leave the details to the supplementary material 2.

### 3.1 Results from the main analysis

Most options have a relatively stable distribution with the 2nd and 3rd quartile ranking and average values within  $\pm 1$  from the median (Fig. 1). The option with the largest variation is *Pellets*, which is the only option affecting all six pollutants.


The relative robustness of the ranking can be further confirmed when focusing on the range around the mode ranking (the most common ranking). In Table 5, it can be seen that all options have rankings occurring around the mode ranking  $\pm 1$  for more than two-thirds of the occasions. The general picture emerging from Fig. 1 and Table 5 is that the ranking is relatively robust with respect to climate metric.

### 3.2 Disaggregation of the main analysis

The results are disaggregated in order to analyse which of the climate metrics that has the strongest impact on the variation of the ranking of the abatement options. The disaggregation of the results also allows for an estimate of which feature of the metric design that could be considered to drive the variation in rank.

#### 3.2.1 Ranking per metric

The mode ranking of cost-effectiveness can be seen to be stable across metrics, with largest variation for *4S S-mob*, followed by *Man. ferm* and *Pellets*. The variation in mode rank for *4S S-mob* is likely caused by the variation in metric values for  $CH_4$  and BC, which are important for the rank range of *Man. ferm*, *Pellets*, and *Sl. cover* (Fig. 2).



**Fig. 1** Distribution of relative ranking of cost-effectiveness ( $\text{€}_{2010}/\text{tonne CO}_{2eq}$ ) for each option. The grey box shows the range of ranks for the 2nd and 3rd quartile of the results, the grey line in the boxes shows the median rank, the cross shows the average, and the error bar shows the 90th percentile range

**Table 5** The mode rank for each option and percent of all ranks that surround the mode rank for the option


Option	Mode rank	% of occurrence around mode $\pm 1$
Mod. boiler	#1	100%
Man. ferm.	#2	92%
Pellets	#2	64%
Sl. cover	#4	89%
4S S-mob.	#5	73%
4S&EI NRMM	#6	94%
Sl. acid.	#7	88%
W. solv.	#8	99%
Mod. NRMM	#9	100%
Mod. boiler	#1	100%

Furthermore, the disaggregated mode ranks show that the stability for the aggregated results (Table 5) is seen also for the disaggregated results (Table 6).

Also, for the 729 CO<sub>2eq</sub> emission levels per option and metric, the difference between max and min rank as a function of metric remains within two rank steps in more than two-thirds of the outcomes for all options (Table 7). Again, the option with the largest variation is *Pellets*.

### 3.2.2 Effects on ranking of regional scope, metric type, and time horizon

In order to improve the understanding of the variation in ranking, we separate out the results into global metrics vs. European metrics; GTP vs. GWP; and 20-year vs. 100-year time horizon. Through these separations we observe: that the option ranking varies more for global



**Fig. 2** The options relative ranking of cost-effectiveness (€<sub>2010</sub>/tonne CO<sub>2eq</sub>) for each climate metric. The coloured boxes show the mode rank of each option, and the error bar shows the 90th percentile range

**Table 6** Percent of all option ranks that are at the Table 5 mode rank  $\pm 1$ , presented per metric

Metric/option	GWP <sub>20-E</sub>	GWP <sub>100-E</sub>	GTP <sub>20-E</sub>	GTP <sub>100-E</sub>	GWP <sub>20-G</sub>	GWP <sub>100-G</sub>	GTP <sub>20-G</sub>	GTP <sub>100-G</sub>
Pellets	61%	61%	51%	70%	67%	67%	67%	67%
Mod. boiler	100%	100%	100%	100%	100%	100%	100%	100%
Man. ferm.	89%	95%	100%	93%	85%	93%	100%	81%
Sl. cover	83%	91%	100%	79%	85%	93%	100%	78%
Sl. acid.	100%	89%	67%	78%	100%	100%	78%	97%
Mod. NRMM	100%	100%	100%	100%	100%	100%	100%	100%
4S S-mob.	64%	73%	78%	75%	67%	74%	89%	67%
4S&EI NRMM	93%	96%	100%	95%	93%	96%	100%	81%
W. solv.	100%	100%	100%	100%	99%	100%	100%	97%

metrics than European metrics (consistent with the larger variation in metric values for global metrics, especially the larger BC and NO<sub>x</sub> metric value variation in GWP<sub>20-G</sub> and GWP<sub>100-G</sub>); that GTP rankings generally are more stable than GWP rankings (consistent with the relatively large BC and NO<sub>x</sub> value variation in GWP<sub>20-G</sub> and GWP<sub>100-G</sub>); and that a 100-year time horizon has a tendency to increase variation in ranking (consistent with most pollutants having slightly larger variation in metric values for the 100-year time horizon than the 20-year time horizon). However, neither the regional scope, nor metric type, nor time horizon has a significant effect on the variation in ranking.

### 3.3 Results from the sensitivity analysis

We present the results from the first (correlation in RF strength of PM species) and third (neglecting RF of PM<sub>res</sub>) sensitivity analysis. The results from the second (no NO<sub>x</sub>), fourth (hypothetical even distribution of abatement costs), and fifth (combination of #2 and #4) sensitivity analyses are shown in the supplementary material 2 and only presented in brief in the main text.

#### 3.3.1 If the radiative forcing of PM-subspecies is correlated

In the first sensitivity analysis, we test the sensitivity of the ranking by letting the low, mid, and high CO<sub>2eq</sub> values (in absolute terms) for PM<sub>res</sub>, BC, and OC in Table 3 be correlated when

**Table 7** Rank variation as a function of metric over all 729 combinations, presented per option in percentage

	Pellets	Mod. boiler	Man. ferm.	Sl. cover	Sl. acid.	Mod. NRMM	4S S-mob.	4S&EI NRMM	W. solv.
0	1%	100%	26%	44%	50%	93%	16%	23%	77%
1	55%	0%	56%	38%	17%	7%	59%	52%	20%
2	28%	0%	11%	18%	33%	0%	25%	23%	3%
3	11%	0%	7%	0%	0%	0%	0%	2%	0%
4	5%	0%	0%	0%	0%	0%	0%	0%	0%
5	0%	0%	0%	0%	0%	0%	0%	0%	0%
6	0%	0%	0%	0%	0%	0%	0%	0%	0%
7	0%	0%	0%	0%	0%	0%	0%	0%	0%
8	0%	0%	0%	0%	0%	0%	0%	0%	0%
Tot	100%	100%	100%	100%	100%	100%	100%	100%	100%

calculating cost-effectiveness and ranking. Overall, the relative ranking of the options is similar in the sensitivity analysis and the main analysis (Fig. 3).


### 3.3.2 Disregarding climate impact of $PM_{res}$

In the third sensitivity analysis, we test the sensitivity of the ranking by setting the  $CO_2$  equivalent emissions of  $PM_{res}$  to zero. The exclusion of  $PM_{res}$  has a stabilising effect on the ranking, with noticeably reduced variability of *Pellets* and *4S S-mob* (Fig. 4).


### 3.3.3 Summary of additional sensitivity analyses

The sensitivity analyses presented in the supplementary material 2 indicate similar results as the main analysis. Regarding the hypothetical equalised cost distribution case (sensitivity analysis #4), the results show an expected decrease in the robustness of ranking compared with the main analysis. The median and average ranking of options are relatively stable and comparable with what they were in the main analysis except for the option *Pellets* that change median ranking from 3 to 5 and average from 3.3 to 5.1.

There is however another result from the sensitivity analysis based on the hypothetical even cost that is of higher interest. When studying the details, it can be observed that the increased variation in rank (noticeable for all options in this hypothetical case) seems especially large and irregular for *Mod. NRMM*. In contrast with the other options, *Mod. NRMM* has a noticeable larger variation in ranking for the global metrics than for European metrics as well as for GWP rather than for GTP (Figure SM2 13 and SM2 14). This is likely explained by the fact that the abatement option affects  $NO_x$  emissions and that the metric value for  $NO_x$  change sign under some circumstances for GWP global (while not for the other metrics). In all cases but one, the metric for  $NO_x$  is negative, but for the case  $GWP_{20-G-Low}$ , it turns positive. This causes the ranking of *Mod. NRMM* to vary with metric choice and assumption about time horizon and forcing strength. Further, an indication of validity of this explanation is that the ranking distribution of *Mod. NRMM* has a bimodal characteristic (Table SM2 3), it most often ranks last or close to last when  $NO_x$  emissions have a negative metric value, but when  $NO_x$  emissions have a positive metric value, the occurrence of ranks around place three almost



**Fig. 3** Distribution of relative ranking for each option if assuming that low, mid, and high  $CO_2$  equivalence of BC, OC, and  $PM_{res}$  are correlated (648 possible ranks per option). The grey box shows the range of ranks for the 2nd and 3rd quartile of the results, the grey line in the boxes shows the median rank, the cross shows the average, and the error bar shows the 90th percentile



**Fig. 4** Distribution of relative ranking for each option if assuming that  $PM_{res}$  emissions have no radiative forcing (243  $CO_2$  equivalence combinations per metric). The grey box shows the range of ranks for the 2nd and 3rd quartile of the results, the grey line in the boxes shows the median rank, the cross shows the average, and the error bar shows the 90th percentile

doubles. To check the robustness of this explanation, we carry out an additional sensitivity analysis (#5—even cost and no  $NO_x$  assumption). However, the expected attenuation in rank variation for *Mod. NRMM* as measured in Figures SM2 16 to 19 is not visible, but the bimodal characteristics of the ranking of *Mod. NRMM* is weakened (Table SM2 4). Hence, the likely explanation of the observed variation of *Mod. NRMM* rank and its bimodal characteristics in sensitivity analysis #4 are due to that the  $CO_2$  equivalent emission of  $NO_x$  change sign for the  $GWP_{20-G-Low}$  metric.

## 4 Discussion and implications

Many different climate metrics have been suggested and are found in the literature. In this paper, we analyse to what extent estimated relative cost-effectiveness of options to reduce  $CO_{2eq}$  emissions through air pollution abatement is affected by the choice of climate metric used in the calculations; a choice that is at least partly normative. We calculate cost-effectiveness for nine NTCF abatement options expected to be available in Sweden using eight different climate metric designs (two regional scopes, two metric types, and two-time horizons), and then compare how the abatement options' relative cost-effectiveness would be affected by the metric-related choices. Overall, our results show comparatively robust relative cost-effectiveness (ranking), with the reservation that options having a large effect on  $NO_x$  might be an exception. This implies that the choice of metric will not strongly affect policy recommendations regarding the relative cost-effectiveness of abatement options affecting only emissions of NTCFs, at least when  $NO_x$  is not heavily affected. However, it also implies that potential effects on  $NO_x$  are important to include when analysing cost-effective NTCF abatement options. In fact, the potential impact of  $NO_x$  provides an indication that care should be taken and that all relevant NTCFs and LLGHGs, including  $SO_2$ ,  $CO$ , and  $CO_2$ , should be included in an analysis of policy measures on NTCFs.

Our results are in apparent contrast to much of the literature on climate metrics for policy analysis. As an example, Grewe and Dahlmann (2015) stress the importance of increasing the precision of the [policy] question asked, as well as finding adequate references, emission scenarios, indicators, and time horizons, since all of these factors will impact which climate metric to use in

analyses. Further, Tanaka et al. (2010) state that the choice of metric will affect the balance of which options to prefer when choosing between ‘multicomponent’ options. The important difference between these studies and our analysis is that all these and other studies concerns comparisons of either different LLGHGs or a combination of different LLGHGs and NTCFs. Herein lays a key message from our paper: it is important to consider the different challenges faced when designing policy for LLGHG and NTCF abatement versus when designing policy for only NTCF abatement where the effect on LLGHG are negligible. When only considering a policy for NTCF abatement, where the LLGHG are not affected, there is likely no need to worry about the effect of metric choices on the outcome (with reservation for  $\text{NO}_x$ ). In other words, if analysing cost-effectiveness of NTCF emission abatement, the existing climate metrics are showing relatively robust results when it comes to the ranking of the relative cost-effectiveness on various abatement options (given that options have insignificant effect on  $\text{NO}_x$  emissions).

Even though the metric values depend strongly on metric design (as is pointed out by Grewe and Dahlmann (2015) and Tanaka et al. (2010)), the relatively robust ranking can be understood through that the metric values for the pollutants are quite strongly correlated over metrics and time horizons, with the exception for  $\text{NO}_x$ .  $\text{CO}_2$  is the reference gas for all metrics used in this analysis, and it has a very long climate perturbation time compared with the perturbation time of NTCFs. The variation in metric values would be less affected by metric design choices if a short-lived gas such as methane would be used as reference gas (Cherubini and Tanaka 2016). For this reason, and given the current policy separation of NTCF and LLGHG abatement, it may make sense to use methane as a reference gas when analysing cost-effectiveness of NTCF abatement options. This will not affect the relative ranking in our analysis to any significant extent, but would make related policy assessments more transparent in the sense that the metric value for each NTCF would be more stable over different metric choices and time horizons.

One limitation is that we analyse only nine NTCF abatement options in a single country, and our results should therefore be complemented with other studies made for more countries. The abatement options though are rather generic and likely available in most countries. Further, we do not include  $\text{SO}_2$  in the analysis, since the options considered in the analysis does not have an effect on  $\text{SO}_2$  emissions. Also, the results are partly dependent on the wide variation in abatement costs of the considered options, which we controlled for in the fourth sensitivity analysis (supplementary material 2).

The results presented in this paper are relevant since there is a risk that decision makers will mentally transfer the existing general message from metric studies—the choice of metric will have an effect, potentially a large effect, on the balance of options—onto decisions regarding NTCF-only abatement that have no (or little) effect on the emissions of LLGHGs. All in all, our results suggest that there is limited need for policymakers to be concerned by the large variations of metrics used in studies of cost-effective solutions for NTCF control as long as only NTCFs (with  $\text{NO}_x$  emissions exempted) and not LLGHGs are affected by the abatement options considered.

**Acknowledgements** The authors thank Rasmus Einarsson (Chalmers University of Technology) for the help with the code preparation. Finally, we thank three anonymous reviewers for constructive comments to drafts of the manuscript.

**Funding information** This research was funded by the Swedish Environmental Protection Agency through the SCAC research programme (grant number NV-03730-1616/87). The funding source had no involvement in the analysis or writing of this paper.

## Compliance with ethical standards

**Conflict of interest** The authors declare that they have no conflict of interest.

**Open Access** This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made.

## References

- Aamaas B, Peters GP, Fuglestvedt JS (2013) Simple emission metrics for climate impacts. *Earth Syst Dynam* 4: 145–170. <https://doi.org/10.5194/esd-4-145-2013>
- Aamaas B, Bernsten TK, Fuglestvedt JS, Shine KP, Bellouin N (2016) Regional emission metrics for short-lived climate forcers from multiple models. *Atmos Chem Phys* 16:7451–7468. <https://doi.org/10.5194/acp-16-7451-2016>
- ACAP (2014) Reduction of black carbon emissions from residential wood combustion in the Arctic - Black Carbon Inventory, Abatement Instruments and Measures. Arctic Contaminants Action Program (ACAP). <http://hdl.handle.net/11374/1365>. Accessed 17 Dec 2018
- Amann M (2014) Updates to the GAINS model databases after the bilateral consultations with national experts in 2014 - TSAP report #14 version 1.0
- Amann M et al (2014) The final policy scenarios of the EU clean air policy package, TSAP report #11
- AMAP, Bluestein J, Rackley J, Baum E (2008) Sources and mitigation opportunities to reduce emissions of short-term Arctic climate forcers. Arctic Monitoring and Assessment Programme (AMAP), Arctic Council (2011) An assessment of emissions and mitigation options for black carbon for the Arctic Council
- Åström S, Bäckström S, Jerksjö M, Fridell E (2013) Åtgärder för att minska utsläpp av NOx och PM2,5 från den svenska transportsektorn 2025 - potential och kostnader, IVL rapport B2111
- Bowerman NHA, Frame DJ, Huntingford C, Lowe JA, Smith SM, Allen MR (2013) The role of short-lived climate pollutants in meeting temperature goals. *Nat Clim Chang* 3:1021–1024. <https://doi.org/10.1038/nclimate2034>
- Chen J et al (2017) A review of biomass burning: emissions and impacts on air quality, health and climate in China. *Sci Total Environ* 579:1000–1034. <https://doi.org/10.1016/j.scitotenv.2016.11.025>
- Cherubini F, Tanaka K (2016) Amending the inadequacy of a single Indicator for climate impact analyses. *Environ Sci Technol* 50:12530–12531. <https://doi.org/10.1021/acs.est.6b05343>
- CLEO (2014) Klimatförändringen och miljömålen - Rapport till Naturvårdsverket inför Fördjupad Utvärdering 2015
- Collins WJ, Fry MM, Yu H, Fuglestvedt JS, Shindell DT, West JJ (2013) Global and regional temperature-change potentials for near-term climate forcers. *Atmos Chem Phys* 13:2471–2485. <https://doi.org/10.5194/acp-13-2471-2013>
- Fesenfeld LP, Schmidt TS, Schrode A (2018) Climate policy for short- and long-lived pollutants. *Nat Clim Chang* 8:933–935. <https://doi.org/10.1038/s41558-018-0328-1>
- Fridell E, Åström S (2009) Analysis of measures to reduce Swedish emissions by 2020 for NOx, PM2.5 and NMVOC - non-road machinery and shipping, IVL report U2617
- Grewe V, Dahlmann K (2015) How ambiguous are climate metrics? And are we prepared to assess and compare the climate impact of new air traffic technologies? *Atmos Environ* 106:373–374. <https://doi.org/10.1016/j.atmosenv.2015.02.039>
- Hellstedt C, Cerruto J, Nilsson M, McCann M (2014) Nordic initiatives to abate methane emissions - a catalogue of best practices. <https://doi.org/10.6027/anp2014-741>
- Henze DK et al (2012) Spatially refined aerosol direct radiative forcing efficiencies. *Environ Sci Technol* 46: 9511–9518. <https://doi.org/10.1021/es301993s>
- Johansson DJA (2011) Economics- and physical-based metrics for comparing greenhouse gases. *Clim Chang* 110:123–141. <https://doi.org/10.1007/s10584-011-0072-2>
- Kindbom K, Åström S, Yaramenka K, Wisell T, Sjödin Å (2015) Kortlivade klimatpåverkande luftföroreningar (SLCP) - Underlag för övervägande om etappmål, IVL-rapport C99

- Kuylenstierna JCI et al (2011) Near-term climate protection and clean air benefits: Actions for Controlling Short-Lived Climate Forcers
- Levasseur A et al (2016) Enhancing life cycle impact assessment from climate science: review of recent findings and recommendations for application to LCA. *Ecol Indic* 71:163–174. <https://doi.org/10.1016/j.ecolind.2016.06.049>
- Manne AS, Richels RG (2001) An alternative approach to establishing trade-offs among greenhouse gases. *Nature* 410:675–677. <https://doi.org/10.1038/35070541>
- Manning M, Reisinger A (2011) Broader perspectives for comparing different greenhouse gases. *Philos Trans R Soc A Math Phys Eng Sci* 369:1891–1905. <https://doi.org/10.1098/rsta.2010.0349>
- Myhre G et al (2013a) Anthropogenic and natural radiative forcing supplementary material. In: Stocker TF et al. (eds) *Climate change 2013: the physical science basis. Contribution of working group I to the fifth assessment report of the intergovernmental panel on climate change*
- Myhre G et al (2013b) Anthropogenic and natural radiative forcing. In: Stocker TF et al (eds) *Climate change 2013: the physical science basis. Contribution of working group I to the fifth assessment report of the intergovernmental panel on climate change*. Cambridge University Press, Cambridge
- Norwegian Environment Agency (2014) Summary of proposed action plan for Norwegian emissions of short-lived climate forcers
- Peters GP, Aamaas B, Berntsen T, Fuglestedt JS (2011) The integrated global temperature change potential (iGTP) and relationships between emission metrics. *Environ Res Lett* 6:044021. <https://doi.org/10.1088/1748-9326/6/4/044021>
- Sarofim MC, Giordano MR (2018) A quantitative approach to evaluating the GWP timescale through implicit discount rates. *Earth Syst Dynam* 9:1013–1024. <https://doi.org/10.5194/esd-9-1013-2018>
- Schmale J, van Aardenne J, von Schneidemesser E (2014) New directions: support for integrated decision-making in air and climate policies – development of a metrics-based information portal. *Atmos Environ* 90: 146–148. <https://doi.org/10.1016/j.atmosenv.2014.03.016>
- Shindell D, Faluvegi G (2009) Climate response to regional radiative forcing during the twentieth century. *Nat Geosci* 2:294–300. <https://doi.org/10.1038/ngeo473>
- Shindell D et al (2012) Simultaneously mitigating near-term climate change and improving human health and food security. *Science* 335:183–189. <https://doi.org/10.1126/science.1210026>
- Shoemaker JK, Schrag DP, Molina MJ, Ramanathan V (2013) What role for short-lived climate pollutants in mitigation policy. *Science* 342:1323–1324
- Stohl A et al (2015) Evaluating the climate and air quality impacts of short-lived pollutants. *Atmos Chem Phys* 15:10529–10566. <https://doi.org/10.5194/acp-15-10529-2015>
- Swedish Board of Agriculture (2012) *Ett klimatvänligt jordbruk 2050*, report 2012:35
- Tanaka K, Peters GP, Fuglestedt JS (2010) Multicomponent climate policy: why do emission metrics matter? *Carbon Manag* 2:191–197
- United Nations Environment Programme (UNEP), World Meteorological Organization (WMO) (2011) *Integrated assessment of black carbon and tropospheric ozone: summary for decision makers*. <https://wedocs.unep.org/rest/bitstreams/12809/retrieve>. Accessed 17 Dec 2018
- Zaelke D, Andersen SO, Borgford-Parnell N (2012) Strengthening ambition for climate mitigation: the role of the Montreal protocol in reducing short-lived climate pollutants. *Rev Eur Commun Int Environ Law* 21(3):231–242. <https://doi.org/10.1111/reel.12010>

**Publisher's note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.