



# First report of *Alternaria* black spot of rose caused by *Alternaria alternata* in China

Peihong Fang<sup>1</sup> · Shaochuan Shi<sup>1</sup> · Xintong Liu<sup>1</sup> · Zhao Zhang<sup>1</sup>

Received: 21 June 2019 / Accepted: 5 September 2019 / Published online: 12 September 2019  
© Società Italiana di Patologia Vegetale (S.I.Pa.V.) 2019

**Keywords** *Rosa hybrida* · *Alternaria* black spot · *Alternaria alternata* · Temperate regions

During fall of 2017, leaf black spot symptoms were observed on rose (*Rosa hybrida*) plants in the campus of China Agricultural University, Beijing, China. Subcircular black spots evolved in leaf discoloration and necrosis, seriously reducing the aesthetic value of rose. Small pieces of diseased tissues were transferred to potato dextrose agar (PDA) plates and incubated at 22 °C. The colonies initially appeared white and then turned dark brown. The conidia ( $n = 50$ ) were ellipsoidal or ovoid and measured 11.9 to 27  $\mu\text{m} \times 7.5$  to 5.1  $\mu\text{m}$  with 0 to 2 longitudinal and 1 to 5 transverse septa. The morphological characters matched those of *Alternaria* spp. (Simmons 2007). For molecular identification, DNA was extracted from the mycelium of a purified single colony, amplified with universal primers ITS4/5 and sequenced. The resulting 515 bp nucleotide ITS segment (MH358396) showed 100% homology with the GenBank accession MF614038.1 of *A. alternata*. Additionally, two nuclear protein coding genes EF-1 $\alpha$  (MH475292) and beta-tubulin 1a/b (MH475293) were sequenced and the sequences all showed 99% identity with those of *A. alternata* strains (MF741188.1 and KF728753.1, respectively). To confirm pathogenicity, we harvested conidia from monoconidial culture on PDA. The conidia were then suspended to a final concentration of  $10^6$  per ml and inoculated artificially on the abaxial surface of healthy detached leaves of *R. hybrida* ‘Samantha’. After 3 days, circular or oval and black disease spots were visible

and the fungus was consistently re-isolated. Previously, *Alternaria* black spot of *Rosa* sp. caused by *A. alternata* has only been recorded in tropical and subtropical regions (South Asia and South Pacific countries; Farr and Rossman 2018). This is the first report of *A. alternata* causing black leaf spot on the *Rosa* genus in temperate regions (East Asia).

## Compliance with ethical standards

**Conflict of interest** The authors declare that they have no conflict of interest.

**Research involving human participants and/or animals** The authors declare that no human participants and animals were involved in this study.

## References

- Farr DF, Rossman AY (2018) Fungal databases, U.S. National Fungus Collections, ARS, USDA. Retrieved from <https://nt.ars-grin.gov/fungal-databases/>. Accessed 26 Apr 2018
- Simmons EG (2007) *Alternaria: an identification manual*. CBS Fungal Biodiversity Centre, Utrecht

**Publisher's note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

✉ Zhao Zhang  
zhangzhao@cau.edu.cn

<sup>1</sup> Beijing Key Laboratory of Development and Quality Control of Ornamental Crops, Department of Ornamental Horticulture, College of Horticulture, China Agricultural University, Beijing 100193, China