

CHAPTER 23

RESPIRATORY DISORDERS

MARTIN L ENGMAN

RODNEY C RICHIE

Introduction	513
Normal Pulmonary Function	513
Arterial Blood Gases Analysis	514
Pulmonary Function Testing	516
Airways Disorders	522
Interstitial and Infiltrative Lung Disease	531
Occupational Lung Disease	536
Immunologic Lung Disease	541
Pulmonary Hypertension	544
Sleep Apnea	545
Pleural Disease	548
Infectious Respiratory Disease	549
Special Clinical Problems	550
References	551

INTRODUCTION

Lung disorders are common in the general population. Chronic bronchitis affects 10–25 percent of the US population, asthma 5 percent, and emphysema about 3 percent. In 1992 these chronic obstructive pulmonary diseases were the fourth leading cause of mortality in the USA, accounting for more than 91 800 deaths.¹ Pneumonia is the sixth leading cause of death in the USA.² It has been estimated that 2–4 million cases of community-acquired pneumonia occur in the USA each year and that as many as 20 percent of these require hospitalization. Occupational lung dis-

ease, cystic fibrosis, hypersensitivity pneumonitis and other respiratory disorders affect many others. Accordingly, life insurance applications from persons with a variety of respiratory disorders are frequently encountered, and the underwriter and medical consultant are called on with some regularity to render assessments of the mortality risk associated with various pulmonary impairments.

NORMAL PULMONARY FUNCTION

Although the lung serves many functions (mechanical filtration of blood, metabolic, immunologic etc.), the main function of the lung

is to support cellular respiration by providing an interface that will allow oxygen to move from ambient air into the vascular system in order that it may be delivered to the tissues, and to allow carbon dioxide, the by-product of cellular respiration, to be excreted back into the air.

Ventilation is the mechanical process by which ambient air is caused to move through the tracheo-bronchial tree into the alveolar spaces and back out again. The tracheo-bronchial tree is a system of branching airways which divide in an asymmetric and dichotomous manner. There are approximately 16 generations of airways from the trachea to the terminal bronchioles. The terminal bronchioles comprise the last generation of conducting airways leading to the terminal respiratory units. The terminal respiratory units, or acini, each consist of approximately three generations of respiratory bronchioles and alveolar ducts, which empty into alveolar sacs. No gas exchange occurs in the conducting airways. It is for this reason that these conducting airways are referred to as the 'anatomic dead space'. Beginning with the respiratory bronchioles, however, alveolar out-pouchings begin to appear from airway walls. The number of these alveolar out-pouchings increases with subsequent airway branching until the airway walls are composed entirely of alveolar openings. At this point the airways are called alveolar ducts. Thus respiratory bronchioles and alveolar ducts have the dual roles of air conduction and gas exchange. Alveolar ducts then open into alveolar sacs which consist of complex geometric clusterings of interconnected alveoli.

Each alveolus is surrounded by a network of capillaries. It is within these alveolar-capillary units that gas exchange occurs through the process of diffusion: oxygen moving down a concentration gradient from the alveolar spaces, across alveolar capillary membranes into the pulmonary capillaries; and carbon dioxide moving down a concentration gradient in the opposite direction.

Ventilation is under the control of the cerebral cortex, brainstem and peripheral chemoreceptors. Although ventilation can be volitionally modified to accommodate such activities as speech, it is also under autonomic control by the respiratory centers located in the pons and medulla. In addition, chemoreceptors located in the medulla and caro-

tid bodies respond to changes in pH (medulla), and $P_a\text{CO}_2$ and $P_a\text{O}_2$ (carotid bodies). Even a slight decrease in pH or increase in $P_a\text{CO}_2$ above the normal range, or a reduction in $P_a\text{O}_2$ below 60 torr will markedly increase ventilation. Conversely, significant reductions in the $P_a\text{CO}_2$, provided pH and $P_a\text{O}_2$ remain normal, will result in a diminution (hypopnea) or pause (apnea) in ventilation until the $P_a\text{CO}_2$ is again normalized.

ARTERIAL BLOOD GASES ANALYSIS

Directly measured arterial blood gas parameters include measurements of arterial hydrogen ion concentration (pH), arterial carbon dioxide concentration (partial pressure of carbon dioxide in arterial blood or $P_a\text{CO}_2$) arterial oxygen concentration (partial pressure of oxygen in arterial blood or $P_a\text{O}_2$) and the degree to which hemoglobin is fully loaded or saturated with oxygen ($S_a\text{O}_2$).

Normal arterial pH is maintained within a narrow range (7.35–7.45) through a complex feedback loop involving ventilatory and metabolic adjustments that result in hyperventilation to lower $P_a\text{CO}_2$ and the retention of HCO_3 by the kidneys during periods of acidosis. Conversely alkalosis will result in hypoventilation and CO_2 retention and cause increased renal HCO_3 excretion.

Normal $P_a\text{CO}_2$ values fall within the range of 35–45 torr. An elevated $P_a\text{CO}_2$ level (hypercarbia) is the result of alveolar hypoventilation either because of a reduction in total ventilation or because of an increase in dead space ventilation at the expense of alveolar ventilation. Normal $P_a\text{O}_2$ values vary inversely with age. For normal subjects aged 20–80 $P_a\text{O}_2$ values determined while seated and breathing ambient air at 1 atmosphere pressure range from 100 to 80 torr.

The term hypoxia refers to the situation in which the $P_a\text{O}_2$ is below the normal range. Sometimes the term is also used to refer to tissue hypoxia, a condition in which cells are deprived of an adequate supply of oxygen either because of low oxygen content in arterial blood or because of cardiac or arterial insufficiency. It should be remembered that most oxygen in blood is bound to hemoglobin in the form of oxyhemoglobin and that only a small fraction is dissolved in the plasma. However, the concentration of oxygen dissolved in

plasma (P_aO_2) is one of the three factors that determines the amount of oxygen that can be bound by hemoglobin, the other factors being the binding affinity of hemoglobin for oxygen and the concentration of hemoglobin in blood. Hypoxemia is the term that refers to low arterial blood oxygen content either due to a low P_aO_2 , a low hemoglobin-oxygen binding affinity or anemia.

It is sometimes useful to classify pulmonary insufficiency into two general types. In Type 1 or hypoxic pulmonary insufficiency, arterial P_aO_2 levels are reduced. If the degree of hypoxia is mild, ventilation may not increase and P_aCO_2 levels may be normal. However, reductions in P_aO_2 levels below 60 torr usually result in compensatory hyperventilation and a reduction in P_aCO_2 . In Type 2 or hypercapnic-hypoxic respiratory insufficiency, alveolar hypoventilation produces elevated alveolar concentrations of carbon dioxide and reduced alveolar concentrations of oxygen which, in turn, result in the situation in which P_aCO_2 is increased despite the presence of arterial hypoxemia. Although this classification is somewhat artificial due to the occurrence of mixed forms of respiratory insufficiency, it has proven useful in helping to identify the underlying disease process (see Table 23.1).

Risk Selection

The relative mortality risk of persons with Type 1 respiratory insufficiency varies greatly, depending

on the nature of the underlying disorder and the severity of hypoxemia. However, the presence of any degree of hypoxemia is a cause for concern and should prompt an investigation into the underlying etiology.

A severe degree of hypoxia ($P_aO_2 < 60$ torr) with or without any degree of hypercapnia ($P_aCO_2 > 45$ torr, Type 2 respiratory insufficiency) indicates the presence of a life-threatening disorder. In the 1970s the British Medical Research Council sponsored randomized trials of the effect of supplemental oxygen therapy in hypoxemic patients ($P_aO_2 < 60$ torr) who had clinical evidence of pulmonary hypertension due to chronic obstructive pulmonary disease (COPD). The mean ages for men and women in these trials were 58 and 56 respectively. Patients receiving no supplemental oxygen had an annual death rate of 30 percent compared with 18 percent for patients who received supplemental oxygen for 15 hours per day. In a complementary study in North America, the Nocturnal Oxygen Therapy Trial demonstrated that hypoxemic COPD patients (mean age 65) who received 12 hours of supplemental oxygen per day were found to have an annual death rate of 20 percent compared with an annual death rate of about 11 percent for patients who received oxygen for at least 19 hours per day.³

Therefore, although supplemental oxygen therapy markedly improved survival in patients with COPD who were hypoxemic, even with

Table 23.1. Causes of respiratory insufficiency.

Type 1 (hypoxic)	Type 2 (hypoxic-hypercapnic)
Alveolar process	Neurologic
Pneumonia	Primary alveolar hypoventilation
Pulmonary edema	Sedation
Pulmonary hemorrhage	Cervical cord injury
Pulmonary alveolar proteinosis	Phrenic nerve injury
Atelectasis	Respiratory muscle weakness
Emphysema	Upper airway obstruction
Interstitial process	
Pulmonary fibrosis	
Pulmonary vascular process	
Thromboembolic	
Right-to-left shunt	

supplemental oxygen their relative death rates were elevated several thousand-fold compared with expected death rates. In more recent studies, subjects with COPD who require long-term oxygen therapy have been reported to have mortality rates of 18–22 percent over 18–24 months of follow-up.^{4,5} Patients having hypoxia due to interstitial lung disease have even worse survival.⁶

PULMONARY FUNCTION TESTING

Pulmonary function testing is a valuable aid in determining the cause of such symptoms as cough and dyspnea, as well as gauging the severity of the pulmonary disease process and estimating prognosis. Many types of pulmonary function tests exist. These include tests designed to measure lung volume, airflow, gas exchange, respiratory muscle strength, ventilatory drive and ventilation-perfusion matching (*see* Table 23.2).

Table 23.2. Pulmonary function tests.

Parameter	Measurement
Lung volume	TLC by plethysmography, gas dilution, planimetry and static lung volume determinations
Airflow	Volume-time curve (timed vital capacity), flow-volume curves
Bronchial reactivity	Specific or non-specific (histamine or methacholine) bronchial provocation challenges
Alveolar-capillary surface area	Carbon monoxide diffusion capacity by steady state (DLCO _{ss}) or single breath (DLCO _{sb}) methods
Respiratory muscle strength	Maximum inspiratory and expiratory pressures
Ventilatory drive	Ventilatory response to induced hypercarbia and hypoxia
Adequacy of ventilation	Arterial carbon dioxide concentration (P _a CO ₂)
Ventilation-perfusion matching	Arterial oxygen concentration (P _a CO ₂), ventilation/perfusion pulmonary scintigraphy

Lung Volume Measurements

Total lung volume, more properly referred to as total lung capacity (TLC), is the total amount of

Table 23.3. Lung volumes.

Measurement	Abbr- eviation	Definition
Tidal volume	TV	Amount of air flowing in and out of the lung during quiet normal breathing
Residual volume	RV	Amount of air remaining in the lung after maximal exhalation
Expiratory reserve volume	ERV	Amount of air that can be maximally exhaled from the end of a normal tidal exhalation
Inspiratory reserve volume	IRV	Amount of air that can be maximally inhaled from the end of a normal tidal inhalation
Functional residual capacity	FRC	Sum of ERV and RV; this is the amount of air remaining in the lung at the end of a normal tidal exhalation
Inspiratory capacity	IC	Sum of TV and IRV; this is the amount of air that can be maximally inhaled from the end of a normal tidal exhalation
Vital capacity	VC	Amount of air than can be maximally exhaled following a maximal inhalation or, alternatively, the amount of air that can be maximally inhaled from the end of a maximal exhalation. Also the sum of IC and ERV.
Total lung capacity	TLC	Sum of VC and RV; the total amount of air in the lungs at full inspiration; also the sum of IRV + TV + ERV + RV, or the sum of IC + FRC etc.




Fig. 23.1. Lung volumes. (See Table 23.3 for key to abbreviations.)

air contained within the tracheo-bronchial tree and alveolar units at the point of maximal inhalation. TLC can be further subdivided into four and three other capacities (technically a capacity is the sum of one or more volumes (see Fig 23.1 and Table 23.3.)).

Disease processes that result in a reduction in TLC are termed restrictive diseases. In general, restrictive diseases are those that primarily affect the alveoli, interstitium, pleura, chest wall or the respiratory muscles (see Table 23.4). Since disease processes that result in reductions in TLC usually result in proportionate reductions in the other lung volume and capacity measurements, VC measurements are often used to substitute for TLC measurements. Thus a reduced vital capacity usually indicates the presence of a restrictive lung disease.

Table 23.4. Examples of restrictive lung diseases.

Interstitial lung disease	Disorders of the chest wall
Idiopathic pulmonary fibrosis	Rib fractures
Sarcoidosis	Kyphoscoliosis
Pneumoconioses	Spondylosis
Pulmonary edema	Obesity
Pleural disease	Neuromuscular disorders
Pneumothorax	Spinal cord injury
Pleural effusion	Myasthenia gravis
Empyema	Myotonic dystrophy
Fibrothorax	Anterior horn cell disease

Occasionally severe airflow obstruction may result in an exception to the rule that reductions in VC are paralleled by proportionate reductions in TLC. Severe airflow obstruction may cause air to become trapped within the lung during exhalation. This may result in an increase in RV and a corresponding reduction in VC. In this situation a reduced VC measurement alone might suggest the presence of restrictive lung disease. However, if TLC is measured, the true nature of the impairment will become clear: the reduction in VC will be seen to have occurred due to an increase in RV as a result of air trapping. At times, airflow obstruction and air trapping are sufficiently severe to cause a marked increase in RV and thereby an increase in TLC.

Airflow Measurements

Dynamic spirometry, also called flow related spirometry, measure rates of airflow. Both inhalation and exhalation flow rates can be measured. Typically these tests are performed by asking the subject to inhale fully and then forcibly exhale until RV is reached, at which time the subject is asked to inhale again as rapidly and completely as possible. The volume of air inhaled and exhaled is recorded as a function of time and plotted as a volume-time curve, also known as timed vital capacity. Flow rates are then calculated by dividing the volume of air exhaled (or inhaled) by the time required for that exhalation (or inhalation) (see Fig. 23.2.)

The amount of air exhaled during the first second of a forced expiratory maneuver, the FEV₁, is the most commonly reported




Fig. 23.2. Normal volume-time curve (timed vital capacity curve).

measurement of airflow. Another commonly reported measurement of airflow is the maximum mid-expiratory flow rate (MMEF). The MMEF has also been termed the mid-flow or the FEF_{25-75} (the forced expiratory flow between 25 percent and 75 percent of the vital capacity). This is the rate of airflow between the points at which 25 percent and 75 percent of the vital capacity are exhaled. It was previously thought that a reduction in the mid-flow in the presence of a normal or nearly normal FEV_1 was indicative of obstruction occurring in the small (< 2 mm diameter) airways. It is now believed that isolated reductions in mid-flow rates indicate mild airflow obstruction and are not specific for disease localized to the small airways.

The incorporation of microprocessors into spirometers has made it possible for instantaneous flow rates to be determined in real time during the forced VC maneuver and for these instantaneous flow rates to be plotted against lung volume. The resulting curve, termed a flow-volume loop, gives the same information contained in a volume-time curve but displayed in a manner that facilitates the visual interpretation of the spirogram. Representative examples of flow-volume loops are shown in Figures 23.3 and 23.4.

It should be recognized that flow rates are dependent upon lung volume: as lung volume decreases so do flow rates. Therefore, impairments that cause a reduction in lung volume




Fig. 23.3. Normal flow-volume loop.

(restrictive disease) and impairments that produce a limitation in airflow (obstructive disease) will cause reductions in FEV_1 . Accordingly, a reduction in FEV_1 is not sufficient in itself for determining whether the process responsible for that reduction is primarily restrictive or obstructive in nature. Rather, a restrictive process is indicated when the FEV_1 and FVC (or better, the TLC) are reduced to a similar degree. The presence of an obstructive process is indicated when




Fig. 23.4. Examples of flow-volume loops in various disease states: A, upper airway obstruction; B, mild asthma; C, severe restrictive disease; D, severe COPD.

Interpretation

Reductions in flow rates and volumes are usually expressed in terms that compare a given subject's flows and volumes with the flows and volumes predicted for a normal individual of the same age, height and sex. Typical criteria used to categorize the degree of pulmonary impairment are shown in Table 23.6.⁷

Various sets of equations to predict normal pulmonary function exist. Most sets of equations predicting FEV₁ and FVC for normal men and women are linear functions of height and age and take the general form [(a × height) − (b × age) + c] where a, b and c are constants. The constants used in these equations have been derived empirically by studying populations presumed to consist of individuals having normal pulmonary function (see Tables 23.7–9). The degree to which these study populations represent individuals who truly have normal pulmonary function varies significantly. One of the earlier studies consisted of male patients in Veteran's Administration hospitals. Because cigarette smokers were included in this study, as well as for other reasons, the predicted values for FVC and FEV₁ derived from this study are substantially lower than those derived from other studies that excluded subjects reporting any history of tobacco use, pulmonary disorders or current respiratory symptoms.

the FEV₁ is reduced to a greater degree than the FVC. This is measured by the FEV₁/FVC ratio, also known as the FEV₁ percent (see Table 23.5).

Table 23.5. Ventilatory defect patterns.

	TLC	FVC	FEV ₁	FEV ₁ /FVC
Restrictive defect	Decreased	Decreased	Decreased	Normal or sometimes slightly increased
Obstructive defect	Normal or sometimes slightly increased	Normal or decreased ^a	Decreased	Decreased or normal ^a

^a Severe airflow obstruction may produce air trapping and cause a reduction in the FVC and a corresponding preservation in the FEV₁/FVC ratio.

Table 23.6. Classification of ventilatory impairment severity.

Degree of impairment	FVC		FEV ₁		FEV ₁ /FVC
No impairment	≥ 80% predicted	and	≥ 80% predicted	and	≥ 70%
Mild impairment	60–79% predicted	or	60–79% predicted		
Moderate impairment	51–59% predicted	or	41–59% predicted		
Severe impairment	≤ 50% predicted	or	≤ 40% predicted		

Table 23.7. Studies used to derive equations predicting normal pulmonary function.

	Kory ⁸	Morris ⁹	Crapo ¹⁰	Knudson ¹¹	Enright ¹²
Year	1961	1971	1981	1982	1993
Number of subjects		988	251	901	288
% males	52%	50%	36%	39%	
Age range	Under 25	16%	7%		0%
	25-60	76%	66%		0%
	60-70	6%	14%		57%
	Over 70	2%	14%		43%
Population	Veterans	Mormons (55%) Seventh Day Adventists (24%)	Mormons (90%)		Elderly urban dwellers
Smokers excluded?	No	Yes	Yes	Regular smokers excluded	Yes
Methods used to exclude	Unknown	Questionnaire	Questionnaire Chest X-ray Physical exam	Questionnaire	Questionnaire
Height measured	Without shoes	Without shoes			
Spirometer	Stead-Wells	Water seal	Pneumotach ATS standards	Water seal ATS standards	ATS standards
Measurements	FEV ₁ , FVC,	FEV ₁ , FVC, FEF ₂₀₀₋₁₂₀₀ FEF ₂₅₋₇₅	FEV ₁ , FVC, FEV ₁ /FVC FEF ₂₅₋₇₅	FEV ₁ , FVC FEV ₁ /FVC FEF ₂₅₋₇₅ , \dot{V}_{50} , \dot{V}_{75}	MIP, MEP

Table 23.8. Prediction equations for FEV₁. FEV₁ = H₁ × Ht (in inches) - A₁ × age (in yr) - F₁.

Study	H ₁	A ₁	F ₁	SEE
Kory				
Males	0.0940	0.0280	1.5900	
Females	0.0711	0.0210	0.0870	
Morris				
Males	0.0920	0.0320	1.2600	0.550
Females	0.0890	0.0250	1.9320	0.470
Crapo				
Males	0.1052	0.0244	2.1900	0.486
Females	0.1052	0.0244	2.1900	0.326
Knudson				
Males	0.1689	0.0292	6.5147	
Females	0.0785	0.0201	1.4050	
Enright				
Males	0.0960	0.0271	1.7300	0.505
Females	0.0714	0.0325	0.0900	0.300

Table 23.9. Prediction equations for FVC. FVC = H × Ht (in inches) - A × age (in yr) - F.

Study	H	A	F	SEE
Kory				
Males	0.1321	0.0220	3.6000	
Females	0.1041	0.0180	2.6900	
Morris				
Males	0.1480	0.0250	4.2410	0.740
Females	0.1150	0.0240	2.8250	0.520
Crapo				
Males	0.1524	0.0214	4.6500	0.644
Females	0.1247	0.0216	3.5900	0.395
Knudson				
Males	0.2144	0.0298	8.7818	
Females	0.1085	0.0174	2.9001	
Enright				
Males	0.1440	0.0206	4.3700	0.505
Females	0.0927	0.0330	0.7000	0.444

Because underwriting criteria for pulmonary function are often based on a percent reduction in FEV₁ and FVC compared with normal predicted values, it is desirable that the set of equations used to specify normal values for pulmonary function, or the tables constructed from these equations, be based on data that directly relate reductions in FEV₁ and FVC to mortality risk. If other sets of normal values are chosen, mortality experience may differ from expected unless underwriting actions are adjusted accordingly.

Risk Selection

The spirometer was invented in 1846 by John Hutchinson who believed that the maximum volume of air that a subject could exhale was a reliable indicator of the risk of subsequent death. Such was his confidence in this belief that he coined the term vital capacity for this measurement and offered his device to the insurance industry of London.¹³ Numerous studies over the years have proven Dr Hutchinson's belief to be correct. However, it now appears that reductions in FEV₁ may be more predictive of subsequent mortality than reductions in VC or the ratio FEV₁/FVC.¹⁴

The Copenhagen City Heart Study followed 12511 subjects for a mean of 6.5 years. Using a proportionate hazards model to predict the risk of cardiovascular death and fatal or non-fatal myocardial infarction (MI), the percent predicted FEV₁ and percent predicted FVC were shown to be significantly related to cardiovascular mortality and the risk of fatal MI. Subjects with a FEV₁ or a FVC less than 60 percent of predicted had a risk of death twice as great as those having values for FEV₁ or FVC greater than 80 percent of predicted.¹⁵

In a study of 2718 British men followed for 20–25 years, the ratio of FEV₁ to the cube of the standing height (FEV₁/H³) measured at the time of entry into the study was shown to predict the risk of death from chronic obstructive lung disease, lung cancer, non-neoplastic respiratory causes, vascular disease and all-cause mortality.¹⁶

Data on 3133 men and women participating in the Framingham studies indicated that the ratio FEV₁/Ht² was inversely related to mortality in

men and women under age 70 and in women older than age 70.¹⁷

In a 24-year follow-up study of 874 men enrolled in the Baltimore Longitudinal Study on Aging, reductions in percent predicted FEV₁ predicted mortality over 24 years of follow-up. After adjusting for the effect of age, smoking, education and race, the relative mortality risk for individuals having an FEV₁ 70–80 percent of predicted compared with individuals having an FEV₁ 100–110 percent of predicted was 125 percent.¹⁸ Individuals who had rapidly declining FEV₁ values had up to a fivefold increase risk of cardiac deaths compared with those who had the lowest rate of FEV₁ decline.¹⁹

Reductions in FEV₁ have also been shown to predict deaths due to lung cancer. For men participating in the Multiple Risk Factor Intervention Trial (MRFIT), lung cancer death rates were increased sevenfold for persons having FEV₁ values in the lowest quintile (≤ 2670 ml) compared with the highest quintile (≥ 3749 ml) between the third and tenth years of follow-up. The authors report that this relationship was not weakened after adjustment for smoking history.²⁰

In an insured lives study performed by Lincoln Re, excess death rates became significantly elevated even with what would be considered mild impairments in FEV₁ (FEV₁ between 60–79 percent of predicted).²¹

However, care should be taken when interpreting timed VC studies to ensure that the reported results reliably reflect the subject's true pulmonary function status. Pulmonary function testing is fraught with potential for error, including errors resulting from faulty equipment, improper spirometer calibration, those introduced by inexperienced technicians and the failure of the subject to inhale to full lung capacity and to completely exhale with maximum force. The importance of this last point cannot be emphasized too strongly: performing a proper timed VC maneuver requires a skilled technician and a capable, cooperative and highly motivated subject. Significant error will result if incorrect instructions are given to the subject, if technical errors are made or if the subject exerts less than maximum effort.

Reviewing the actual timed VC curves or flow-volume loop tracings will aid in identifying many

sources of error. Fortunately most common errors (with the exception of calibration errors) tend to produce falsely low values for FVC and FEV₁. Accordingly, if errors other than calibration are suspected, it may be prudent to consider reported values for FVC and FEV₁ to represent the lower bound of what is likely to be the subject's true pulmonary function.

AIRWAYS DISORDERS

Airways disorders encompass the following diseases: asthma, asthmatic bronchitis, chronic bronchitis, emphysema, bronchiectasis, bronchiolitis and cystic fibrosis.

Asthma

Asthma is a disease characterized by airway inflammation and bronchospasm, which results in reversible airflow obstruction. Although our understanding of the pathophysiology of asthma is incomplete, it is clear that airway inflammation plays a central role in the pathogenesis and perpetuation of asthma by causing the production and release of mediators and mitogens that produce structural remodeling of the lung. Over time, this remodeling may produce fixed airway narrowing and airway hyperresponsiveness that is refractory to treatment.²² Accordingly, early and chronic treatment with inhaled anti-inflammatory agents (inhaled corticosteroids, chromolyn sodium or nedocromil) is now recommended for essentially all asthma patients, with perhaps the exception of those having only the mildest forms of asthma and those who can easily avoid exposure to identifiable allergens.^{23,24}

In the past, asthma has been classified as intrinsic or extrinsic depending on whether airway hyperresponsiveness occurred in an atopic individual (i.e. an individual in whom an allergy to airborne substances could be demonstrated). Atopic individuals with bronchial hyperresponsiveness, elevated IgE levels and peripheral blood eosinophilia were said to have extrinsic asthma; those without atopic features were said to have intrinsic asthma.

No difference in histology or response to corticosteroids has been demonstrated between sub-

jects with intrinsic and extrinsic asthma, and these terms are used less commonly now than in the past. Indeed some investigators believe that if the causal stimuli were known for every patient, all cases of asthma would be extrinsic.²⁵

Today, the causes of asthma are often discussed in terms of 'inducers' and 'triggers'. Inducers are those substances that cause airway inflammation. These include allergens, viral and mycoplasma infections and exposure to noxious gases such as chlorine. Triggers are those stimuli that provoke bronchoconstriction and symptoms in asthmatics but that do not cause airway inflammation. Examples of triggers include exercise, cold air, emotional upset, laughter and inhaled irritants such as perfumes and second hand cigarette smoke. Long-term exposure to environmental tobacco smoke has also been suspected to be an inducer of inflammation in susceptible persons, such as young children.^{26, 27}

The prevalence of asthma has increased worldwide over the past 15 years, as have asthma death rates.^{28,29} Data for the USA indicate that asthma mortality rates reached a nadir in 1977 but have progressively increased since then.³⁰ The reasons for this increase are uncertain, but rising levels of urban air pollution, indoor air pollution with second hand tobacco smoke, overuse of inhaled bronchodilators, changing social and economic factors and limited access to health care have all been proposed as possible causes.³¹⁻³⁴

It is noteworthy, however, that the increase in asthma hospitalization rates and mortality rates have not been evenly distributed across all localities in the USA. Rather, there has been a disproportionate increase in asthma morbidity and mortality in specific counties, among specific social and ethnic groups, and in specific zip code areas within large cities such as New York and Boston.³⁵⁻³⁷ Poverty, lower educational attainment, lack of treatment with inhaled corticosteroids, coexisting alcohol and substance abuse and genetic predisposition have all been postulated as factors to explain this pattern.³⁸

Mortality Studies

In a longitudinal study of 1075 asthmatics between the years 1974 and 1990, all-cause mortality was found to be increased to 240 percent (95

percent confidence intervals: 160–340 percent) compared with age- and sex-matched controls.³⁹ Multiple regression analysis revealed that age, a history of cigarette smoking, the presence of eosinophilia, percent predicted FEV₁ and the degree of reversible airway obstruction were all predictors of death. No association was found between previous hospital admissions for asthma and subsequent death (see Table 23.10)

Table 23.10. Risk factors for asthma deaths.^a

Variable	RR death from asthma	95% CI
Age		
< 39	1.0	
40–69	2.5	0.8–8.1
> 70	8.5	2.2–33.7
Tobacco use		
Lifelong non-smokers	1.0	
< 20 pack-years	2.6	1.0–6.8
> 20 pack-years	5.9	2.3–15.0
Blood eosinophils		
< 450/mm ³	1.0	
> 450/mm ³	7.4	2.8–19.7
FEV₁ % predicted		
> 70%	1.0	
40–69%	4.9	2.0–11.8
< 40%	3.3	0.8–14.5
Reversibility		
15–24%	1.0	
25–49%	2.7	0.9–7.9
> 50%	7.0	2.4–21.0

a Source: Ulrick and Frederiksen.³⁹

In another long-term study of 2499 community residents with asthma living in Olmsted county, Minnesota, overall survival was found to be not significantly different than expected for the general population over a mean of 14 years of follow-up. Survival, however, was found to be less than expected for subjects who were 35 years of age or older when their asthma was diagnosed and in those who had another coexisting lung disease.⁴⁰

In contrast to the relatively favorable mortality rates reported in community studies of asthma are those studies following persons with severe asthma. Marquette *et al.* followed, for periods

ranging from 1 to 75 months, 147 patients with life-threatening asthma who required mechanical ventilation. In-hospital mortality was 16.5 percent. Among the 121 patients discharged from the intensive care unit (ICU), 18 subsequently died, 17 from a new asthma attack. Post-hospitalization mortality was 10.1 percent. Nearly two-thirds of these post-hospital deaths occurred within the year following discharge from the ICU. Smoking and age over 40 year were associated with a higher risk of mortality.⁴¹

The Medical Impairment Study 1983 included a relatively large number of insured lives with a history of asthma. In this study 125 377 policies issued at standard or substandard rates to persons with asthma between the years 1952 and 1976 were traced to 1977 policy anniversaries or death. Only policies on persons free from other impairments were included. Overall cumulative mortality ratios were found to be 118 percent for men and 183 percent for women. For men who were issued policies at standard rates, mortality ratios were 108 percent of expected. Comparable figures for women were 160 percent of expected. For men and women who were issued policies at substandard rates, mortality ratios were 211 percent for men and 281 percent for women. For men mortality ratios were somewhat lower for issue ages 30–39 (178 percent) and 40–49 (196 percent) compared with ages 15–29 (239 percent) or 50–69 (230–237 percent).⁴²

In 1984 Swiss Re reported the results of an asthma mortality study in persons issued insurance between 1958 and 1982. Mortality ratios were 255 percent for men and women classified as having chronic, severe asthma and 105 percent for those classified as having occasional or seasonal asthma compared to standard insured lives expectations.⁴³

In another insured lives study from Lincoln National, policyholders with asthma were found to have an overall mortality ratio of 194 percent compared to standard insured lives expectations.⁴⁴

Selection of Risks

Clinical criteria to classify asthma in terms of severity have been proposed (see Table 23.11). Data from recent studies indicated that mild

asthma may be associated with relatively little excess mortality, especially if access to medical care is readily available, if therapy is directed toward ameliorating airway inflammation and if there is good compliance with the prescribed treatment regimen. Certain subgroups of persons with asthma have a higher risk of death. These appear to include women, persons with severe asthma, persons with a past episode of life-threatening asthma, those with other coexisting lung diseases, persons with psychiatric impairments⁴⁵ and those with alcohol or substance abuse disorders.

The use of inhaled steroids appears not to be a marker for an increased risk of asthma mortality. On the contrary, patients who regularly use inhaled steroids appear to have a significantly improved survival compared to those who do not.⁴⁶ The regular use of oral or parenteral corticosteroids is a marker for asthma of at least moderate severity. The potential complications of

chronic corticosteroid use should also be considered as part of total mortality risk.⁴⁷

Chronic Bronchitis and Emphysema

Chronic bronchitis and emphysema are the two other common forms of COPD.

Chronic bronchitis has been defined on clinical grounds as the presence of a chronic productive cough on most days of the month for 3 months in each year for 2 consecutive years in an individual in whom other causes of cough (such as chronic infection, lung cancer and congestive heart failure) have been excluded. Most cases of chronic bronchitis are due to the chronic inhalation of tobacco smoke or, less commonly, other substances.

The histologic changes seen in chronic bronchitis consist of bronchial gland hyperplasia and hypertrophy, patchy areas of squamous metaplasia in bronchi and an increase in the amount of

Table 23.11. Clinical classification of asthma by severity (Modified from Woolcock and Jenkins⁴⁸).

Symptoms	Score	Bronchodilator Use		Steroid Use		PF or FEV ₁ Variability ^a %	Score
		Use	Score	Use	Score		
Life-threatening	4	< 6×/day	4	Daily oral for > 30 days per yr	4	> 50	4
Nocturnal symptoms or hospitalization or ER visit for asthma within the yr	3	4–6×/day	3	Daily oral < 30 days per yr	3	30–50	3
Symptomatic if not using bronchodilators	2	1–4×/day	2	Daily inhaled	2	20–30	2
Rare wheezing or cough-variant asthma of exercise-induced symptoms, only	1	< 1×/day	1	Occasionally inhaled	1	10–20	1

^a PF or FEV₁ Variability = (best - worst)/best over preceding yr

Scoring: add up scores for each category

	Total score excluding variability scores	Total score including variability scores
Severe	≥ 8	≥ 10
Moderate	4–7	5–9
Mild	1–3	1–4

airway smooth muscle. Bronchioles, the small airways lacking cartilage, show varying degrees of inflammation, mucous plugging, smooth muscle hypertrophy and distortion. Bronchiolar distortion is related to airflow obstruction; however, not all cases of chronic bronchitis have significant degrees of airflow obstruction. Those cases without significant airflow limitation have been termed chronic simple bronchitis.

Emphysema is characterized by destruction of the terminal respiratory units or acini, the gas exchanging portions of the lung. Three general types of emphysema have been described, depending on which portions of the acini have been primarily affected: centriacinar, panacinar, and distal acinar (or paraseptal). Centriacinar emphysema affects primarily the proximal portions of the acini, just beyond the terminal bronchioles. This is the form of emphysema most commonly associated with cigarette smoking. Chronic dust inhalation can produce localized forms of centriacinar emphysema, also known as focal emphysema. In centriacinar emphysema the upper and posterior portions of the lung are most extensively affected.

Pan acinar emphysema involves destruction of the entire acinus. It most often involves the basal portions of the lung to a greater extent than the apices. The lung bases of cigarette smokers often have changes of panacinar emphysema whilst the apices may show changes of centriacinar emphysema. A diffuse severe form of panacinar emphysema is seen in persons with alpha 1 anti-protease deficiency.

Distal acinar (or paraseptal) emphysema is a localized form of emphysema involving the sub-pleural airspaces and the airspaces adjacent to the interlobular septae. It is this form of emphysema that causes the apical bullae that result in spontaneous pneumothoraces often seen in young persons.

Although emphysema is primarily a disease of the acinus and alveolar capillary units, the destruction of elastic tissue causes a loss of pulmonary elastic recoil and results in airway collapse and airflow obstruction.

Emphysema, chronic bronchitis and asthma are often present together to varying degrees. This common occurrence has led to use of the

term chronic obstructive pulmonary disease (COPD). However, although some degree of bronchial hyperreactivity is common in many cases of obstructive pulmonary disease, the term asthma should be reserved for cases where there is complete reversibility of airway obstruction (at least in the early stages of the disease) and that lack the other features of chronic bronchitis and emphysema (see Fig. 23.5).




Fig. 23.5 Relationship of various forms of COPD.

Relationship with Tobacco Smoke

In 1964 the first US Surgeon General's Report on Smoking and Health established an association between cigarette smoking, symptoms of chronic cough and sputum production, and findings of impaired lung function in smokers over the age of 40.⁴⁹ Since then, cigarette smoking has been identified as the major cause of COPD in the USA.⁵⁰

Tobacco smoke causes many pathologic changes in the lungs of smokers, including loss of cilia; proliferation of goblet cells; mucous gland hyperplasia; squamous metaplasia of bronchial mucosa which may progress to bronchogenic carcinoma; inflammation, atrophy and mucous plugging of small airways; destruction of peribronchiolar alveoli; and activation of macrophages and neutrophils.⁵¹

Epidemiologic studies have indicated that on average, lung function, as measured by the FEV₁, normally declines in non-smokers at an average rate of 20–30 ml per year beginning at

about age 25. In smokers, FEV₁ may decline at a somewhat faster rate (25–80 ml per year).^{52–55} Of smokers, 10–15 percent may be particularly susceptible to the effects of cigarette smoke and may lose lung function at a considerably faster rate (e.g. an annual FEV₁ loss on the order of 150 ml per year). At that rate, a susceptible smoker who began cigarette smoking at age 25 may have an FEV₁ under 1 before age 60. It has been suggested that a smoker who has an FEV₁ lower than 1 l below predicted after 20 years of smoking is a susceptible smoker.⁵⁶ Although smoking cessation does not result in any gain in lung function, the rate of loss appears to be reduced to rates comparable to non-smokers (20–50 ml per year).^{51,55}

Some individuals appear to be genetically resistant to the detrimental effects of tobacco smoke. In fact, only 10–15 percent of smokers develop clinically significant COPD. Therefore it has been suggested that if lung function is normal in an individual over the age of 40 who has smoked for more than 20 years, the probability of that individual developing significant obstructive airways disease over the remainder of his lifetime is low.

Although environmental tobacco smoke can trigger asthma in susceptible adults, its role in the development of other forms of COPD is uncertain. Several studies have demonstrated mild degrees of pulmonary function impairment in the children of parents who smoke, a higher incidence of respiratory illnesses, and have implicated environmental tobacco smoke in the development of childhood asthma.

The American Cancer Society's Cancer Prevention Studies I and II (CPS-I and CPS-II) both demonstrated that cigarette smoking is associated with excess deaths in males and female

from COPD as well as from all causes, coronary heart disease, stroke and cancer. CPS-I covered the years 1959–1965 and involved 1 051 038 subjects in 25 states. CPS-II covered the years 1982–1988 and involved 1 083 600 subjects in all 50 states, the District of Columbia, Puerto Rico and Guam. Age-adjusted death rates for smokers and non-smokers were calculated and compared to determine relative and excess death rates. Both studies demonstrated that deaths due to COPD were increased markedly for male and female smokers compared to non-smokers. During the 23-year interval between studies, death rates due to COPD remained stable for non-smokers but increased by 1.5-fold for male smokers and 3.3-fold for female smokers.⁵⁷

Alpha-1 Protease Inhibitor Deficiency

Alpha-1 protease inhibitor (API) was formerly called alpha-1 antitrypsin. It is a protein, normally found in the serum, capable of inhibiting the action of several proteolytic enzymes including trypsin, chymotrypsin, elastase, neutrophilic elastase and collagenases. API is an acute-phase reactant which appears to play a role in inflammation.

The gene coding for API is located on chromosome 14 and is highly pleomorphic: 75 different alleles have been identified. Some alleles are associated with normal levels of normally functioning AP; other alleles are associated with low or absent amounts of API or abnormally functioning API. The amount and kind of API present in an individual is determined by the interaction of the two co-dominant alleles inherited from both parents. The normal API allele is termed M. Other common alleles and their corresponding genotypes and phenotypes are shown in Table 23.12.

Table 23.12. Protease inhibitor genotypes.^a

Genotype	Serum API concentration	Risk of emphysema	
		Non-smokers	Smokers
PI MM	20–48 μ mol (150–350 mg/dl)	Not increased	Increased
PI MZ	12–35 μ mol (90–260 mg/dl)	Not increased	Increased
PI SS	15–33 μ mol (100–240 mg/dl)	Not increased	Increased
PI SZ	8–19 μ mol (60–140 mg/dl)	Slight increase	Increased
PI ZZ	2.5–7 μ mol (20–50 mg/dl)	Increased	Marked increase

^aSource: Snider *et al.*⁵⁸

A threshold protective effect of an API against emphysema in non-smokers is felt to occur at a serum concentration of 11 μmol (80 mg/dl or 35 percent of normal).

It is postulated that API normally protects the lung from damage by neutrophilic elastase and other proteases liberated during inflammatory processes and that an imbalance between proteolytic enzymes and antiproteolytic enzymes, such as occurs in hereditary API deficiency, can result in the destruction of lung tissue and lead to emphysema. It is also felt that cigarette smoking may cause a similar protease-antiprotease imbalance by producing chronic inflammation that leads to an increase in protease activity and to a simultaneous inactivation of API through the action of oxidants.

API deficiency is also associated with other disease states. PI-Z mutations result in aggregation of insoluble PI-Z protein in the endoplasmic reticulum of hepatocytes, eventually producing cirrhosis. In addition, various forms of vasculitis may be seen in API deficiency states.⁵⁹

Diagnosis

Chronic bronchitis and emphysema are diagnosed on the basis of history, physical examination and pulmonary function studies. Shortness of breath is the usual complaint in both chronic bronchitis and emphysema. Persons with emphysema may have few complaints other than the gradual and steady progression of shortness of breath over many years. Chronic bronchitis is diagnosed clinically on the basis of the presence of a chronic cough and sputum production. Findings of lung hyperinflation, wheezing and prolonged exhalation may be present in asthma, chronic bronchitis and emphysema. Cyanosis and signs of right heart failure may also be present in severe cases, but they are more common in chronic bronchitis.

Pulmonary function testing will show airflow obstruction in emphysema, most cases of chronic bronchitis and intermittently in asthma. For this reason, expiratory flow rates alone are not useful in distinguishing between these forms of obstructive airway disease. Because emphysema results in the destruction of alveolar-capillary units, whereas chronic bronchitis and asthma affect primarily the airways themselves, tests of carbon monoxide

capacity, which measure the functioning alveolar-capillary surface area, are often used as an aid to distinguish emphysema from other obstructive airway diseases.

It should be emphasized that the finding of lung hyperinflation on a plain chest radiograph is not a sensitive or specific marker for obstructive airway disease. Lung hyperinflation can be seen in normal individuals as well as in persons with a variety of types of obstructive airway diseases. Adding other radiographic criteria, such as rapid tapering of airways, paucity of vascular markings, peribronchial cuffing and bronchial wall thickening, may result in some increase in specificity. However, plain chest radiography is rarely useful in diagnosing any but the most severe cases of emphysema.⁶⁰

Mortality Studies on COPD

In 1987 Burrows reported the results of a 10-year follow-up study of 45 subjects having COPD and values for FEV₁ less than 65 percent predicted.⁶¹ Butz compared the survival rates reported by Burrows to the 1975–1979 Group Life tables and estimated excess death rates and relative mortality ratios. Over the interval, 0–5 years, the excess death rate was calculated to be 26 per 1 000 per year and the relative mortality ratio 215 percent. For the interval, 0–5 years, the corresponding figures were 67 per 1 000 per year and 385 percent. Over the entire 10-year duration, the mean annual excess death rate was 46 per 1 000 per year and the relative mortality ratio 280 percent.⁶²

One of the largest insured lives studies of the mortality risk associated with COPD is the *Medical Impairment Study 1983* covering the years 1962–1977. Mortality was analyzed separately for chronic bronchitis and emphysema and for males and females. Males with chronic bronchitis who were offered policies at standard rates had a relative mortality ratio of 137 percent (standard mortality = 100 percent based on modified 1965–1970 Basic tables); those rated slightly substandard (up to 175 percent) had a mortality ratio of 239 percent; and those rated moderately substandard (180–250 percent) had a mortality ratio of 281 percent. The overall mortality ratio for substandard male issues was 240 percent. The overall mortality ratio for substandard females (235

percent) was similar to that for males. Males with emphysema who were offered policies at standard rates had a relative mortality ratio of 128 percent; those rated slightly substandard (up to 175 percent) had a mortality ratio of 309 percent; and those rated moderately substandard (180–250 percent) had a mortality ratio of 341 percent. The overall mortality ratio for substandard male issues was 323 percent, the second highest relative mortality in the Medical Impairment Study 1983 after MI.

The number of policies issued on women with COPD were far fewer than the number issued on males. This precluded a detailed mortality analysis. For standard and substandard issues combined, women having chronic bronchitis had a relative mortality ratio of 235 percent. The corresponding figure for women having emphysema was 252 percent.⁶³

Brackenridge and Mills have reported that in a smaller intracompany study of COPD by the Swiss Reinsurance Company, the relative mortality ratio associated with chronic bronchitis compared with standard risks over the same period of observation (1958–1982) was 210 percent. The corresponding figure for emphysema was 235 percent. When emphysema was combined with chronic bronchitis the relative mortality rose to 345 percent.⁶⁴ Brackenridge and Mills have also stated that this adverse effect of emphysema combined with chronic bronchitis was also seen in the study by the Prudential Assurance Company of London where among insureds aged 39–49 at entry the excess death rate was 13.5 per 1000 and the relative mortality ratio 323 percent. Among insureds aged 50 and older the excess death rate was 20.3 per 1000 and the relative mortality ratio 196 percent.⁶⁹

More recently Fessel has reported a study of 1481 European males having emphysema and/or chronic bronchitis who were issued life policies between the years 1956 and 1985.⁷⁰ The average duration per policy issued was slightly more than 7.5 years. Overall, the mortality ratio of this cohort relative to standard insured lives over this duration was 230 percent with an annual average excess death rate of 6.7 per 1 000 per year. Males issued policies between the ages 20 and 44 had a mortality ratio of 190 percent. The mortality ratio

for policies issued between ages 45 and 54 was 235 percent, and for policies issued between ages 55 and 84 the mortality ratio was 265 percent.

Risk Selection

Depending on the severity of airflow obstruction, excess mortality may range from mild to extreme. The results of pulmonary function testing are helpful in stratifying risk. Continuing tobacco use in the presence of obstructive airways disease is associated with continuing loss of lung function at an accelerated pace. The presence of hypoxemia, pulmonary hypertension and cor pulmonale are also adverse prognostic factors.

Bronchiectasis

Bronchiectasis is a condition in which the muscular, elastic and cartilaginous components of the walls of medium or large airways are damaged or destroyed. This results in an abnormal dilatation (ectasia) of the bronchus and impaired mucociliary clearance. Bronchiectasis may result from bronchial obstruction or infection, or be a complication of a systemic disorder (such as cystic fibrosis, primary ciliary dyskinesia (Kartagener's syndrome), allergic bronchopulmonary aspergillosis, IgG deficiency, alpha-1 protease inhibitor deficiency or Marfan's syndrome).

If caused by bronchial obstruction or acute pneumonia, bronchiectasis may be localized to discrete areas of the lung. Some cases of localized bronchiectasis may even resolve. However, it is more common for bronchiectasis to be the result of a diffuse process, to be chronic, persistent, and involve multiple areas of the lungs or be generalized. Impaired mucociliary clearance results in secondary bacterial colonization, infection, and inflammation resulting in further bronchial damage and fibrosis.

Bronchiectasis may be asymptomatic or result in a chronic cough productive of copious amounts of muco-purulent sputum ('wet bronchiectasis'). If bronchiectasis is localized to the upper lobes of the lungs, good bronchial drainage may be preserved and cough may be non-productive ('dry bronchiectasis').

Common complications of bronchiectasis include recurrent pneumonia, lung abscess,

pneumothorax and hemoptysis. Hemoptysis usually follows a mild respiratory infection. It is said to be more often a complication of dry bronchiectasis than of wet bronchiectasis. Usually the hemoptysis is mild and self-limited. However, because bronchial arteries contain blood under systemic pressure, hemoptysis may be massive. Despite this, it has been stated that patients with bronchiectasis rarely die from hemoptysis.⁶⁷

Treatment for bronchiectasis includes the following components: controlling infection with antibiotics, promoting adequate mucociliary clearance, and identifying and treating any underlying disease processes (e.g. immunoglobulin administration for patients having IgG deficiency). Medical management is effective for most patients. Surgical resection should be reserved for patients with focal bronchiectasis who have life-threatening hemoptysis, uncontrollable infection or who are unwilling or unable to take antibiotics. Double lung or heart-lung transplantation is being performed for patients with severe, generalized bronchiectasis, most often that associated with cystic fibrosis.

In the pre-antibiotic era, survival in patients with bronchiectasis was dismal and the quality of life poor. However, with the advent of antibiotics survival has improved dramatically.

The Medical Impairment Study 1983 reported insured lives mortality experience on 5300 life insurance policies issued between 1952 and 1977 on persons with a history of bronchiectasis. Most of the exposure occurred in males (n = 76) and in policies issued standard (n = 26) or mildly (n = 36) or moderately substandard (n = 18). There were 87 deaths in this cohort. For males the standardized mortality ratio compared with insured lives expectations was 132 percent for standard issues, 134 percent for mildly substandard issues and 153 percent for moderately substandard issues. The mortality ratio was higher for males aged 40 or older the time of issue compared with younger males (167 percent vs 81 percent).

Risk Selection

Mortality risk in patients with bronchiectasis is influenced by the extent of the disease; the presence of complications (such as life-threatening hemoptysis); pulmonary function; the coexistence

of asthma, emphysema or chronic bronchitis; and the presence of underlying disease processes such as immune deficiency states and cystic fibrosis. Patients who have localized bronchiectasis, whose overall pulmonary function remains good, who have not had serious complications and who have no other pulmonary disease would be expected to have the best prognosis.

Bronchiolitis and Bronchiolitis Obliterans

Bronchiolitis is the term applied to inflammatory processes primarily affecting the small (< 2 mm diameter) non-cartilagenous airways. A wide variety of toxic, irritative, infectious, immunologic and inflammatory processes can cause bronchiolitis. Infectious bronchiolitis occurs commonly in children. Respiratory syncytial virus, parainfluenza virus, adenovirus, influenza A and B viruses, *Mycoplasma pneumonia* and *Legionella pneumophila* are common etiologic agents. Infectious bronchiolitis is uncommon in adults.

In adults bronchiolitis is more often the result of inhaling toxic gases. Oxides of nitrogen (NO, NO₂, N₂O₄) produced by anaerobic fermentation of silage are responsible for the bronchiolitis of Silo Filler's disease. Fire victim's or firefighter's lung results from the inhalation of a wide variety of toxic fumes liberated by combustion. Depending on the degree of exposure, the resulting illness may be mild with minor cough and wheeze, or severe with resulting respiratory failure, pulmonary hemorrhage, adult respiratory distress syndrome and death. Fortunately most patients recover from acute bronchiolitis.

Bronchiolitis may also be caused by the inhalation of organic dusts. Bronchiolitis is a prominent histopathologic feature (along with the findings of alveolitis, interstitial inflammation and fibrosis) of hypersensitivity pneumonitis.⁶⁸ Examples of hypersensitivity pneumonitis include farmer's lung disease, pigeon breeder's disease and bagassosis.

In some patients bronchiolitis may become chronic. Fibrous scarring of the small airways and exudative plugging of airway lumens may occur. This condition, termed bronchiolitis obliterans, has been reported to follow respiratory

infection, toxic fume inhalation, as a result of medications, as a complication of transplantation or in association with a connective tissue disorder. If the exudative and inflammatory process extends into the alveoli, the condition is termed bronchiolitis obliterans with organizing pneumonia (BOOP).

Bronchiolitis obliterans begins insidiously with a non-productive cough and mild shortness of breath. Over time it may progress to disabling dyspnea and respiratory failure. Chest radiographs may be normal or show a nodular or reticulonodular pattern. BOOP is associated with patchy ground-glass infiltrates in the majority of cases.⁶⁹ Spirometry reveals an obstructive defect in pure bronchiolitis and a restrictive defect in BOOP, usually with an impairment in the carbon monoxide diffusion capacity.⁷⁰ Early treatment with corticosteroids or other immunosuppressants, such as azathioprine or cyclophosphamide, may suppress the inflammatory process and ameliorate or prevent severe obstruction. If severe obstruction and fibrosis are present, however, treatment is less likely to be effective.⁷¹ Corticosteroid treatment appears to be effective in most cases of BOOP.

Risk Selection

Acute infectious bronchiolitis may be fatal in children, especially those under the age of 2. The most common sequela of acute bronchiolitis in children is the development of bronchial hyperreactivity, which may last for years after the initial infection.

There have been few long-term mortality studies on bronchiolitis obliterans. In one study of allogeneic bone marrow transplant recipients, the 5-year survival rate of 35 patients with obstructive airways disease was approximately 18 percent compared with a 54 percent survival at 5 years for 412 concurrent patients with chronic graft-versus-host (GVHD) disease who survived more than 80 days after transplantation but who had normal pulmonary function.⁷²

Cystic Fibrosis (CF)

CF is a multisystem disorder caused by mutations of a gene, termed the cystic fibrosis transmem-

brane conductance regulator (CFTR), located on the long arm of chromosome 7. The major manifestations of CF are obstructive airways disease, recurrent pulmonary infection and exocrine pancreatic insufficiency. Other manifestations may include obstructive azoospermia, pansinusitis, mucoid appendiceal impaction, distal intestinal obstruction or intussusception in children and young adults, salty-tasting skin and hyponatremic dehydration.

CF is inherited in an autosomal recessive pattern. The prevalence of the carrier state is 2–5 percent. The homozygous state has been estimated to occur in 1 in 2500 live births in the Caucasian population. CF is much less prevalent in other populations and it is rare in Asians, native American Indians and black natives of Africa.

The CFTR regulates cAMP activated chloride conductance across membranes. Mutations of the CFTR gene result in the production of a defective CFTR protein and impairment in transmembrane chloride ion conductance. This results in abnormalities in salt and water transport, leading to alterations in the composition of secretions of the lung, pancreas, gastrointestinal tract, sweat glands and other exocrine organs. Respiratory tract secretions become more viscous, and mucociliary transport becomes impaired, predisposing to bacterial colonization of airways with *Staphylococcus aureus*, *Hemophilus influenzae* and *Pseudomonas aeruginosa*. Chronic infection leads to the development of a combination of bronchiectasis, bronchial hyperreactivity, airflow obstruction, cystic dilatation of airspaces and organizing pneumonitis.⁷³

Over 500 mutations of the CFTR gene have been described.⁷⁴ The 70 most common mutations account for more than 90 percent of all cases.⁷⁵ All reported CFTR gene mutations are associated with elevated chloride concentrations in sweat. Many mutations (including the most common mutation, $\Delta F508$, a three base pair deletion at position 508) are associated with pancreatic exocrine insufficiency. The relation between genotype and severity of pulmonary disease has not been as clearly defined. It has been suggested that this lack of tight correlation may be due to the other genetic and environmental influences and the lack of a good system for classifying pulmonary disease severity in CF.⁷⁶

The standard screening test for CF remains the sweat chloride test, in which the concentration of chloride in sweat is measured by ion-selective electrodes or gravimetric methods. Chloride concentrations of 60 mmol/l and typical clinical manifestations are usually diagnostic of CF.⁷⁵

Chest radiographs taken early in the course of the disease demonstrate bronchial thickening, hyperinflation and segmental or subsegmental atelectasis. Later, evidence of bronchiectasis appears. In advanced cases, radiographs show nodular opacities, hyperinflation, cystic airspace dilatation, honeycombing and evidence of right heart failure.

Pulmonary complications of CF include pneumothorax, hemoptysis (which can be minor or life-threatening), respiratory failure, pulmonary hypertension and cor pulmonale.

Current treatment strategies are directed toward maintaining adequate nutrition, controlling respiratory tract infections, promoting mucus clearance and combating inflammation. Lung transplantation for end-stage pulmonary disease is being performed more often.^{77,78} Investigative treatments include employing agents that modulate ion transport, gene therapy and direct introduction of normal CFTR proteins into lung epithelia.

The natural history of CF varies markedly, ranging from death during the neonatal period to relatively asymptomatic and prolonged survival well into the third, fourth and even fifth decades of life. A recent report from the Danish Cystic Fibrosis Center indicates that with aggressive treatment there is a 80.4 percent probability that a newborn child with CF will live until age 45 (confidence interval = 76.5–84.6 percent).⁷⁹

INTERSTITIAL AND INFILTRATIVE LUNG DISEASE

A broad group of diseases of varying etiology falls under this category, ranging from connective tissue disorders, drug induced pulmonary diseases and diffuse alveolar hemorrhage, to idiopathic pulmonary fibrotic disorders. The common feature of these diseases is their ability to produce an inflammatory response in the pulmonary interstitial spaces (i.e. the area between the alveolar base-

ment membrane and the alveolar capillary endothelium).

In general these disorders are characterized by an increase in the elastic recoil of the lungs, a restrictive pattern on pulmonary function testing, no increase in flow resistance, reduction in carbon monoxide diffusing capacity due to diminished alveolar-capillary surface area and reduced values for P_aO_2 . The chest radiograph can reveal a variety of patterns depending on the nature and severity of the disorder. Although clinical history, physical examination, laboratory tests, pulmonary function and imaging studies can provide valuable clues as to the nature of the disorder, definitive diagnosis often requires a tissue diagnosis by transbronchial, transthoroscopic or open lung biopsy.

Idiopathic Pulmonary Fibrosis (IPF)

IPF is known by a variety of other names including cryptogenic fibrosing alveolitis, diffuse interstitial pulmonary fibrosis, chronic interstitial pneumonia and idiopathic interstitial pneumonia. Other names applied to this disorder refer to specific histologic changes seen with light microscopy that may have a somewhat better (desquamative interstitial pneumonia) or worse prognosis (usual interstitial pneumonitis). The use of the term 'Hamman-Rich syndrome' should be restricted to describing a rapidly progressive and fatal form of idiopathic pulmonary fibrosis. The term cryptogenic fibrosing alveolitis, used more commonly outside the USA, is a particularly descriptive term which emphasizes the unknown etiology, the inflammatory cell infiltration of the alveolar septa and alveolar spaces, and the relentless progression to pulmonary fibrosis characteristic of most forms of this disorder.

Although the underlying cause of IPF is unknown, it is clear that chronic inflammation (alveolitis) plays a central role. It is postulated that after some, as yet, undefined initial damage occurs, immune complexes are formed that attract and activate neutrophils, macrophages and fibroblasts resulting in the deposition and/or modification of collagen within the pulmonary interstitium.

IPF is a disorder of middle to old age, and rarely occurs before age 40. Affected individuals

complain of the insidious onset and progression of non-productive cough and dyspnea. Most patients have had symptoms for 2 or more years before the diagnosis is established. In most patients, fine inspiratory crackles are heard at the lung bases upon chest auscultation. Of patients 50 percent may have clubbing of the digits. Chest radiographs typically exhibit a reticular or reticulonodular pattern. In the late stages of the disease, a honeycomb pattern may be seen. Pulmonary function studies show a restrictive defect. Carbon monoxide diffusion is reduced and varying degrees of hypoxemia are present.

The risk of pneumothorax, pneumonia, bronchogenic carcinoma, cor pulmonale, ischemic heart disease and pulmonary embolism are all increased in patients who have IPF. Death is usually due to progressive respiratory insufficiency. Treatment is directed toward treating complications and suppressing inflammation through the use of corticosteroids or other immunosuppressants such as cyclophosphamide, azathioprine, penicillamine or colchicine.

Histologic examination of lung tissue reveals a variety of patterns. Liebow and Carrington⁸⁰ described five patterns of interstitial lung disease which they termed usual interstitial pneumonia (UIP), desquamative interstitial pneumonia (DIP), lymphocytic interstitial pneumonia (LIP), giant cell interstitial pneumonia (GIP) and bronchiolitis obliterans with interstitial pneumonia (BIP).

LIP a rare disorder characterized by a pattern of lymphocytic infiltration of the alveolar interstitium. It must be differentiated from lymphomas, pseudolymphomas and other lymphocytic disorders. Its natural history has not been well studied. Marked improvement or resolution with steroids has been reported; in other cases, progressive fibrosis leading to cor pulmonale occurs. GIP is also rare and has been reported in association with measles and other viral infections. BIP is more commonly referred to as bronchiolitis obliterans with organizing pneumonia (BOOP) (*see above*).

Risk Selection

Carrington *et al.* reported that patients with the DIP pattern had a much better survival than patients with the UIP pattern.⁸¹ They followed

53 patients with UIP (mean age 50.9 years) and 40 patients with DIP (mean age 42.3 years) for an average of 6.7 years (UIP) to 9.1 years (DIP). The majority of subjects were male (33 with UIP, 28 with DIP). Of UIP subjects 38 percent were smokers; 36 percent of DIP subjects were smokers. At 5 years of follow-up 4.8 percent of DIP patients and 44.6 percent of UIP patients had died. At 10 years of follow-up 30.4 percent of DIP patients and 71.5 percent of UIP patients had died.

More recently Erbes *et al.* reported data on 99 patients with IPF. Survival rates at 2, 5 and 10 years were 80 percent, 64 percent and 45 percent. From these data it is possible to calculate that interval mortality ratios were approximately 1000 percent in years 0–2, 650 percent in years 2–5, 400 percent in years 5–10 and 300 percent in years 0–10. The authors also presented data showing that TLC and VC measurements at the time of initial evaluation were predictive of the subsequent course, but that diffusion studies were not.⁸²

Sarcoidosis

Sarcoidosis is systemic inflammatory disorder of unknown etiology. It can affect virtually every organ in the body including the lungs, brain, eyes, heart, liver, kidneys, lymph nodes, bone and skin.

The pathologic hallmark of the disorder is the non-caseating granuloma. However, the presence of non-caseating granulomas is not specific for sarcoidosis: they may be seen in mycobacterial and fungal infections, extrinsic allergic alveolitis, bronchogenic carcinoma, leukemia, Hodgkin's and non-Hodgkin's lymphoma, Crohn's disease, Whipple's disease and pneumoconioses and they are also related to drugs and foreign bodies.⁸³

Although many granulomas resolve over time, some undergo progressive fibrosis. There is abundant evidence that sarcoidosis is an immunologic process which begins by exposure to an as yet unidentified antigen or antigens and is mediated by cellular immune mechanisms involving T cells, interleukins, macrophages, mast cells, neutrophils, natural killer cells and fibroblasts.

Sarcoidosis occurs worldwide. Although it can affect persons of all ages and races, it is more

common in persons under the age of 40, blacks and those of Irish and Scandinavian decent. For unclear reasons, the incidence of sarcoidosis is higher in the winter and early spring.⁸⁴ Although there is evidence from family and twin studies that genetic predisposition plays a role in the development of sarcoidosis, it is unlikely to be due to a single gene mutation.

Sarcoid pulmonary disease has been described in terms of five stages. The International Congress on Sarcoid refers to Stages 0, 1, 2A, 2B and 3. Other investigators refer to Stages 0–4, designating Stage 2B as Stage 3, and Stage 3 as Stage 4. The International Congress staging system will be used here.

Stage 0 is characterized by the absence of abnormal chest radiographic findings. However, active alveolitis is invariably present. Stage 1 is characterized by the presence of bilateral symmetric hilar and paratracheal lymphadenopathy without pulmonary parenchymal involvement. Stage 2A is characterized by pulmonary and hilar lymph node enlargement and interstitial disease. Stage 2B is characterized by interstitial pulmonary disease without lymph node enlargement. Stage 3 is characterized by radiographic evidence of pulmonary fibrosis (diffuse honeycombing, hilar retraction, etc.) and the absence of pulmonary adenopathy. Of sarcoid patients 5–10 percent present with stage 0, 40–50 percent with Stage 1 and 40 percent with Stage 2A. The remaining 5–10 percent present with Stage 2B or 3. Endobronchial sarcoid is common. At times, endobronchial granulomas are large enough to cause airway obstruction.

Pulmonary function studies typically show restrictive defects. The presence of airflow obstruction should suggest the presence of endobronchial obstruction or another coexisting disease. Diffusing capacity is often diminished. At times, pulmonary function studies may be normal despite the presence of radiographic infiltrates. The presence of resting hypoxemia usually indicates advanced disease.⁸⁵

Pulmonary complications include progressive fibrosis, respiratory failure and cor pulmonale. Other uncommon complications include pneumothorax, hemoptysis and pleural effusion.

Extrapulmonary involvement is common. Some of the common extrapulmonary sites of involvement are shown in Table 23.13.

Table 23.13. Sarcoidosis: frequency of occurrence and types of extrapulmonary manifestations.^a

Site	Frequency	Manifestation
Nervous system	5%	Cranial nerve palsy
		Vestibular disturbance
		Peripheral neuropathy
		Meningitis
		Seizures
		Cerebellar lesions
		Spinal cord
Heart	5–10%	Heart block
		Ventricular ectopy
		Dilated cardiomyopathy
		Restrictive cardiomyopathy
		Cor pulmonale
		Sudden death
		Pericardial effusion
Liver	40–70%	Pericarditis
		Hepatic granulomas, usually asymptomatic
		Cholestasis
		Portal fibrosis
Eyes	22%	Uveitis
		Conjunctival and lachrymal granulomas
		Optic neuritis
Reticuloendothelial	28%	Lymphadenopathy
Kidney	15–40%	Splenomegaly
		Renal granulomas (usually asymptomatic)
		Nephrolithiasis
Endocrine	10%	Nephrocalcinosis
	<5%	Hypercalcemia
	10–20%	Hypercalcuria
	60%	Hypopituitarism
	<1%	Diabetes insipidus
Skin	18%	Syndrome of inappropriate ADH secretion
		Erythema nodosum
		Subcutaneous nodules
Musculoskeletal	4%	Lupus pernio
		Bone cysts
		Arthralgias, myalgias
		Polyarthritis

a Source: Sittzbach *et al.*⁸⁶

The diagnosis of sarcoidosis usually rests on typical clinical findings and biopsy evidence of non-caseating granulomas. Common biopsy sources include the skin, conjunctiva, minor salivary glands, lymph nodes and lung (bronchial or transbronchial). Other studies such as cytologic examination of bronchial alveolar lavage samples, serum angiotensin-converting enzyme levels and gallium-67 scanning may be of prognostic value but are not diagnostic.

Siltzbach *et al.*⁸⁶ reported a retrospective series of 1609 patients with sarcoidosis from London, New York, Paris, Los Angeles and Tokyo; 70 percent of patients were under age 40 at the time of presentation, 55 percent were women, 54 percent were white, 22 percent were black and 18 percent were Japanese. Overall approximately 10 percent of patients presented with Stage 0 disease, 40 percent with Stage 1, 35 percent with Stage 2 and 15 percent with Stage 3. Chest radiograph abnormalities resolved in 63 percent of patients with Stage 1, 50 percent of patients with Stage 2 and 18 percent of patients with Stage 3.

Sharma has reported that cardiac involvement occurs in up to 20 percent of patients; however, it is clinically recognized in only 5 percent.⁸⁷ Complete heart block is the most common abnormality. Ventricular ectopic beats and sustained or unsustained ventricular tachycardia are also common. Other conduction defects and supraventricular arrhythmias may also occur. Sudden death due to ventricular tachyarrhythmias or conduction defects is stated to account for 30–50 percent of deaths due to sarcoidosis. Progressive cardiac failure has been estimated to cause 25 percent of cardiac deaths and is the second most common cause of death in patients with cardiac sarcoidosis.⁸⁷

Neurologic involvement may manifest as headache, vertigo, facial weakness, hemiparesis, ataxia, seizures, visual disturbance or dementia. Cranial and peripheral neuropathies, meningitis, and mass lesions in the cerebral hemispheres, basal ganglia, brain stem, cerebellum or spinal cord may occur.

Hepatic granulomas are common in sarcoidosis but are usually asymptomatic. Active liver disease is rare. The two hepatic complications, portal hypertension and cholestasis, are almost exclus-

ively seen in black males over the age of 40 who have extensive pulmonary involvement.^{88, 89}

Corticosteroids remain the primary mode of therapy for sarcoidosis. The indications for corticosteroid treatment have been recently reviewed^{84,90} Patients with Stage 1 and 2A disease who are asymptomatic and have good pulmonary function should be followed without treatment until their disease progresses. Symptomatic Stage 1 and 2A patients, Stage 2A patients who have significant impairment in lung function and Stage 2B and Stage 3 patients should receive treatment. Patients with Stage 3 disease may be candidates for pulmonary transplantation. Cardiac, neurologic, upper airway, ocular involvement and significant hypercalcemia or hypercalcuria should also be treated. Other agents such as chloroquine and methotrexate are sometimes used as an adjunct to corticosteroids, but further studies are needed to define their effectiveness and proper role.

Risk Selection

Siltzbach *et al.*⁸⁶ reported a 6 percent overall mortality in their series: 5 percent as a direct result of sarcoidosis and 1 percent from unrelated causes. Patients who present with Lofgren's syndrome (erythema nodosum, arthralgias and bilateral hilar adenopathy) almost invariably have a good prognosis. Patients with Stage 1 pulmonary disease have a better prognosis than those with Stage 2 who, in turn, have a better prognosis than those with Stage 3. Failure of radiographic resolution over 1 year has been stated to be a poor prognostic sign. Radiographic resolution lasting for 2 or more years has been said to indicate a cure.⁹¹ A worse prognosis may be anticipated for patients over the age of 40 at disease onset, those who have cardiac sarcoid, neurosarcoid, sarcoidosis involvement of more than 3 organ systems, and those who have Stage 3 pulmonary disease. Although the overall prognosis for sarcoidosis is good, as many as 50 percent of patients have residual organ impairment.

Pulmonary Eosinophilic Granuloma

Pulmonary eosinophilic granuloma is one of the entities comprising the Langerhans cell granulomatosis (histiocytosis X). It is a disorder of unknown etiology in which there is a proliferation

of Langerhans cells around bronchioles early in the disease process; it later progresses to involve the interstitium surrounding alveolar units. The radiographic appearance of pulmonary eosinophilic granuloma is that of diffuse reticular, nodular and reticular nodular infiltrates primarily involving the mid-lung zones.

Pulmonary eosinophilic granuloma is a disease of young and middle-aged adults and is more common in heavy smokers. Approximately 25 percent of patients may be asymptomatic at the time of diagnosis, the diagnosis being made in the course of evaluating an abnormal chest radiograph. Other patients present with the symptoms of cough and dyspnea. Pneumothorax is a common complication, occurring in 14 percent of patients. Pulmonary function testing reveals a combination of obstructive and restrictive ventilatory defects. The adult form of eosinophilic granuloma usually remains confined to the lungs. Bone involvement may occur in 5 percent of patients and hypothalamic involvement in 10 percent. This is in contrast to the multifocal forms of Langerhans cell granulomatosis which manifest in childhood and include the entities of Hand-Schuller-Christian disease (bone defects, diabetes insipidus and exophthalmos) and Letterer-Siwe disease (disseminated differentiated histiocytosis).

The prognosis of pulmonary eosinophilic granuloma is variable. Some cases resolve spontaneously and others progress to respiratory failure. Poor prognostic factors include either young or old age, multi-organ involvement, extensive pulmonary disease, low carbon monoxide diffusing capacity and recurrent pneumothoraces. Because of the association of pulmonary eosinophilic granuloma with tobacco use, all patients should quit smoking. Although treatment with corticosteroids and cytotoxic agents may be tried, results are usually unsatisfactory.⁹²

Adult Respiratory Distress Syndrome (ARDS)

ARDS refers to Type 1 respiratory failure resulting from the accumulation of protein and water in the alveolar spaces following acute lung injury. A small amount of water normally moves from the vascular space into the lung interstitium and

alveolar spaces. The amount of water moving from one space to the other is determined by the pulmonary venous pressure and the concentration of protein in the vascular space. The central event in ARDS is damage to the capillary endothelium producing increased permeability to protein and water, which in turn results in the abnormal accumulation of water and protein in the lung. An intense inflammatory response also occurs. ARDS is a disorder affecting the total body rather than a dysfunction limited to the lungs. In fact, some physicians now describe this process as 'increased permeability edema with acute lung injury'.

A large number of insults may cause ARDS, and it is not within the scope of this chapter to review these initiating events. Suffice it to say that the vast majority of ARDS patients are critically ill and spend days to weeks requiring ventilator support. Approximately 40 percent of patients who develop ARDS die, but in the majority of the 60 percent who survive, there is an eventual return towards normal health.⁹³ Of those patients who have residual lung damage, the severity of pulmonary impairment can best be assessed by complete pulmonary function studies. The most common residual of ARDS is an abnormal diffusion capacity (D_LCO). Other ARDS survivors are left with reactive airways disease.⁹⁴ The diffusion capacity (expressed as a percentage of predicted) is inversely correlated with the degree of lung damage (i.e. the lower the diffusion capacity the greater the residual damage).

Of greatest relevance to underwriting an applicant who has survived ARDS is to define as accurately as possible the degree of residual impairment and how that impairment might affect future morbidity and mortality. Lung function improves most rapidly during the first 3–6 months following recovery from acute injury and proceeds more gradually during the next 6 months. Pulmonary function reaches a plateau by 1 year after ARDS onset. Therefore abnormal clinical findings during the first 6 months may prove reversible, but abnormalities that persist for 1 year are unlikely to resolve.⁹⁵ However, pulmonary fibrosis demonstrated on chest radiographs or on computerized chest tomography usually does not reverse.⁹⁶

Pulmonary Alveolar Proteinosis (PAP)

PAP is classified as primary (idiopathic) or secondary (associated with or provoked by another disorder). Secondary PAP may be associated with pulmonary infections (*Nocardia*, *Mycobacteria*, *Pneumocystis carinii*), leukemia, lymphoma and inhalation of dusts and fumes (silica, aluminum, titanium, insecticides).⁹⁷

Primary pulmonary alveolar proteinosis is a disorder of unknown etiology in which proteinaceous material is deposited in alveoli, producing hypoxemia. Little inflammation occurs and pulmonary fibrosis is minimal.

The natural history of idiopathic PAP is variable: one-third of patients are said to die due to progressive hypoxia or superinfection, one-third remain stable and one-third spontaneously improve.⁹⁸ Although various treatments have been tried, including the corticosteroids, potassium iodide, trypsin, acetylcysteine and heparin, only whole lung lavage has proven of benefit. Whole lung lavage is performed under general anesthesia. The patient is intubated with a double lumen Carlens endotracheal tube, and while one lung is ventilated, the other is lavaged 10 or 15 times with saline over a period of 1–2 hours. The other lung is lavaged the next day. This treatment has been shown to be very effective; however, some patients require repeat whole lung lavage at intervals of 6–12 months before the disease becomes quiescent. Although no large series have been reported, the prognosis of patients treated with whole lung lavage is said to be good. In a series of 21 patients with primary PAP followed for periods ranging from 1 to 19 years, a total of 3 patients died, 2 from motor vehicle accidents and 1 from pulmonary fibrosis.⁹⁸

OCCUPATIONAL LUNG DISEASE

Estimates of the rate of development of occupational lung disease in the USA range from 1.25 to 4 per 1000 workers. As most of these estimates come from physicians reporting of occupational illnesses, and as most states do not require such reporting, there is a general feeling by experts in the field that these estimates are conservative and that the real incidence may

even be tenfold higher. For instance, a series of studies have provided support for the conclusion that up to 15 percent of adult-onset asthma is occupationally related.⁹⁹ NIOSH (National Institute for Occupational Safety and Health) estimates that in the USA, 3–9 million people are exposed to carcinogens in the workplace, and that 2–4 percent of all cancers are occupational in origin.²

Methods of assessing occupational lung disease include use of the chest radiograph and pulmonary function testing. Efforts to try and standardize interpretation of chest X-rays in occupational disease led to the adoption in 1980 of the International Labour Organization (ILO) Classification (see Table 23.14). In the USA, physicians who undergo training and accreditation for the ILO Classification are called 'A' readers if they have attended the course, and 'B' readers if they have taken and passed the very difficult certification test. Using the ILO system requires that readers have

Table 23.14. Classification of occupational lung disorders.

I	Interstitial lung disorders (pneumoconioses)
	A Fibrogenic (e.g. silica, asbestos, silicates)
	B Non-fibrogenic (e.g. tin, iron, barium)
II	Airway disorders
	A Upper respiratory tract irritation (e.g. formaldehyde)
	B Industrial bronchitis (e.g. coal dust)
	C Laryngeal edema (e.g. ammonia)
	D Allergic occupational asthma (e.g. TDI, red cedar)
	E Non-allergic occupational asthma (e.g. RADS or VCD)
	F Bronchiolitis obliterans (e.g. nitrogen dioxide)
	G Byssinosis, etc. (e.g. cotton, grain, flax dust)
III	Alveolar disorders
	A Adult respiratory distress syndrome (e.g. phosgene)
	B Chemical pneumonitis (e.g. cadmium fumes)
	C Granulomatous disorders (e.g. berylliosis, extrinsic allergic alveolitis)
	D Organic dust toxic syndrome (e.g. fungal dust)
	E Fume fever (e.g. metal or polymer fumes)
IV	Pulmonary infectious disorders
	A Indoor (e.g. <i>Legionella</i> , tuberculosis)
	B Outdoor (e.g. histoplasmosis, anthrax)
V	Lung cancer
	A Bronchogenic (e.g. asbestos, bis-chloromethyl ether)
	B Pleural (e.g. mesothelioma)

available to them the standard set of radiographs, which are fundamental to the interpretation.

Lung parenchymal abnormalities are classified into large (>1 cm) and small opacities. Small opacities are further defined by assigning letters for increasing size and shape: *p*, *q*, and *r* for small, rounded opacities, and *s*, *t*, and *u* for small, irregular opacities. There may be a combination of two designations, with the first indicating the most numerous change. A report of *s/t* in a patient with asbestosis, for instance, would indicate that the *s* was the primary opacity and *t* was the secondary small, irregular opacity noted.⁹⁹

Spirometry is the major form of pulmonary function testing and is perfectly adequate for screening and for minor abnormalities. However, if there is question about significant abnormalities, full pulmonary function studies, usually available only in hospital laboratories, are necessary to measure actual lung and diffusion capacity.

'Pneumoconiosis' is derived from the Greek—*pneumo* (lung) and *Konis* (dust)—and describes lung diseases resulting from the inhalation of mineral dusts. Over the course of time it has been adapted to indicate the nature of the dust (e.g. silicosis (silica), asbestosis (asbestos), stannosis (tin)); to indicate the occupational risk (e.g. coal workers' pneumoconiosis); and to include organic dusts (e.g. byssinosis).¹⁰⁰

Asbestos Lung and Pleural Disease

Asbestos is a term applied to mineral silicates with a 'fibrous habit' (i.e. they occur naturally in bundles of parallel, radiating or interlaced fibers that can be readily separated into javelin-like fibers. These fibers have in common a remarkable resistance to deterioration due to heat, cold, salt water or a variety of other influences, making them ideal insulating material. Their fibrous nature also allows them to be spun.

World use of asbestos increased exponentially between the two world wars, but then peaked in the late 1970s and 1980s in the USA and other developed countries when the ill-health effects of exposure became known. Since asbestos is a cheap, durable material with remarkable insulating properties, its production and use have continued to increase in emerging markets.

It has been estimated that between the years 1940 and 1979 roughly 19 million US workers had significant occupational asbestos exposure. Of this number, it has been estimated that several hundred thousand deaths have already occurred, primarily due to lung cancer, and to a lesser extent to mesothelioma and other cancers.^{101,102}

Asbestos fiber inhalation results in three major health consequences. First, the fiber causes pleural disease. The pleura is the lining membrane of the chest wall (parietal pleura) and of the lung (visceral pleura). Presumably the javelin-like asbestos fibers are inhaled lengthwise into the lung parenchyma, then migrate to the pleural surfaces. The resulting reaction causes pleural plaques, discrete areas of pleural thickening that may be evident on routine chest X-rays. Occasionally, benign pleural effusions may also be the presumed consequence of asbestos inhalation. Although there is no precise dose-response relationship between exposure to asbestos fibers and consequent pleural disease, there does appear to be a threshold below which no risk occurs and above which there is an increase in proportion affected with increasing exposure. This threshold is thought to be the OSHA (Occupational Safety and Health Administration) standards for workplace exposure limits, but in reality the actual threshold for 'safe' exposure limits is unknown.

Asbestos-caused pleural plaques and benign pleural effusions, in and of themselves, are usually of no major concern for significant morbidity or mortality. However, workers with pleural plaques are more likely to develop asbestos-induced parenchymal fibrosis than similarly exposed workers without plaques. Also, there is an increased risk of lung cancer among similarly exposed workers who develop pleural disease. Thus from an underwriting standpoint a finding of pleural plaques (or benign pleural effusions) thought to be a consequence of asbestos exposure should serve as a 'red flag' for potential development of fibrotic lung disease or cancer.

A second consequence of asbestos fiber inhalation is the development of fibrosis. This process is termed asbestosis and is most prominent in the lower lobes of the lungs and in the subpleural areas (in contrast to silicosis and coal workers' pneumoconiosis, which predominantly involve

the upper lung zones). Microscopically, the fibrosis seen is indistinguishable from the fibrotic reaction to other fibrogenic dusts and idiopathic pulmonary fibrosis, except for the presence of the asbestos fibers. The actual fibers are invisible on routine light microscopy and require electron microscopy or phase contrast microscopy for identification. However, some of these fibers will be surrounded and 'coated' with alveolar macrophages and other inflammatory cells which will stain for iron, and which are referred to as ferruginous bodies. Most commonly, there are hundreds or even thousands of uncoated fibers for every ferruginous body seen. It is a comment on the ubiquitous use of asbestos that autopsy studies of unexposed persons invariably reveal the presence of asbestos fibers, if a thorough search for them is performed.

The pulmonary fibrosis following asbestos exposure requires more than just the exposure history to confirm causality. There must be a minimum of at least 6 months of a moderate intensity of exposure and at least 20 years from time of exposure to appearance of clinical and roentgenographic disease. In general, the grade of pulmonary fibrosis relates to the fiber burden carried by the lungs. Radiologic asbestosis may remain static or progress, following cessation of exposure. Regression does not occur. Dreesen *et al.*, studying dust and disease in a South Carolina textile factory, concluded that the threshold for development of asbestosis occurred at dust levels above 5 million particles per ft³ (probably equivalent to 15–30 fibers/ml).^{103,104}

Asbestosis is most commonly heralded by shortness of breath (dyspnea). Cough is also commonly reported, and is often non-productive in non-smokers and productive in smokers. On physical examination, the most common finding is auscultatory crackles (rales) over the lung bases. Although the presence of 'small irregular opacities' of fine reticular or ground-glass appearance in the mid-and lower lung fields ('shaggy heart border') remains the *sine qua non* for pulmonary asbestosis, high-resolution computer tomography (HRCT) will reveal interstitial abnormality before it is visible by routine chest radiography.

Asbestosis is classically described as a restrictive lung process, with less-than-predicted lung

volumes, decreased diffusing capacity and impaired gas exchange on pulmonary function testing. However, airway obstruction is also commonly seen, both because cigarette smoking is common in this population and because asbestos fibers apparently cause airway obstruction even without the confounding influence of smoking.

A third consequence of asbestos fiber inhalation is the development of malignancy. Most commonly the malignancy developed is lung cancer, but in fact there is an increased incidence of malignant tumor of any tissue site that may be impacted by the asbestos fiber, including the larynx, esophagus and stomach. There is a striking relationship between asbestos exposure and the development of a primary malignancy of the pleura, called malignant mesothelioma. Some federal courts have even judged the occurrence of this neoplasm to signal significant asbestos exposure, recognized or not. Epidemiologic studies of lung cancer mortality in exposed work forces have confirmed the excess risk of asbestosis and have shown it to be related to the intensity and duration of exposure, thus providing evidence for causality. The increased association of cancer with cigarette smoking and with asbestos exposure has been found to be multiplicative, not just additive. The increased risk of asbestos workers who smoke for developing lung cancer is estimated to be as high as 50–90 times that of non-smoking, non-asbestos-exposed persons.

Although human exposure to asbestos is probably universal, significant exposure sufficient to cause asbestosis or cancer is usually occupational. Recently in the USA there has been concern about the risk of living or working in a building or home originally built with asbestos products (especially in some one-third of US schools). The lifetime risk for premature cancer death from such exposure has been estimated at only 4–6 per million exposed.

Risk Selection

The diagnosis of asbestos lung or pleural disease, either admitted on the application or incidentally found in the attending physician's statement (APS), should be significant and unequivocal. Litigious and compensation-related concerns have caused many workers exposed to asbestos

to be 'diagnosed' as having asbestos disease, when in reality they may only have, at most, asbestos pleural disease or very early or mild fibrotic change. Certainly there remains some risk for development of malignancy in this cohort, especially among the smokers, but unless they have significant symptoms or chest X-ray changes most will certainly remain insurable. Chest X-ray reports and, if available, pulmonary function studies should confirm either the absence of significant disease or the presence of mild dysfunction.

The presence of significant symptoms or chest X-ray signs of asbestos fibrotic disease, on the other hand, suggests significant morbidity and mortality concerns, especially with the observation that progressive disease may occur despite cessation of exposure to asbestos.

Finally, the presence or history of any asbestos-related malignancy is cause for the greatest concern and caution by underwriters. Many of these malignancies are not curable and occur in the setting of significant asbestosis.

Silicosis

Silicosis refers to lung disease caused by inhalation of silica (silicon dioxide) in crystalline form, usually as quartz. Silica is the earth's most abundant mineral. Minerals with high amounts of free silica include quartz (also granite), flint, chert, opal, chalcedony and diatomite. Combined forms of silica (*silicates*) include, among others, asbestos, talc, mica and kaolin. Whenever the earth's crust is broken and silica-containing rock or sand is gathered, milled and processed, or used, there are potential risks for the men and women whose occupation(s) involve them in these processes.¹⁰⁵

Silicosis is a disease defined by the chest X-ray. Simple silicosis appears as small (less than 10 mm in diameter) rounded opacities, usually in the upper lung fields, and often accompanied by enlarged and calcified hilar lymph nodes. Progressive massive fibrosis (PMF) is the result of coalescence of these small nodules into larger opacities, also typically in the upper lung fields. These disease processes require a lifetime of occupational exposure to develop. Rarely, in cases of massive exposure to dust with very high quartz content, an acute form of silicosis, silicoproteino-

sis, may be seen, with disease evolving over a time period of months to years.

There are three requirements for the diagnosis of silicosis: (1) a history of silica exposure; (2) the presence on chest X-ray of characteristic small rounded opacities in upper lung fields; and (3) the absence of other illnesses that might mimic silicosis, such as miliary tuberculosis or fungal disease. Usually exposure for 10–30 years is required before these chest radiographic changes are noted. Although host factors, such as genetics, smoking and underlying diseases, may play a part in the development of silicosis, the most important factors include silica dust in ambient air, duration of dust exposure, the polymorphic structure of the silica particles, the percentage of free silica and the particle size.¹⁰⁶

Silicosis is incorrectly considered by many physicians to be a radiologic abnormality dissociated from impairment, disability or excessive mortality. In fact, survival may be reduced in persons with dyspnea, sputum production or abnormal breath sounds on physical exam, and a reduced vital capacity (reduced FVC % predicted) on spirometric testing. Spirometry more commonly indicates obstruction or combined obstruction and restriction, rather than just pure restriction, even in non-smokers. Up to one-half of cases of simple or conglomerate silicosis demonstrate progression on chest X-ray even after exposure has ceased. There is a questionable slight increased incidence of lung cancer in persons with more advanced silicosis. This association remains controversial, however, due to the difficulty in controlling for other carcinogenic exposures common to the population of workers exposed to silica, such as radon, asbestos and smoking. Also, it may be that the respiratory impairment in the more severe cases of silicosis results in premature deaths and masks the role of silica as a carcinogen in some of the negative studies.

The chest radiograph in silicosis is rather distinctive in that the infiltration of small rounded densities are usually confined to the upper lung zones, and the hilar lymph nodes calcify in a peculiar peripheral fashion, often referred to as 'egg-shell calcification'. Conglomeration of the opacities into PMF, also usually confined to the upper lung zones, results in retraction of the hilar

pulmonary vasculature into the upper lungs. Pleural plaques occur but are not a common feature of silicosis.

An interesting association exists between silicosis and tuberculosis. Rates for active tuberculosis in silicotic workers range from 5 percent to 43 percent, depending on whether radiologic or autopsy criteria are used, an increase of 2- to 30-fold more than in non-silicotic miners. The presence of cough, hemoptysis, and the systemic symptoms of weight loss and fever in a person with silicosis should always suggest the possibility of mycobacterial disease. The incidence of tuberculosis increases from simple silicosis to PMF, with the highest incidence occurring in acute silicosis (alveoloproteinosis) and commonly accounting for death in this setting.

Although silicosis is associated with increased prevalence of positive autoimmune serologies, antinuclear antibodies and elevated immunoglobulin levels have not correlated with the baseline profusion category of the chest radiograph, the rate of chest radiographic progression or the rate of lung function decline in sandblasters with silicosis. Rheumatoid nodules may appear in the upper lung zones of silica workers with rheumatoid arthritis, a coincidence termed Caplan's syndrome.

Risk Selection

Once silicosis develops, the primary risks to the applicant are progression of disease and the development of mycobacterial infection. Simple silicosis is associated with little mortality significance. Conglomerate silicosis (PMF), especially when accompanied by significant symptoms and pulmonary function abnormalities, calls for much greater concern in underwriting. Acute silicoproteinosis is such a severe disease that it would be unlikely for anyone with this disorder to be an applicant for insurance.

Coal Workers' Pneumoconiosis (CWP)

Coal is the end result of a gradually formed sedimentary process beginning with decaying vegetable matter and evolving through peat, lignites, sub-bituminous and bituminous coals to anthracite through the application of heat and great

pressure over time. The carbon content increases progressively from peat through various stages to anthracite, and as the percentage of carbon increases the amount of quartz and other minerals decreases. The degree of risk to which an applicant for insurance has been exposed therefore depends on the type of coal mined as well as the type of mining done (surface strip mining versus underground mining).

CWP is a distinct pathological entity resulting from the inhalation of coal dust into the lungs. As coal dust also contains a variable amount of silica, some degree of silicosis may also be found in many of these persons. The deposition of the dust in the distal airways and lung parenchyma results in two lesions. Deposition of dust in the distal airways and lung appears to cause bronchitis and emphysema similar to that seen with cigarette smoking, with expiratory airflow obstruction and hyperinflation on pulmonary function testing. The dark pigment localizes in lung macrophages that coalesce at the level of respiratory bronchioles, causing scarring and destruction of adjacent alveolar walls and forming the characteristic dust macule that is considered pathognomonic of CWP.

With increasing dust deposition, the pigmented macule progressively enlarges, becoming solid and palpable. Coalescence of multiple dust macules results in the formation of nodular lesions, which may cause vascular damage to surrounding lung tissue.¹⁰⁷ Coalescence of many nodular lesions results in PMF and is referred to as complicated CWP. These lesions are greater than 1-2 cm in diameter and are usually seen in the upper portions of the lungs.

Almost all miners with complicated CWP also have severe bronchitis and emphysema. Coal dust appears to be far less fibrogenic than other bioactive dusts such as silica and asbestos.¹⁰⁸ Thus the total dust burden in the lung may be much higher in CWP than in other pneumoconiosis, yet with less consequence. Classic silicotic lesions are seen when lung dust residue contains 18 percent or more quartz.

There were still 175 000 miners working in the USA in 1986.¹⁰⁹ Although the number of coal miners is decreasing worldwide, the increased mechanization of mining is resulting in generation

of finer dusts. This fact, along with the observation that there is a long latency period from exposure to the dust to evolution of lung pathology, means that insurance underwriters will continue to see applicants for insurance for several decades to come.

Overall, the life expectancy of coal miners appears to be comparable to that of the general population.¹¹⁰ Survival rates for men with PMF are substantially reduced, and this disorder is also associated with increased morbidity, impairment of pulmonary function and disability.¹¹¹ Although current dust-control measures in mines have resulted in lower prevalence of CWP, the lifetime risk is still 1.4–14 percent.¹¹² Miners exposed to the current US standard for respirable coal mine dust of 2 mg/m³ or less over a working lifetime are at increased risk of death from pneumoconiosis or from chronic bronchitis or emphysema.¹¹³

Risk Selection

Miners and ex-miners with simple pneumoconiosis survive as well as those with no evidence of pneumoconiosis at all. PMF, the prevalence of which is now low, affects mortality in only a minority of subjects, most of whom have chronic air-flow obstruction related to smoking. The dust effects in miners who smoke cigarettes appear to be additive; no disproportionate effect of smoking has been identifiable.

IMMUNOLOGIC LUNG DISEASE

Hypersensitivity Pneumonitis

Organic material existing as fine particulate matter or aerosols may be inhaled and, if antigenic to the person inhaling this material, may cause acute and chronic reactions in the lungs. These organic agents include vegetable dusts (especially fungal spores), proteins of animal and avian origin, various micro-organisms (bacteria, *Rickettsia* and *Chlamydia*) and certain organic chemicals. The acute reaction consists of constitutional symptoms, the generation of specific precipitating antibodies (precipitins), and lymphocytic infiltration and sarcoid-type granulomas in the walls of alveoli and small airways. The chronic reaction is char-

acterized by an irreversible and often progressive diffuse intrapulmonary fibrosis.

One important characteristic of these various allergens is that the particles are respirable (i.e. they are small enough to be inhaled and deposited in the gas-exchange portion of the lung parenchyma). The disease is usually associated with an occupation or hobby, because a high level of exposure to the allergen is required to induce sensitization. The classic form of this illness is farmer's lung, caused by the inhalation of both thermophilic bacteria and several fungal species.

Risk Selection

A history of a limited number of episodes of acute hypersensitivity pneumonitis would be expected to be associated with little excess mortality if current pulmonary function testing is normal and if subsequent exposure to offending antigens is avoided. However, continuing exposure may result in chronic low-grade pulmonary inflammation and progressive pulmonary disease in the absence of acute episodes. In these cases, the mortality risk of hypersensitivity pneumonitis is related to the degree of pulmonary impairment present and to the degree to which antigen exposure is likely to continue.

A study from Mexico City of 74 female and 4 male patients (mean age 41 years) with chronic pigeon breeder's lung revealed a survival rate of 71 percent 5 years from the time of diagnosis. The median survival from the beginning of symptoms was 134 months. At the time of diagnosis, mean predicted FVC was 45 percent and the mean predicted TLC was 70 percent. Using a Cox proportionate hazards model and after controlling for the effects of age, sex, clubbing, FVC and TLC, the investigators found that the extent of fibrosis and the presence of honeycombing on chest radiographs remained significantly and inversely related to survival with relative mortality risks of 220 percent and 290 percent respectively.¹¹⁴

Goodpasture's Syndrome

Goodpasture's syndrome is an autoimmune disease characterized by diffuse alveolar hemorrhage and glomerulonephritis. In this disorder, circulating IgG antibodies become bound in a linear

fashion to glomerular and alveolar capillary basement membranes. Accordingly, this disease is sometimes referred to as anti-basement membrane antibody disease. Circulating anti-basement membrane antibody can be detected in over 90 percent of cases.

Histopathologic examination of lung tissue reveals diffuse alveolar hemorrhage, hemosiderin laden macrophages, mild to moderate interstitial fibrosis and proliferation of type II pneumocytes. Renal lesions consist of a focal segmental necrotizing glomerulitis with crescent formation. Most often pulmonary and renal disease occur simultaneously; however, in 25–33 percent of cases, renal disease occurs alone.

Young adult males are most commonly affected, and alveolar hemorrhage occurs with increased frequency in smokers. Hemoptysis, cough and dyspnea usually precede the manifestations of renal disease. Fatigue is also common, and some degree of iron deficiency anemia is usual. Chest radiographs reveal patchy or diffuse alveolar infiltrates corresponding to alveolar hemorrhage. With resolution of the hemorrhage, reticular-nodular infiltrates may develop. A restrictive pattern is seen on pulmonary function testing. Due to the presence of blood in the air-spaces, carbon monoxide diffusion capacity is increased during episodes of acute alveolar hemorrhage.

Treatment of Goodpasture's syndrome consists of corticosteroids, cytotoxic agents and plasmapheresis. Renal transplantation is recommended for anuric patients. Despite advances in treatment, the prognosis remains poor. Overall, the 2-year survival rate is 50 percent. Most patients die from alveolar hemorrhage. Severe renal failure is associated with a 6-month survival rate of 50 percent.¹¹⁵

Wegener's Granulomatosis

Wegener's granulomatosis is a systemic necrotizing vasculitis that affects the upper airways, lungs, glomeruli, eyes, heart, joints, and peripheral and central nervous systems. Sinusitis is the most common initial manifestation. Some 95 percent of patients have upper airway disease, over 90 percent have lung disease and 85 percent have

glomerulonephritis. The term limited Wegener's granulomatosis refers to those cases of Wegener's granulomatosis in which glomerulonephritis is absent. Limited Wegener's granulomatosis is felt to have a better prognosis than classical Wegener's. Pericarditis, myocarditis or cardiomyopathy occur in approximately 10 percent of patients. Other manifestations include episcleritis, uveitis, retro-orbital masses, arthralgias, arthritis, mononeuritis multiplex, cranial nerve palsies and subcutaneous nodules.

Pulmonary manifestations of Wegener's include pulmonary infiltrates and multiple bilateral or solitary nodules which may or may not cavitate. Atelectasis, pleural effusions and hemoptysis may also occur.

Although the diagnosis of Wegener's granulomatosis is usually made on the basis of biopsy evidence of a necrotizing granulomatous vasculitis, the presence of serum antineutrophil cytoplasmic antibodies (ANCA) has been shown to be a highly specific marker for Wegener's granulomatosis. Two patterns of neutrophil staining by these antibodies have been reported: diffuse cytoplasmic (C-ANCA) and perinuclear (P-ANCA). P-ANCA staining is seen in other vasculitides including Churg-Strauss vasculitis, idiopathic glomerulonephritis, polyarteritis nodosum and systemic lupus erythematosus (SLE). In contrast, C-ANCA is highly specific for Wegener's granulomatosis.

In the past, Wegener's granulomatosis was fatal within a few months, death usually resulting from renal failure. However, treatment with cytotoxic agents, such as cyclophosphamide alone or in combination with corticosteroids, has significantly improved survival. Yet relative mortality ratios remain high. In a study of 77 patients with Wegener's granulomatosis enrolled in the American College of Rheumatology vasculitis classification study and followed for a mean of 7.1 years, the standardized mortality ratios were 680 percent for females, 400 percent for males and 470 percent for males and females combined.¹¹⁶

Pulmonary Eosinophilia

This is a heterogeneous group of disorders characterized by abnormal accumulation of eosinophils either in lung tissue or in the blood stream, or both.

Transient Pulmonary Infiltrates with Eosinophilia (Loffler's Syndrome)

Persons with this disorder are usually asymptomatic. The disorder is often discovered in the course of evaluating an abnormal chest radiograph obtained as part of a routine examination. Common symptoms include transient cough, mild fever and malaise. Pulmonary infiltrates are seen on chest radiographs. These infiltrates are typically bilateral, peripheral in location and transient, usually resolving spontaneously in 2–4 weeks.

Loffler's syndrome has been shown to result from exposure to a variety of agents, including parasitic infections and exposure to drugs and chemicals.

Ascaris infections (*Ascaris suum* or *Ascaris lumbricoides*) are classically associated with Loffler's syndrome. Pulmonary infiltrates are seen 10–16 days following ingestion of *Ascaris* larvae when the larvae have migrated through the intestinal wall into the mesenteric lymphatics and venules, eventually to be filtered out in the pulmonary microcirculation. The larvae migrate into the alveoli and up the bronchial tree to be swallowed and to re-enter the intestinal tree for maturation. Parasites and ova will not appear in the stool until 6–12 weeks later, so the diagnosis is often missed. Corticosteroids offer dramatic relief, and if the ingested larvae are *Ascaris*, a single dose of piperazine (4 g) is curative. Other parasites (*Entamoeba histolytica*, *Strongyloides stercoralis* etc.) may also cause this disorder.

A similar illness may occur as a consequence of inhaling organic and inorganic chemicals or ingesting medicines that elicit an inflammatory or hypersensitivity reaction. A large number of medicines have been implicated in Loffler's syndrome. Usually such reactions are self-limited and resolve with discontinuance of the medication, but not always. The best-described instance of this reaction causing permanent lung damage is with the drug nitrofurantoin, a medicine commonly used as a suppressant against chronic urinary tract infections.

Tropical Pulmonary Eosinophilia

Tropical Pulmonary Eosinophilia is a disorder related to Loffler's syndrome but is more serious. Tropical pulmonary eosinophilia is caused by

mosquito-borne parasites that produce an eosinophilic alveolitis. The classic causes of tropical pulmonary eosinophilia are due to the filarial nematodes *Wuchereria bancrofti* and *Brugia malayi*. The blood eosinophilia is much more severe than in simple Loffler's, and patients are usually much more symptomatic. Prompt treatment (Diethylcarbamazine at 6 mg/kg per day for 3 weeks) is usually curative, but chronic disease or delayed treatment may lead to significant pulmonary fibrosis.

Acute Eosinophilic Pneumonia

This is a newly described entity, usually occurring in young people and presenting with acute dyspnea and ground-glass or reticulonodular infiltrates on chest X-ray. Half of affected patients have pleural effusions. The diagnosis is made by bronchoalveolar lavage or transbronchial lung biopsy. There is rapid clinical improvement and roentgenographic clearing within days of instituting corticosteroid therapy. No long-term sequelae have been reported.^{117–119}

Chronic Eosinophilic Pneumonia (Carrington's Eosinophilic Pneumonitis)

This disorder occurs most commonly in women in the fourth or fifth decades of life but may occur in either sex at any age. It is characterized by the coexistence of blood eosinophilia, elevated sedimentation rates, and pulmonary infiltrates due to the accumulation of eosinophils in alveolar and interstitial spaces. Asthmatic symptoms may also occur. The chest X-ray presentation is distinctive and consists of peripheral infiltrates in the upper lobes, the so-called 'photographic negative of pulmonary edema'. The cause is unknown, and the disease is usually curable with corticosteroids, although prolonged therapy may be required.

Allergic Bronchopulmonary Mycoses

Inhalation of fungal spores in patients with asthma may result in endobronchial colonization of the respiratory tree. The clinical picture is that of asthma with recurring lung infiltrates or areas of atelectasis due to mucous-plugging. The most common of these syndromes is allergic bronchopulmonary aspergillosis (ABPA) caused by inhalation and colonization with *Aspergillus fumigatus*.

Therapy consists of oral and inhaled corticosteroids. This syndrome may cause some increased morbidity and should therefore be of concern to health insurance underwriters, but it would be a rare cause of mortality by itself.

Hypereosinophilic Syndrome

This is a rare disorder in which there is massive infiltration of eosinophils in the lung, heart, CNS, skin and other organs. Males are predominantly affected, and symptoms are often severe. The heart is involved in more than half of all cases, and severe dysrhythmias and valvular incompetencies are a usual cause of death. Pulmonary embolism is also a common cause of death. Although an initial response may be seen with corticosteroid treatment, relapse and death ultimately occur.

Allergic Granulomatosis with Vasculitis (Churg-Strauss Syndrome)

This is a vasculitic syndrome which often results in granuloma formation and necrosis in lung tissue. Persons with this disorder have asthma, marked eosinophilia and systemic symptoms. The diagnosis is established by lung biopsy demonstrating an eosinophilic vasculitis affecting small and medium-sized arteries and veins. High-dose prednisone therapy (and, in resistant cases, azathioprine) may achieve 5-year survival rates of 60 percent, with 50 percent surviving 9 years.¹²⁰

Pulmonary Eosinophilic Granuloma

This disorder is part of the clinical spectrum of histiocytosis X, a group of disorders characterized by proliferation of Langerhans' histiocytes. Peripheral eosinophilia is not part of the clinical picture of eosinophilic granuloma, but the infiltrates on chest X-ray are due to eosinophilic accumulations (For a further discussion of pulmonary eosinophilic granuloma, see 'Interstitial and Infiltrative Lung Disease' above.)

PULMONARY HYPERTENSION

The pulmonary arterial tree is a low-pressure system. Normal mean pulmonary arterial pressure is only 14 torr, about 1/6 that of the systemic arterial pressure. Pulmonary arterial pressure normally

rises very little with exercise, due to the capacity of the pulmonary vasculature to recruit vessels normally not perfused at rest. When pulmonary systolic pressure rises above 30 torr, or mean pulmonary pressure rises above 18 torr, pulmonary arterial hypertension is said to be present. Pulmonary venous hypertension is present when mean pulmonary venous pressure rises above 12 torr.¹²¹ As pulmonary pressure rises, the right ventricular workload increases and right ventricular hypertrophy may occur. Elevation of right ventricular end diastolic pressure is indicative of right ventricular failure. The term *cor pulmonale* has been defined variously as meaning right ventricular hypertrophy (the preferred definition of the World Health Organization) or right ventricular failure.¹²²

The etiologies of pulmonary hypertension have classically been described in terms of precapillary and postcapillary causes (see Table 23.15).

Table 23.15. Causes of pulmonary hypertension.

Precapillary causes	Postcapillary causes
Vascular	Cardiac
Left-to-right shunt	Left ventricular failure
Primary pulmonary hypertension	Mitral stenosis
Pulmonary embolic disease	Left atrial myxoma
Pulmonary vasculitis	Cor triatriatum
Low ambient oxygen tension (e.g. high altitude)	
Pleuro-pulmonary disease	Pulmonary-venous
Emphysema	Congenital stenosis
Interstitial disease	Pulmonary veno-occlusive disease
Fibrothorax	Anomalous pulmonary venous return
Kyphoscoliosis	
Alveolar hypoventilation	

In most instances the presence of pulmonary hypertension is associated with increased mortality.

Primary pulmonary hypertension (PPH) is a disease of unknown etiology. Diagnostic criteria require a mean pulmonary artery pressure above 25 torr at rest or above 30 torr during exercise. PPH is for the most part a diagnosis of exclusion: all other known causes of pulmonary hypertension

must be ruled out. However, it should also be noted that characteristic histopathologic changes have been described in muscular pulmonary arteries. These changes include medial hypertrophy, intimal proliferation and fibrosis, vascular dilatation and occlusion, plexiform lesions and necrotizing arteritis.

Although PPH has traditionally been regarded as a disease of young women, it affects both men and women of all ages. The prevalence of PPH has been estimated to be 1.3 per 1000 in an autopsy series.¹²³ Some cases of PPH may have been related to ingestion of toxins or medications. Aminorex, an appetite suppressant, was suspected of causing cases of PPH in Europe in the late 1960s. More recently another appetite suppressant, fenfluramine, has been shown to increase the risk of PPH.¹²⁴

The prognosis of PPH is poor. In 1991, survival data from the Patient Registry for the Characterization of Primary Pulmonary Hypertension were published.¹²⁵ The estimated median survival was 39 months for men and 32 months for women.

SLEEP APNEA

The sleep apnea syndromes are a group of disorders characterized by the presence of periodic pauses in breathing during sleep. These pauses may either be complete, during which airflow totally ceases (apneas), or partial, during which airflow is significantly diminished but does not entirely cease (hypopneas). Sleep apnea may be the result of a failure of the brainstem to continue to stimulate breathing during sleep (central sleep apnea or Ondine's curse) or it may be due to airflow obstruction at the level of the pharynx or hypopharynx (obstructive sleep apnea). The combination of central apneas followed by obstructive apneas is termed mixed sleep apnea. By far the most common form of sleep apnea is the obstructive sleep apnea syndrome (OSAS) which consists of pure obstructive apneas and/or mixed apneas.

The consequences of sleep apnea include: repetitive drops in blood oxygen levels during sleep (oxygen desaturations); complete or partial arousals from sleep (sleep fragmentation); excessive daytime sleepiness (EDS or hypersomno-

lence); arrhythmias (sinus pauses, transient heart block, and atrial and ventricular arrhythmias during apneas); hypertension; impaired neuropsychological functioning (including altered libido, impotence, irritability, personality disturbances, inability to concentrate, fatigue and an increased risk of industrial and driving accidents); and sudden death.

An uncommon but particularly severe form of sleep apnea is the Pickwickian syndrome, which is also called the obesity hypoventilation syndrome (OHS). This syndrome consists of obesity, sleep apnea, daytime hypersomnolence, hypercarbia and hypoxemia during wakefulness, and evidence of pulmonary hypertension or right heart failure.

Diagnosis

Obstructive sleep apnea is usually suspected when a history of loud snoring is present in addition to one or more of the following features: (1) witnessed apneic episodes, (2) hypersomnolence, (3) change in personality or ability to concentrate, (4) insomnia, and (5) significant obesity (however, many persons may have sleep apnea without being significantly overweight).

Even though sleep apnea may be suspected from clinical symptoms, the diagnosis requires confirmation by nocturnal polysomnography. Typically a full polysomnographic study consists of monitoring and recording EEG activity, extraocular eye movements, submental EMG activity, nasal and oral airflow, respiratory effort (by impedance plethysmography, respiratory strain gauge or esophageal balloon), ear oximetry, ECG and lower extremity EMG activity. An accurate diagnosis of sleep apnea requires at least 6 hours of monitoring and includes recording of sleep stage, airflow, respiratory effort, blood oxygen level (oxygen saturation) and ECG.

Sleep apnea severity is usually expressed in terms of an apnea index (AI) or a respiratory disturbance index (RDI). The AI is defined as the average number of apneas occurring per hour of sleep. The RDI, also known as an apnea-hypopnea index, is defined as the average number of apneas and hypopneas occurring per hour of sleep. Normal values for AI and RDI increase with age. For young adults, an AI less than 5 or a RDI

less than 10 is considered by most authorities to be normal.

Consequences

Apneas and hypopneas result in episodic oxygen desaturations. The severity of desaturations correlates with the length and frequency of apneas and the presence of underlying pulmonary disease. Desaturations are commonly defined as drops in oxygen saturation 4 percent or more below the baseline oxygen saturation level. In cases of severe sleep apnea, these desaturations may occur frequently and be prolonged and severe. Desaturations to 40 percent or lower, occurring at a frequency of 40 or more per hour, and lasting 60 seconds or more, are not uncommonly seen in sleep laboratories.

The consequences of recurrent hypoxemia include bradycardia, other cardiac dysrhythmias, pulmonary hypertension, congestive heart failure, sleep arousal and fragmentation, and hypoxic cerebral depression.¹²⁶ However, little is known about the severity or frequency of the desaturations that are necessary to produce these adverse consequences. There is little evidence to suggest that oxygen desaturations occurring during sleep in the absence of waking hypoxemia contribute to pulmonary hypertension or cor pulmonale,¹²⁷⁻¹²⁹ and whilst nocturnal oxygen desaturations may eventually be shown to be the cause of the excess mortality associated with sleep apnea syndrome, there is presently no definitive evidence to confirm this hypothesis.

There is some evidence, however, that nocturnal oxygen desaturations are at least in part responsible for the diminished daytime vigilance seen with sleep apnea syndrome. Oxygen saturations of 60 percent or lower during the night have been shown to correlate with a mean time to sleep onset of less than 5 minutes, which is indicative of pathological sleepiness. Lowest nocturnal oxygen saturations of 60-95 percent have been shown to be associated with mean sleep latencies in the intermediate range of 5-10 minutes.

Arrhythmias are common in OSAS. Over 75 percent of patients with OSAS have sinus bradycardia and 10 percent of patients may have sinus pauses up to 13 seconds. Mobitz Type 2 AV block

may occur in 4-8 percent of patients. Ventricular ectopy occurs in 57-74 percent of OSAS patients. However, this frequency is not different from what would be expected in a cross section of middle-aged men. When oxygen saturations drop to less than 60 percent, however, ventricular ectopy increases threefold. Above a saturation of 60 percent there is no increase in ventricular ectopy.^{130,131}

Recently studies have implied that nocturnal hypertension may be the mechanism by which sleep apnea causes excess mortality. Normally blood pressure dips at night. However, microarousals during sleep may result in cardioacceleration and nocturnal hypertension. In some patients, nocturnal microarousals may also contribute to waking hypertension.¹³²

In a study from Japan, Noda *et al.* studied 51 patients with sleep apnea, 34 of whom had waking hypertension. Polysomnograms, echocardiograms and 24-hour blood pressure monitoring were performed. Patients with an apnea index (AI) of more than 20 had a higher incidence of left ventricular hypertrophy (LVH) than those with an AI of less than 20. They also found that levels of total plasma norepinephrine after waking were significantly increased compared with levels before sleep and that this increase was correlated with the duration of oxygen saturation below 90 percent but not with the AI. The authors postulated that sleep apnea may cause nocturnal hypertension through the stimulation of the adrenergic nervous system and thereby produce LVH.¹³³

Neuropsychiatric complications of sleep apnea are common. Excessive daytime sleepiness (EDS) is probably the most common presenting symptom in patients with obstructive sleep apnea. Cognitive impairment is another well recognized complication of obstructive sleep apnea. Sleep fragmentation and nocturnal hypoxemia have been implicated as causes for both. As a consequence of EDS and cognitive impairments, patients with obstructive sleep apnea have lower job performance scores and higher accident rates.¹³⁴ Findley *et al.* found that patients with obstructive sleep apnea had motor vehicle accident rates 7 times greater than controls and 2.6 times greater than all licensed drivers in Virginia.¹³⁵ Stoohs reported that long-haul commer-

cial truck drivers with sleep-disordered breathing had a twofold higher accident rate per mile than drivers without sleep-disordered breathing.¹³⁶

Mortality Risk

Only a few studies have been reported that attempt directly to assess the mortality risk associated with obstructive sleep apnea. One of the earliest studies was reported by Gonzales-Rothi *et al.*¹³⁷ They compared the course of 91 patients having polysomnographically documented sleep apnea between July 1978 and June 1986 with a control group of 35 patients who had symptoms but no polysomnographic evidence of sleep apnea. Nine of the sleep apnea patients and four of the control patients died by the end of the study, a statistically insignificant difference. This study has been criticized on the basis of its small size, the difference in age between the two study groups and an excessively high death rate among controls.

Partinen *et al.* reported on 190 patients with obstructive sleep apnea who were either advised to lose weight (127 patients) or treated with tracheostomy (71).¹³⁸ Despite the fact that patients treated with tracheostomy had more severe sleep apnea (AI = 69 vs 43) all of the 14 deaths that occurred during the 5-year study occurred in the group that was only advised to lose weight.

In 1988 He *et al.* reported the results of a study of 706 patients with obstructive sleep apnea who were followed for up to 9 years.¹³⁹ Their data indicate that mortality is increased for men who had an AI greater than 20. The relative mortality risk for males in this study over the age of 50 having an AI greater than 20 is estimated at 168 percent compared with mortality expectations derived from 1979–1981 US life tables. Similarly, for males under age 50 who have an AI greater than 20 the relative mortality risk is estimated to be 425 percent. The authors of this study also reported that treatment with nasal continuous positive airway pressure (NCPAP) restored mortality to normal whereas surgery designed to remove redundant tissue from the posterior pharynx (uvulopalatopharyngoplasty or UPPP) did not.

More recently data have been reported suggesting that some of the risk attributed to sleep apnea

may be due to the presence of comorbid factors such as hypertension, heart disease and lung disease. Lavie *et al.* reported a series of 1456 men and 164 women with sleep apnea between the years 1976–1988.¹⁴⁰ They found that sleep apnea syndrome, defined as an AI greater than 10, was associated with significant excess mortality in men between the ages of 30 and 50. The relative mortality ratios for ages 20–30 and 50–70 were also greater than unity, but not significantly so. The authors then used a Cox proportionate hazard model to determine independent risk factors for subsequent mortality. They found that although the AI was identified as an independent predictor of death from all causes, it was not a predictor of death from cardiopulmonary causes or of death due to MI. Even in the analysis of deaths from all causes, the AI made the least contribution to the model.

In agreement with the study reported by Lavie, other studies have also indicated that sleep apnea may be less of a risk factor for mortality in the elderly than in younger persons. In 1988 Bliwise *et al.* reported that in their study of 198 persons having a mean age of 66.6 years, an RDI of 10 or greater was associated with a 2.67 relative risk of mortality, but the 95 percent confidence intervals included unity.¹⁴¹

In 1990 Pollack *et al.* studied sleep problems in 1855 community-living elderly. For females the presence of symptoms of sleep apnea was not a predictor of mortality. Although the relative mortality risk for males who reported a history of sleep apnea was 131 percent, this increase was not statistically significant.¹⁴²

More recently Mant *et al.* followed a cohort of 163 non-demented community-dwelling individuals (mean age = 80.6) for 4 years. The presence of sleep apnea (defined as a RDI \geq 15) was not found to be predictive of mortality.¹⁴³

Treatment

Central apnea may be treated with diaphragm pacemakers, nocturnal ventilation or medication. Obstructive apnea may be treated by varying the position of the patient during sleep (position therapy), weight loss, medication (protriptyline, medroxyprogesterone, theophylline, almitrine,

oxygen), surgery (UPPP, nasal, maxillary or mandibular surgery, or tracheostomy) by various appliances (tongue retaining devices), or by nasal continuous positive airway pressure (NCPAP) or continuous positive airway pressure (CPAP). All these therapies require repeat polysomnography in order to determine whether they have effectively eliminated apneas. Only tracheostomy and NCPAP have been demonstrated in clinical studies to significantly reduce mortality. NCPAP therapy is often perceived as disagreeable by the patient and a significant percentage of patients prescribed NCPAP will eventually discontinue its use. NCPAP compliance is about 75 percent at 6 months and 66 percent at 1 year; ultimately 50 percent will discontinue the use of NCPAP at 2 years or beyond.^{144, 145}

Risk Selection

In summary, the few studies on the mortality risk of sleep apnea provide evidence that sleep apnea is associated with excess mortality, especially in those under 50 years of age at the time of diagnosis. Studies indicate that although NCPAP is an effective treatment for sleep apnea and restores mortality to normal, there is relatively poor long-term compliance with this form of treatment. Other studies indicate that although upper airway surgery such as UPPP can improve obstructive apneas, significant improvement occurs in only about half of those so treated and that follow-up nocturnal polysomnographic studies are indicated to determine if significant improvement has occurred and whether additional treatment is warranted. There is also evidence to suggest that a portion of the risk attributed to sleep apnea may be due to the presence of co-morbid factors, such as hypertension, heart disease and lung disease, that may have already been considered in the underwriting process.

PLEURAL DISEASE

Although many disease processes affect the pleural linings of the lungs, the pleura respond in only a few ways: pain (pleurodynia), effusion, the development of a pneumothorax and fibrosis. The importance of these manifestations is determined

by the underlying etiology and the degree to which pulmonary function is impaired.

Pleuritis and pleurisy are non-specific terms referring to pain produced by pleural inflammation. Pleuritis can have many causes. Pleural inflammation without effusion has been termed 'dry pleurisy'. Pleural pain usually diminishes as an effusion develops ('wet pleurisy'). One of the most common causes of dry pleurisy is infection with Coxsackie virus. Pleural inflammation caused by Coxsackie virus is so severe, spasmodic and paroxysmal in nature that this disorder has been called 'the devils gripe'. The infection is self-limited and only symptomatic treatment is required. It has no significant mortality implications.

Pleural effusions may result from many causes. The mortality implications of a pleural effusion are determined primarily by the underlying disorder. Pleural effusions have been characterized as transudates or exudates depending on their composition. An exudative pleural effusion associated with a bacterial pneumonia, bronchiectasis or lung abscess is termed a parapneumonic effusion. Parapneumonic effusions are described as 'complicated' if a tube thoracostomy is required for them to resolve. An empyema is defined as pus within the pleural space and almost always requires drainage.

Pneumothoraces may result from many causes. Spontaneous pneumothoraces are those that occur in the absence of obvious trauma or underlying lung disease. Secondary pneumothoraces are those that occur as a result of trauma (traumatic pneumothorax, iatrogenic pneumothorax) or underlying lung disease (secondary spontaneous pneumothorax).

The mortality implication of a pneumothorax depends on the underlying cause and the extent to which pulmonary function is compromised. Primary spontaneous pneumothorax is fairly common, occurring at an incidence of 7.4 per 100 000 per year in men and 1.4 per 100 000 per year in women. Most are believed to be caused by the rupture of apical subpleural blebs. Smokers are at seven times the risk of non-smokers for primary spontaneous pneumothorax. Many primary spontaneous pneumothoraces resolve spontaneously or are easily treated by aspiration or

tube thoracostomy. Persons who have already had one primary spontaneous pneumothorax are at increased risk of having recurrences. Pleural sclerosis or open parietal pleurectomy has been recommended for persons having more than two primary spontaneous pneumothoraces and for those having an episode of bilateral pneumothorax. With few exceptions, one or more episodes of primary spontaneous pneumothorax that have been adequately treated would be expected to result in little if any excess mortality.

Secondary spontaneous pneumothoraces, by definition, occur in the presence of underlying lung disease, and accordingly the mortality risk is increased. COPD is the most common underlying disease, and the risk of pneumothorax increases as the severity of COPD increases. The many other causes include pulmonary tuberculosis, pulmonary fibrosis, sarcoidosis, pneumoconiosis, lung abscess, bronchogenic carcinoma and asthma.

Risk Selection

The Medical Impairment Study 1983 recorded 328 deaths out of 10 180 policies issued from 1952 to 1976 to persons having histories of pleurisy, pleural effusion or empyema within the preceding 3 years. The overall relative mortality ratio was 118 percent compared with standard insured lives mortality. Males over age 50 at the time of policy issue had a relative mortality ratio of 148 percent.¹⁴⁶ The study also recorded 291 deaths out of 16 804 policies issued to persons having histories of spontaneous pneumothorax from 1952 to 1976. Only subjects who were thought to be free of other diseases were included in the study. The overall relative mortality ratio was 140 percent compared with standard insured lives mortality. There did not appear to be a significant difference in mortality by issue age or by duration.¹⁴⁷

INFECTIOUS RESPIRATORY DISEASE

Infectious respiratory diseases comprise a large number and variety of disorders. Infectious bronchitis, bronchiolitis, pneumonia, lung abscess, parapneumonic effusions and empyemas fall into

this category. In general the short-term prognosis of infectious lung disease depends on the immune status of the individual, the infecting organism, the nature and severity of the infection and the adequacy of treatment. The long-term prognosis depends primarily on the immune status of the individual, the degree of residual pulmonary impairment and the presence of coexisting disease.

Acute Pulmonary Infections

Many acute respiratory infections are associated with substantial mortality risk during the stage of active infection. Overall, community-acquired pneumonia is associated with a case-fatality rate of 6–24 percent, and hospital-acquired pneumonia has a case-fatality rate of 2–53 percent.¹⁴⁸ The mortality due to anaerobic lung abscess has been reported to be as high as 15 percent,¹⁴⁹ and the mortality due to empyema varies: 8–15 percent in healthy young persons and 40–70 percent in elderly patients and those with serious underlying disease.¹⁵⁰

Pulmonary Tuberculosis

The worldwide incidence of tuberculosis diminished steadily from the 1960s and reached a nadir in the mid-1980s. Since then, the incidence has again been increasing. It is estimated that currently one-third of the world's population and one-fifth of the US population are infected with *Mycobacterium tuberculosis* as indicated by a positive tuberculin skin test. The World Health Organization has estimated that by the year 2000 the worldwide annual incidence of tuberculosis will rise to over 10 million new cases, an increase of over 35 percent compared with the 1990 incidence. Of the 3.5 million deaths from tuberculosis that are expected in the year 2000, 85 percent will occur in Africa, South-East Asia and the western Pacific.

The rising tuberculosis incidence and death toll are related, in large part, to the spread of the HIV epidemic. Other factors include poverty, crowding, homelessness, malnutrition, immigration and the lack of effective tuberculosis control programs. As the prevalence of tuberculosis increases, so

does the prevalence of drug-resistant mycobacterium. In some countries primary resistance may reach 25 percent and acquired resistance 75 percent.¹⁵¹

Nevertheless the overall mortality of adequately treated tuberculosis remains low. Prior to the availability of effective antimycobacterial chemotherapy, 50 percent of patients died within 2 years, 25 percent remained chronically infected and acted as reservoirs of infection within the community, and 25 percent experienced a spontaneous cure. Now, with good therapy, 98 percent of patients are cured, 1.2 percent still die and 0.8 percent continue to have chronic active disease.¹⁵¹ Recommended treatment regimens for non-resistant cases of *Mycobacterium tuberculosis* consist of (1) 9 months of isoniazid and rifampin or (2) 2 months of isoniazid, rifampin and pyrazinamide followed by 4 months of isoniazid and rifampin.¹⁵²

Risk Selection

Persons infected with *Mycobacterium tuberculosis*, as evidenced by a positive tuberculin skin test, but who do not have active disease, are at an increased risk of developing active tuberculosis some time in the future unless prophylactic treatment is given. The annual risk of developing active tuberculosis is relatively small and has been estimated to be 0.5–1 cases per 1000 per year. The risk is substantially higher, however, if infection has been acquired within the preceding 2 years (recent converters) or if there are coexisting conditions such as intravenous drug abuse, silicosis, chronic use of immunosuppressant agents or oral or parenteral corticosteroids, diabetes, end-stage renal disease, Hodgkin's disease or other hematologic malignancies, gastrectomy, HIV infection, malnutrition or substantial rapid weight loss.¹⁵³ It should also be noted that false-positive reactions to tuberculin occur, notably among persons who have received immunization with BCG.

Two insured lives mortality studies of tuberculosis have been reported. The *Medical Impairment Study 1983* reported on 49 511 life insurance policies issued on persons having a history or evidence of tuberculosis. The average annual death rate in this cohort compared with that of standard insured lives was 0.3 per 1000 per year for men

and 0.7 per 1000 per year for women. Relative mortality ratios were 107 percent and 127 percent for men and women respectively.

In a study of male policyholders issued life insurance by the Prudential Assurance Company of London who were coded as having pulmonary tuberculosis at the time of policy issue, no excess mortality was seen in those coded as having mild disease. Those coded as having moderate or severe disease and those who were treated with pneumothorax or surgery had mortality ratios ranging between 121 percent and 144 percent compared with those of standard insured lives.

SPECIAL CLINICAL PROBLEMS

Hemoptysis

The significance and the mortality implications of hemoptysis depend on the underlying cause. The amount of blood produced may range from a few flecks to cupfuls. With few exceptions, every case of new or appreciable hemoptysis should be carefully evaluated. Some of the underlying causes of hemoptysis are listed in Table 23.16.

Table 23.16. Selected causes of hemoptysis.

Naso-pharyngeal or upper airway bleeding
Chronic bronchitis
Bronchiectasis
Bronchial adenoma
Bronchogenic carcinoma
Tuberculosis
Bacterial pneumonia
Lung abscess
Fungal infections
Pulmonary infarction
Pulmonary arterio-venous fistula
Congestive heart failure
Mitral stenosis
Goodpasture's syndrome
Wegener's granulomatosis
Bleeding diatheses

Solitary Pulmonary Nodule (SPN)

The incidental finding of a SPN on a chest radiograph is not uncommon. A SPN is defined as a

well circumscribed radiographic density less than 3 cm in diameter that is completely surrounded by aerated lung. The differential diagnosis of a SPN is long and includes bronchogenic and metastatic carcinoma (see Table 23.17) Densities 3 cm or

greater in diameter have been termed solitary pulmonary masses and are more likely to be malignant. Features favoring benign and malignant processes are listed in Table 23.18. Any solitary non-calcified nodule in a person over 40 should be considered as possibly malignant until proven otherwise.

Table 23.17. Selected causes of SPN.

Bronchogenic carcinoma
Metastatic cancer
Granulomatous disease
Tuberculosis
Coccidiomycosis
Histoplasmosis
Resolving pneumonia
Lung abscess
Rounded atelectasis
Hamartoma
Pulmonary infarction
Pulmonary arteriovenous malformation
Pulmonary contusion
Pulmonary sequestration
Bronchogenic cyst
Wegener's granulomatosis
Kaposi's sarcoma
<i>Pneumocystis carinii</i> pneumonia

Table 23.18. Characteristics of SPN.

Suggestive of benign process	Suggestive of malignancy
Age under 35	Age over 35
No use of tobacco	Tobacco use
Diameter \leq 2 cm	Diameter $>$ 2 cm
Central calcification, lamellar or punctate calcification	No calcification or eccentric calcification
Smooth well defined margins	Lobulated or irregular margin
No enlargement for 2 or more years	Documented enlargement
Very rapid growth*	Intermediate growth rate
No prior history of malignancy	Prior history of malignancy
CT density $>$ 185 Hounsfield units	

* Very rapid growth is suggestive of infection or inflammatory process.

REFERENCES

- 1 Kochanek KD, Judson BL. Advance report of final mortality statistics, 1992. *Monthly Vital Statistics Report* 1995; 43: 1-73, 1995.
- 2 Campbell CD. Overview of community acquired pneumonia. *Med Clin North Am* 1994; 78: 1035-48.
- 3 Anthonison NR. Home oxygen therapy in chronic obstructive pulmonary disease. *Clin Chest Med* 1986; 7: 673-8.
- 4 Clini E, Vitacca M, Foglio K *et al.* Long-term home care programmes may reduce hospital admissions in COPD with chronic hypercapnia. *Minerva Anesthesiologica* 1996; 62(3): 57-64.
- 5 Dallari R, Barozzi G, Pinelli G *et al.* Predictors of survival in subjects with chronic obstructive pulmonary disease treated with long-term oxygen therapy. *Respiration* 1994; 1(1): 8-13.
- 6 Strom K, Boman G: Long-term oxygen therapy in parenchymal lung disease: an analysis of survival. *Eur Respir J* 1993; 8: 1264-70.
- 7 *American Medical Association Guide to the Evaluation of Permanent Impairment*. 4th edn. American Medical Association, 1993; 5, 162.
- 8 Kory RC. The Veterans Administration Army Cooperative study of pulmonary function. I. Clinical spirometry in normal men. *Am J Med* 1961; 3: 243-58.
- 9 Morris JF, Koski A, Johnson LC. Spirometric standards for healthy nonsmoking adults. *Am Rev Respir Dis* 1971; 103: 57-67.
- 10 Crapo RO, Morris AH, Gardner RM. Reference spirometric values using techniques and equipment that meet ATS recommendations. *Am Rev Respir Dis* 1981; 123: 659-64.
- 11 Knudson RJ, Lebowitz MD, Holberg CJ *et al.* Changes in the normal maximal expiratory flow-volume curve with growth and aging. *Am Rev Respir Dis* 1983; 127: 725-34.
- 12 Enright PL, Kronmal RA, Higgins M *et al.* Spirometry reference values for men and women 65 to 85 years of age: Cardiovascular Health Study. *Am Rev Respir Dis* 1993; 147: 125-33.

- 13 Petty TL. It's never too late to stop smoking. *JAMA* 1993; 269: 2785.
- 14 Bang KM, Gergen PJ, Kramer R *et al.* The effect of pulmonary impairment on all-cause mortality in a national cohort. *Chest* 1993; 103: 536-40.
- 15 Lange P, Nyboe J, Jensen G *et al.* Ventilatory function impairment and risk of cardiovascular death and of fatal or non-fatal myocardial infarction. *Eur Respir J* 1991; 4: 1080-7.
- 16 Peto R, Speizer FE, Cochrane AL, *et al.* The relevance in adults of air-flow obstruction, but not of mucous hypersecretion, to mortality from chronic lung disease. *Am Rev Respir Dis* 1983; 128: 491-500.
- 17 Sorlie PD, Kannel WB, O'Connor G. Mortality associated with respiratory function and symptoms of advanced age. *Am Rev Respir Dis* 1989; *et al.* 140: 379-84.
- 18 Beaty TH, Newill CA, Cohen BH *et al.* Effects of pulmonary function on mortality. *J Chron Dis* 1985; 38: 703-10.
- 19 Tockman MS, Pearson JD, Fleg JL *et al.* Rapid decline in FEV₁, a new risk factor for coronary heart disease mortality. *Am J Respir Crit Care Med* 1995; 151: 390-8.
- 20 Kuller L, Ockene J, Meilahn E *et al.* Relation of forced expiratory volume in one second (FEV₁) to lung cancer mortality in the multiple risk factor intervention trial. *Am J Epidemiol* 1990; 132: 265-72.
- 21 Pinkham A. Personal communication. Lincoln National Reinsurance, 1996.
- 22 Pare P. Conference summary: asthma structure and function. *Chest* 1995; 107 (suppl): 163S-169s.
- 23 Bone RC. Goals in asthma management, a step-care approach. *Chest* 1996; 109: 1056-65.
- 24 National Asthma Education Program Expert Panel Report: guidelines for the diagnosis and management of asthma. 1991 USDHHS (PHS) Publication number 1991-3042.
- 25 Woolcock AJ. Asthma. In: Murray JF, Nadel JA (eds). *Textbook of Respiratory Medicine*. 2nd edn. Philadelphia: WB Saunders Company, 1994; 1288-91.
- 26 Weiss ST. The origins of childhood asthma. *Monaldi Archives for Chest Disease* 1994; 49: 154-8.
- 27 Byrd JC. Environmental tobacco smoke. *Med Clin North Am* 1992; 76: 377-98.
- 28 Kivity S, Shochat Z, Bressler R *et al.* The characteristics of bronchial asthma among a young adult population. *Chest* 1995; 108: 24-7.
- 29 Banner AS. The increase in asthma prevalence. *Chest* 1995; 108: 301-2.
- 30 Wobig EK, Rosen R. Death from asthma: rare but real. *J Emerg Med* 1996: 223-40.
- 31 Weiss KB, Gergen PJ, Wagener DK. Breathing better or wheezing worse: the changing epidemiology of asthma morbidity and mortality. *Ann Rev Publ Health* 1993; 14: 491-513.
- 32 Taylor DR, Sears MR, Cockcroft DW. The beta-agonist controversy. *Med Clin North Am* 1996; 80: 719-43.
- 33 Leikauf GD, Kline S, Albert RE *et al.* Evaluation of a possible association of urban air toxins and asthma. *Environ Health Perspect* 1995; 103 (suppl 6): 253-71.
- 34 Koren HS. Associations between criteria air pollutants and asthma. *Environ Health Perspect* 1995; 103 (suppl 6): 235-42.
- 35 Weiss K, Wagener DK. Changing patterns of asthma mortality: identifying target populations at high risk. *JAMA* 1990; 264: 1683-7.
- 36 De Palo VA, Mayo PH, Friedman P *et al.* Demographic influences on asthma hospital admission rates in New York City. *Chest* 1994; 106: 447-51.
- 37 Gottlieb DJ, Beiser AS, O'Connor GT. Poverty, race and medication use are correlates of asthma hospitalization rates, a small area analysis in Boston. *Chest* 1995; 108: 28-35.
- 38 Greenberger PA, Miller TP, Lifschultz B. Circumstances surrounding deaths from asthma in Cook county (Chicago), Illinois. *Allergy Proc* 1993; 14: 321-7.
- 39 Ulrik CS, Frederiksen J. Mortality and markers of risk of asthma death among 1075 outpatients with asthma. *Chest* 1995; 108: 10-15.
- 40 Silverstein MD, Reed CE, O'Connell EJ *et al.* Long-term survival of a cohort of community residents with asthma. *N Engl J Med* 1994; 331: 1573-41.
- 41 Marquette CH, Saulnier F, Leroy O *et al.* Long-term prognosis of near-fatal asthma: a 6-year follow-up study of 145 asthmatic patients who underwent mechanical ventilation for a near-fatal attack of asthma. *Am Rev Respir Dis* 1992; 146: 76-81.
- 42 *Medical Impairment Study* 1983. Vol 1. Boston: Society of Actuaries and Association of Life Insurance Medical Directors of America, 1986; 74-5.
- 43 Fessel M. Mortality of policyholders suffering from chronic obstructive pulmonary disease. 36th meeting of Swiss Life Assurance Medical Directors, Zurich, 1984. Cited in: Mills RJ, Brackenridge RDC. Respiratory disorders. In: Brackenridge RDC, Elder WJ (eds). *Medical Selection of Life Risks*. 3rd edn. New York: Stockton Press, 1992; 489.

- 44 Pinkham A. Personal communication. Lincoln National Reinsurance, 1993
- 45 Picado. *J Asthma* 1989; 26: 231-6.
- 46 Ernst P, Spitzer WO, Suissa S *et al*. Risk of fatal and near-fatal asthma in relation to inhaled corticosteroid use. *JAMA* 1992; 268: 3462-4.
- 47 Bond WS. Toxic reactions and side effects of glucocorticoids in man. *Am J Hosp Pharmacy* 1977; 34: 479-85.
- 48 Woolcock AJ, Jenkins CR. Assessment of bronchial responsiveness as a guide to prognosis and therapy in asthma. *Med Clin North Am* 1990; 74: 753-5.
- 49 US Public Health Service: Smoking and Health. A report of the Advisory Committee to the Surgeon General of the Public Health Service. US Dept of Health, Education, and Welfare, Public Health Service, Center for Disease Control. PHS (PN) 1103, 1964.
- 50 US Dept of Health, Education, and Welfare. The Health Consequences of Smoking. Chronic Obstructive Lung Disease. US Department of Health and Human Services, Public Health Service, Office of Smoking. Rockville, MD 20857, DHHS (PHS) 84-50205, 1984.
- 51 Sherman CB. The health consequences of smoking. *Med Clin North Am* 1992; 76: 355-75.
- 52 Dockery DW, Speizer FE, Ferris BG *et al*. Cumulative and reversible effects of lifetime smoking on simple tests of lung function in adults. *Am Rev Respir Dis* 1988; 137: 286-92.
- 53 Burrows B, Knudson RJ, Cline MG *et al*. Quantitative relationships between cigarette smoking and ventilatory function. *Am Rev Respir Dis* 1977; 115: 195-205.
- 54 Beck GJ, Doyle CA, Schachter EN. Smoking and lung function. *Am Rev Respir Dis* 1981; 123: 149-55.
- 55 Burchfiel CM, Marcus EB, Curb D *et al*. Effects of smoking cessation on longitudinal decline in pulmonary function. *Am J Respir Crit Care Med* 1995; 151: 1778-85.
- 56 Snider GL, Faling LJ, Rennard SJ. Chronic bronchitis and emphysema. In: Murray JF, Nadel JA (eds). *Textbook of Respiratory Medicine*. 2nd edn Philadelphia: WB Saunders Company, 1994: 1342-3.
- 57 Thun MJ, Day-Lally CA, Calle EE. Excess mortality among cigarette smokers: changes in a 20 year interval. *Am J Public Health* 1995; 85: 1223-30.
- 58 Snider GL, Faling LJ, Rennard SJ. Chronic bronchitis and emphysema. In: Murray JF, Nadel JA (eds). *Textbook of Respiratory Medicine*. 2nd edn Philadelphia: WB Saunders Company, 1994; 1344.
- 59 Eriksson S.A 30 year perspective on alpha 1 antitrypsin deficiency. *Chest* 1996; 110: 237s-242s.
- 60 Clausen JL. The diagnosis of emphysema, chronic bronchitis, and asthma. *Clin Chest Med* 1990; 11: 405-16.
- 61 Burrows B. The course and prognosis of different forms of chronic airways obstruction in a sample from the general population. *N Engl J Med* 1987; 317: 1309-14.
- 62 Butz RH. Mortality associated with chronic pulmonary disease. *J Insur Med* 1988; 20: 46-7.
- 63 *Medical Impairment Study 1983*. Vol 1. Boston: Society of Actuaries and Association of Life Insurance Medical Directors of America, 1986; 76-7.
- 64 Fessel M. Mortality of policyholders suffering from chronic obstructive pulmonary disease. 36th meeting of Swiss Life Assurance Medical Directors, Zurich, 1984. Cited in: Mills RJ, Brackenridge RDC. Respiratory disorders. In: Brackenridge RDC, Elder WJ (eds). *Medical Selection of Life Risks*. 3rd edn. New York: Stockton Press, 1992; 485.
- 65 Clarke RD. Mortality of impaired lives. *J Inst Actuar*. Cited in: Mills RJ, Brackenridge RDC. Respiratory disorders. In: Brackenridge RDC, Elder WJ (eds). *Medical Selection of Life Risks*. 3rd edn. New York: Stockton Press, 1992; 485.
- 66 Fessel M. Chronic obstructive pulmonary disease in European insureds. *J Insur Med* 1991; 23: 38-9.
- 67 Luce JM. Bronchiectasis. In: Murray JF, Nadel JA (eds). *Textbook of Respiratory Medicine* 2nd edn. Philadelphia: WB Saunders Company, 1994; 1402.
- 68 Selman-Lama M, Padilla RP. Airflow obstruction and airway lesions in hypersensitivity pneumonitis. *Clin Chest Med* 1993; 14: 699-714.
- 69 Epler GR, Colby TV, McLoud TG *et al*. Bronchiolitis obliterans organizing pneumonia. *N Engl J Med* 1985; 312: 152-8.
- 70 Wells AU, du Bois RM. Bronchiolitis is association with connective tissue disorders. *Clin Chest Med* 1993; 14: 657-9.
- 71 Lazarus SC. Disorders of the intrathoracic airways. In: Murray JF, Nadel JA (eds). *Textbook of Respiratory Medicine*. 2nd edn. Philadelphia: WB Saunders Company, 1994; 1476.
- 72 Clark JG, Crawford SW, Madtes DK *et al*. Obstructive lung disease after allogenic marrow transplantation. *Ann Intern Med* 1989; 111: 368-76.
- 73 Fraser RG, Pare JAP, Pare PD *et al*. *Diagnosis of Diseases of the Chest*. 3rd edn. Philadelphia: WB Saunders Company, 1990; 2212.
- 74 Zielenski J, Tsui LC. Cystic fibrosis: genotypic and phenotypic variations. *Annu Rev Genet* 1995; 26: 777-807.

- 75 Stern RC. The diagnosis of cystic fibrosis. *New Engl J Med* 1997; 336: 487-91.
- 76 Tsui L. The cystic fibrosis transmembrane conductance regulator gene. *Am J Respir Crit Care Med* 1995; 151: S47-S53.
- 77 Egan TM, Detterbeck FC, Mill MR. Improved results of lung transplantation for patients with cystic fibrosis. *J Thorac Cardiovasc Surg* 1995; 109: 224-35.
- 78 Whitehead BF, Rees PG, Sorensen K *et al.* Results of heart-lung transplantation in children with cystic fibrosis. *Eur J Cardiothorac Surg* 1995; 9: 1-6.
- 79 Frederiksen B, Lanng S, Koch C *et al.* Improved survival in the Danish center-treated cystic fibrosis patients: results of aggressive treatment. *Pediatr Pulmonol.* 1996; 21: 153-8.
- 80 Liebow AA, Carrington CB. The interstitial pneumonias. In: Simon M, Potchen EJ, LeMay M (eds). *Frontiers of Pulm Radiol.* New York: Grune & Stratton, 1969; 102.
- 81 Carrington CB, Gaensler EA, Coutu RE *et al.* Natural history and treated course of usual and desquamative interstitial pneumonia. *N Engl J Med* 1978; 298: 801-9.
- 82 Erbes R, Shaberg T, Loddenkemper R. Lung function tests in patients with idiopathic pulmonary fibrosis: are they helpful for predicting outcome? *Chest* 1997; 111: 51-7.
- 83 Fraser RG, Pare JAP, Pare PD *et al.* *Diagnosis of Diseases of the Chest.* 3rd edn. Philadelphia: WB Saunders Company, 1990; 2611.
- 84 Newman LS, Rose CS, Maier LA. Sarcoidosis. *N Engl J Med* 1997; 336: 1224-35.
- 85 Fanburg BL, Lazarus DS. Sarcoidosis. In: Murray JF, Nadel JA (eds). *Textbook of Respiratory Medicine.* 2nd edn. Philadelphia: WB Saunders Company, 1994; 1873-88.
- 86 Siltzbach LE, James DG, Turiaf J *et al.* Course and prognosis of sarcoidosis around the world. *Am J Med* 1974; 57: 847-52.
- 87 Sharma OP, Maheshwari A, Thaker K. Myocardial sarcoidosis. *Chest* 1993; 253-8.
- 88 James DG. Granulomatous liver disease. In: Schiff L, Schiff ER (eds). *Diseases of the Liver.* 7th edn. Philadelphia: JB Lippincott Company, 1993; 1500.
- 89 *Disease of the Liver and Biliary System.* 9th edn. London: Blackwell Scientific Publications, 1993; 463.
- 90 Sharma OP. Pulmonary sarcoidosis and corticosteroids. *Am Rev Respir Dis* 1993; 147: 1598-1600.
- 91 Fraser RG, Pare JAP, Pare PD *et al.* *Diagnosis of Diseases of the Chest.* 3rd edn. Philadelphia: WB Saunders Company, 1990; 2646.
- 92 Douglas NJ, Goetzel EJ. Pulmonary eosinophilia and eosinophilic granuloma. In: Murray JF, Nadel JA (eds). *Textbook of Respiratory Medicine.* 2nd edn. Philadelphia: WB Saunders Company, 1994; 1926-9.
- 93 McHugh LG, Milberg JA, Whitcomb ME *et al.* Recovery of function in survivors of the acute respiratory distress syndrome. *Am J Respir Crit Care Med* 1994; 150: 90-4.
- 94 Peters JI, Bell RC, Prihoda TJ *et al.* Clinical determinants of abnormalities in pulmonary function in survivors of the adult respiratory distress syndrome. *Am Rev Respir Dis* 1989; 139: 1163-8.
- 95 Elliott CG. Pulmonary sequelae in survivors of the adult respiratory distress syndrome. *Clin Chest Med* 1990; 11: 789-800.
- 96 Zapol WM, Trelstad RC, Coffey JW *et al.* Pulmonary fibrosis in severe acute respiratory failure. *Am Rev Respir Dis* 1979; 119: 547-55.
- 97 Wang BM, Stern EJ, Schmidt RA *et al.* Diagnosing pulmonary alveolar proteinosis. *Chest* 1997; 111: 460-6.
- 98 Wasserman K, Mason GR. Pulmonary alveolar proteinosis. In: Murray JF, Nadel JA (eds). *Textbook of Respiratory Medicine* Philadelphia: WB Saunders Company, 1994; 1933-46.
- 99 Rosenstock L, Cullen MR. *Textbook of Clinical Occupational and Environmental Medicine.* Philadelphia: WB Saunders Company, 1994; 195.
- 100 Rom WN (ed). *Environmental and Occupational Medicine.* 2nd edn. Boston: Little, Brown and Co, 1992, 1-1493.
- 101 Murray JF, Nadel JA. *Textbook of Respiratory Medicine.* 2nd edn. Philadelphia: WB Saunders Company, 1994; 1977.
- 102 Nicholson WJ, Perkel G, Selikoff IJ. Occupational exposure to asbestos: population at risk and projected mortality, 1980-2030. *Am J Ind Med* 1982; 3: 259-311.
- 103 Copes R, Thomas D, Becklake MR. Temporal patterns of exposure and nonmalignant pulmonary abnormality in Quebec chrysotile workers. *Arch Environ Health* 1985; 40: 80-87.
- 104 Dreessen WC, Dallavalle JM, Edwards TI *et al.* A study of asbestosis in the asbestos textile industry. Public Health Bulletin No. 241. US Govt Printing Office, Washington, 1938.
- 105 Murray JF, Nadel JA (eds). *Textbook of Respiratory Medicine.* 2nd edn. Philadelphia: WB Saunders Company, 1994; 1962.
- 106 Rosenstock L, Cullen MR (eds). *Textbook of Clinical Occupational and Environmental Medicine.* Philadelphia: WB Saunders Company, 1994; 265.

- 107 Kleinerman J. Pathology standards for coalworkers' pneumoconiosis. Report of Pneumoconiosis Committee of the College of American Pathologists. *Arch Pathol Lab Med* 1979; 103: 375-85.
- 108 Rom WN. Basic mechanisms leading to focal emphysema in coal workers' pneumoconiosis. *Environ Res* 1990; 53: 16-28.
- 109 US Dept of Health and Human Services. Work-Related Lung Disease Surveillance Report. Public Health Service, Centers for Disease Control, National Institute for Occupational Safety and Health Publication No. 91-113, 1991; 78.
- 110 Morgan WKC, Seaton A. *Occupational Lung Diseases*. 2nd edn. Philadelphia: WB Saunders Company, 1984; 334.
- 111 Pneumoconiosis redefined (editorial). *BMJ* 1972; 2: 552.
- 112 Attfield MD, Seixas NS. Prevalence of pneumoconiosis and its relationship to dust exposure in a cohort of US bituminous coal miners and examiners. *Am J Ind Med* 1995; 27: 137-51.
- 113 Kuempel ED, Sayner LT, Attfield MD *et al*. Exposure-response analysis of mortality among coal miners in the United States. *Am J Ind Med* 1995; 28: 167-84.
- 114 Perez-Padilla R, Salas J, Chapela R *et al*. Mortality in Mexican patients with chronic pigeon breeder's lung compared with those with usual interstitial pneumonia. *Am Rev Respir Dis* 1993; 148: 49-53.
- 115 Schwarz MI, Cherniak RM, King TE. Diffuse alveolar hemorrhage and other rare infiltrative disorders. In: Murray JF, Nadel JA (eds). *Textbook of Respiratory Medicine*. 2nd edn. Philadelphia: WB Saunders Company, 1994; 1899.
- 116 Matteson EL, Gold KN, Bloch DA *et al*. Long-term survival of patients with Wegener's granulomatosis from the American College of Rheumatology Wegener's granulomatosis classification criteria cohort. *Am J Med* 1996; 101: 129-34.
- 117 Pope-Harmon AL, Davis WB, Allen ED *et al*. Acute eosinophilic pneumonia. A summary of 15 cases and review of the literature. *Medicine* 1996; 75(6): 334-42.
- 118 Cheon JE, Leeks, Chung MH *et al*. Acute eosinophilic pneumonia: radiographic and CT findings in six patients. *AJR Am J Roentgenol* 1996; 167(5): 1195-9.
- 119 King MA, Pope-Harmon AL, Allen ED *et al*. Acute eosinophilic pneumonia: radiologic and clinical features. *Radiology* 1997; 203(3): 715-19.
- 120 Chumbley LC, Harrison EG, DeRemee RA: Allergic granulomatosis and angitis. *Mayo Clin Proc* 1977; 52: 477-84.
- 121 Fraser RG, Pare JAP, Pare PD *et al*. *Diagnosis of Diseases of the Chest*. 3rd edn. Philadelphia: WB Saunders Company, 1990; 1826.
- 122 Butler J, Agostoni P. Cor pulmonale. In: Murray JF, Nadel JA (eds). *Textbook of Respiratory Medicine*. 2nd edn. Philadelphia: WB Saunders Company, 1994; 1778.
- 123 Rubin LJ. Pulmonary vasculitis and primary pulmonary hypertension. In: Murray JF, Nadel JA (eds). *Textbook of Respiratory Medicine*. 2nd edn. Philadelphia: WB Saunders Company, 1994; 1693.
- 124 Abenhaim L, Moride Y, Brenot F *et al*. Appetite-suppressant drugs and the risk of primary pulmonary hypertension. *N Engl J Med* 1996; 336: 609-16.
- 125 D'Alonzo GE, Barst RJ, Ayres SM *et al*. Survival in patients with primary pulmonary hypertension. *Ann Intern Med* 1991; 115: 343-9.
- 126 Lombard RM, Zwillich CW. Medical therapy of obstructive sleep apnea. *Med Clin N Am* 1985; 69(6): 1325.
- 127 Bradley TD, Rutherford R, Grossman RF *et al*. Role of daytime hypoxemia in the pathogenesis of right heart failure in the obstructive sleep apnea syndrome. *Am Rev Respir Dis* 1985; 131: 835-9.
- 128 Fletcher EC, Schaaf JW, Miller J *et al*. Long-term cardiopulmonary sequelae in patients with sleep apnea and chronic lung disease. *Am Rev Respir Dis* 1987; 135: 525-33.
- 129 Weitzenblum E, Krieger J, Apprill M *et al*. Daytime pulmonary hypertension in patients with obstructive sleep apnea syndrome. *Am Rev Respir Dis* 1988; 138: 345-59.
- 130 Parish JM, Shepard JW. Cardiovascular effects of sleep disorders. *Chest* 1990; 97(5): 1220-6.
- 131 Shepard JW. Cardiovascular consequences of obstructive sleep apnea. *Clin Chest Med* 1992; 13(3): 442-9.
- 132 Hedner J. Nocturnal hypertension — under the cover of darkness. *Blood Press* 1995; 4: 197-8.
- 133 Noda A, Okad T, Yasuma F *et al*. Cardiac hypertrophy in obstructive sleep apnea syndrome. *Chest* 1995; 107: 1538-44.
- 134 Klitzman D, Miller A. Obstructive sleep apnea syndromes. *Mount Sinai J Med* 1994; 61: 113-21.
- 135 Findley L, Unverzadt M, Suratt P. Automobile accidents involving patients with obstructive sleep apnea. *Am Rev Respir Dis* 1988; 138: 337-40.

- 136 Stoohs RA, Guilleminault C, Itoi A *et al.* Traffic accidents in commercial long-haul truck drivers: the influence of sleep disordered breathing. *Sleep* 1994; 17: 619-23.
- 137 Gonzales-Rothi RJ, Foresman GE, Block AJ. Do patients with sleep apnea die in their sleep? *Chest* 1988; 94: 531-8.
- 138 Partinen M, Jamieson A, Guilleminault C. Long-term outcome for obstructive sleep apnea syndrome patients. *Chest* 1988; 94: 1200-4.
- 139 He J, Kryger MH, Zorick FJ *et al.* Mortality in obstructive sleep apnea. *Chest* 1988; 94: 9-14.
- 140 Lavie P, Herer P, Peled R *et al.* Mortality in sleep apnea patients: a multivariate analysis of risk factors. *Sleep* 1995; 18(3): 149-57.
- 141 Bliwise DL, Bliwise NG, Partinen M *et al.* Sleep apnea and mortality in an aged cohort. *Am J Public Health* 1988; 78: 544-7.
- 142 Pollack CP, Perlick D, Linsner JP *et al.* Sleep problems in the community elderly as predictors of death and nursing home placement. *J Community Health* 1990; 15: 123-35.
- 143 Mant A, King M, Saunders NA *et al.* Four-year follow-up of mortality and sleep related respiratory disturbance in non-demented seniors. *Sleep* 1995; 18(6): 433-8.
- 144 Fleury B *et al.* Long-term compliance to continuous positive airway pressure therapy (nCPAP) set up during a split night polysomnography. *Sleep* 1994; 17: 512-15.
- 145 Hoffstein V, Viner S, Mateika S *et al.* Treatment of obstructive sleep apnea with nasal continuous positive airway pressure. *Am Rev Respir Dis* 1992; 145: 841-5.
- 146 Brackenridge RDC. Respiratory system diseases. In: Lew EA, Gajewski J (eds). *Medical Risks: trends in mortality by age and time elapsed*. New York: Praeger, 1990; 8-60 to 8-61.
- 147 Brackenridge RDC. Respiratory system diseases. In: Lew EA, Gajewski J (eds). *Medical Risks: trends in mortality by age and time elapsed*. New York: Praeger, 1990; 8-56 to 8-57.
- 148 Donowitz GR, Mandell GL. Acute pneumonia. In: Mandell GL, Bennet JE, Dolin R (eds). *Principles and Practice of Infectious Disease*. 4th edn. New York: Churchill Livingstone, 1995; 629.
- 149 Finegold S. Lung abscess. In: Mandell GL, Bennet JE, Dolin R (eds). *Principles and Practice of Infectious Disease*. 4th edn. New York: Churchill Livingstone, 1995; 646.
- 150 Bryant RE. Pleural effusion and empyema. In: Mandell GL, Bennet JE, Dolin R (eds). *Principles and Practice of Infectious Disease* 4th edn. New York: Churchill Livingstone, 1995; 640.
- 151 Enarson DA, Murray JF. Global epidemiology of tuberculosis. In: Rom WM, Garay SM (eds). *Tuberculosis*. Boston: Little, Brown and Co, 1996; 57-75.
- 152 American Thoracic Society. Treatment of tuberculosis and tuberculosis infection in adults and children. *Am J Respir Crit Care Med* 1994; 149: 1359-75.
- 153 Colice GL. Decision analysis, public health policy, and isoniazid chemoprophylaxis for young adult tuberculin skin reactors. *Arch Intern Med* 1990; 150: 2517-22.