

Reprinted from

**BULLETIN**

Volume **5** (1971)

of the Australian Mathematical Society

Number **2**

C. C. Heyde and J. R. Leslie Alan Lambert	<i>Improved classical limit analogues for Galton-Watson processes with or without immigration</i>	145
A. Brown	<i>Unitary equivalence and reducibility of invertibly weighted shifts</i>	157
J. R. Giles	<i>On the differential equations for tide-well systems</i>	175
Kurt Mahler	<i>A Banach space with support homeomorphism is reflexive</i>	187
Alan Brace and D. E. Daykin	<i>An arithmetic remark on entire periodic functions</i>	191
Jennifer Wallis	<i>A finite set covering theorem</i>	197
Masako Izumi and Shin-ichi Izumi	<i>A skew-Hadamard matrix of order 92</i>	203
A. R. Amir-Moéz and C. R. Perry	<i>Integrability of trigonometric series III</i>	205
J. B. Diaz and R. Výborný	<i>Remarks on theorems of Thompson and Freede</i>	221
M. R. Vaughan-Lee	<i>On some mean value theorems of the differential calculus</i>	227
John A. Beachy	<i>On product varieties of groups</i>	239
J. R. Blake	<i>Cotorsion radicals and projective modules</i>	241
R. M. Bryant and L. G. Kovács	<i>Self propulsion due to oscillations on the surface of a cylinder at low Reynolds number</i>	255
C. Christensen	<i>A note on generalized characters</i>	265
Susan Dancs	<i>Rings with a few more zero-divisors</i>	271
J. N. Ward	<i>The sub-near-field structure of finite near-fields</i>	275
Abstracts of Australasian Paul Pentony	<i>On finite groups admitting automorphisms with nilpotent fixed-point group</i>	281
Jennifer Wallis	<i>Ph.D. theses</i>	
B. E. Clancy	<i>Laws in torsion-free nilpotent varieties with particular reference to the laws of free nilpotent groups</i>	283
	<i>Combinatorial matrices</i>	285
	<i>Multigroup neutron transport theory in plane geometry</i>	287

---

Bull. Austral. Math. Soc. Vol. 5 (1971) No. 2, pp. 145-288

THE UNIVERSITY OF QUEENSLAND PRESS  
ST LUCIA, QUEENSLAND 4067

## **THE AUSTRALIAN MATHEMATICAL SOCIETY**

President: G. E. Wall      University of Sydney  
Secretary: W. Pye        Secondary Teachers' College, Victoria, 3052  
Treasurer: V. G. Hart    University of Queensland  
(These officers serve until May 1972)

### **Membership and correspondence**

Applications for membership, notices of change of address or title or position, and other correspondence, except as noted below, should be sent to:

The Secretary, Australian Mathematical Society,  
Secondary Teachers' College,  
Swanston Street, Parkville, Victoria 3052, Australia.

Subscriptions and correspondence related to accounts should be addressed to:

The Treasurer, Australian Mathematical Society,  
Department of Mathematics,  
University of Queensland,  
St Lucia, Queensland 4067, Australia.

Correspondence about the distribution of the Society's JOURNAL and BULLETIN should be addressed to:

The Business Manager, Australian Mathematical Society,  
Department of Mathematics,  
University of Queensland,  
St Lucia, Queensland 4067, Australia.

The annual subscription is \$50 for Institutions and \$16 in 1971 for Ordinary Members. Junior Members (undergraduate and junior graduate students) may ask to have their subscription reduced to \$6 in 1971. Members of other mathematical societies with which the Australian Mathematical Society has entered into reciprocity agreements are admitted to membership and its privileges at half the current rates, if they are not normally resident in Australia.

Members of the Society receive copies of the JOURNAL free, and may subscribe to the BULLETIN at the price of \$4 per volume (\$8 p.a.). Members may elect to receive the BULLETIN instead of the JOURNAL.

### **The JOURNAL of the Australian Mathematical Society**

The JOURNAL began publication in 1959. From 1972, two volumes of four parts will be published annually, at a price of \$25 per volume. Subscription orders and orders for back numbers should be sent to the Business Manager (address above). Manuscripts for publication should be sent to the Editor:

Professor B. Mond,  
Department of Mathematics,  
La Trobe University,  
Bundoora, Victoria 3083, Australia,

or to one of the Associate Editors listed in a recent part of the JOURNAL.

*(Continued inside back cover)*

## **The BULLETIN of the Australian Mathematical Society**

The BULLETIN began publication in 1969. Two volumes of three numbers each are published annually, at a subscription of \$25 in 1972. Subscription orders and orders for back numbers should be sent to the Business Manager (address inside front cover). Manuscripts for publication should be sent to the Editor:

Professor B. H. Neumann,  
Department of Mathematics,  
Institute of Advanced Studies,  
Australian National University,  
POB 4, Canberra, ACT 2600, Australia,

or to the Associate Editor, Dr L. G. Kovács, at the same address. Authors should keep a complete copy of their manuscripts, because manuscripts are not normally returned to them.

Authors receive 50 free offprints of their papers and may order more at cost. Correspondence about proofs and offprint orders should be addressed to one of the Assistant Editors:

Associate Professor A. McMullen, or	Dr S. Yamamuro,
Department of Mathematics,	Department of Mathematics,
Royal Military College,	Institute of Advanced Studies,
Duntroon, ACT 2600, Australia	Australian National University,
	POB 4, Canberra, ACT 2600, Australia.

### **Editorial Policy**

The BULLETIN of the Australian Mathematical Society aims at quick publication of original research in all branches of Mathematics. Only manuscripts that are suitable for quick publication will be accepted, and only if space permits. Manuscripts must be in their final, publishable form when submitted; refereeing will be kept to a minimum, and no revision of manuscripts will be allowed.

The BULLETIN is printed by a photo-offset process from a carefully prepared typescript. The type founts and special symbols available are, accordingly, restricted.

To facilitate the work of reviewing journals and the Mathematical Offprint Service (MOS) of the American Mathematical Society, the BULLETIN publishes an abstract at the head of each paper that is not itself an abstract, and also a tentative MOS classification in the top right-hand corner of the first page of each paper. Authors are invited to assist by supplying abstracts of their papers and supplying or checking MOS classification numbers. Copies of the MOS classification scheme are available to authors on request from MOS, American Mathematical Society, POB 6248, Providence, R.I. 02904, USA. The scheme and instructions for its use can be also found as an appendix to Mathematical Reviews, Index to Volume 39 (1970).

### **Copyright**

The material in this BULLETIN may be quoted in full or in part with due acknowledgement of the source.

The BULLETIN of the Australian Mathematical Society is published for the Society by

The University of Queensland Press,  
St Lucia, Queensland 4067,  
Australia,

and printed by

Watson Ferguson & Co.,  
Brisbane, Queensland 4101,  
Australia.

# Improved classical limit analogues for Galton-Watson processes with or without immigration

C.C. Heyde and J.R. Leslie

It has recently emerged that the central limit theorem and iterated logarithm law for random walk processes have natural counterparts for Galton-Watson processes with or without immigration. Much of the work on these counterparts has previously involved the imposition of supplementary moment conditions. In this paper we show how to dispense with these supplementary conditions and in so doing make the analogy with the random walk results complete.

## 1. Introduction

Let  $Z_0 = 1, Z_1, Z_2, \dots$  denote a super-critical Galton-Watson process with  $1 < EZ_1 = m$  and  $0 < \text{var}Z_1 = \sigma^2 < \infty$ . It is well-known that there exists a non-degenerate random variable  $W$  such that  $\lim_{n \rightarrow \infty} W_n = W$

almost surely, where  $W_n = m^{-n}Z_n$  (for example, Harris [2], p. 13).

Furthermore, some central limit analogues have been established in this context by Heyde [5] and Bühler [1]. These results are that

$(m^2 - m)^{\frac{1}{2}} \sigma^{-1} Z_n^{-\frac{1}{2}} m^n (W - W_n)$  conditional on  $Z_n > 0$  and

$(m^2 - m)^{\frac{1}{2}} \sigma^{-1} m^{-\frac{1}{2}j} (m^j - 1)^{-\frac{1}{2}} Z_n^{-\frac{1}{2}} \left( Z_{n+j} - m^j Z_n \right)$  conditional on  $Z_n > 0$  (fixed  $j$ )

are both asymptotically normal  $N(0, 1)$ . See [5] for an explanation of

why these results can be regarded as central limit analogues.

Under the further restriction that  $EZ_1^3 < \infty$ , rates of convergence in the central limit analogues cited above have been given in [7]. These have been used in [6] to obtain almost sure convergence results for the Galton-Watson process which are analogues of the law of the iterated logarithm for random walks. In this paper we shall show how the restriction that  $EZ_1^3 < \infty$  may be removed. In Section 2 we shall obtain convergence rates in the central limit analogues and also iterated logarithm analogues under the basic condition that  $EZ_1^2 < \infty$ .

In Section 3 of this paper we shall deal with the Galton-Watson process with immigration. The development of the corresponding limit results in this case has followed the pattern described previously for the case without immigration. Heyde and Seneta [8] have obtained central limit analogues under  $EZ_1^2 < \infty$  and rate results and iterated logarithm analogues under  $EZ_1^3 < \infty$ . We shall again show how to dispense with the moment restriction and will obtain the rate results and iterated logarithm analogues under  $EZ_1^2 < \infty$ .

## 2. The process without immigration

We shall establish the following theorems. The reader is referred to the papers [5], [6] and [7] for background details.

**THEOREM 1.** *Let  $1 < m = EZ_1$  and  $0 < \text{var}Z_1 = \sigma^2 < \infty$ . Then*

$$\sup_x \left| P \left[ \left( (m^2 - m)^{\frac{1}{2}} \sigma^{-1} u_{Z_n}^{-1} Z_n^{-\frac{1}{2}} m^n (W - \bar{W}_n) \leq x \mid Z_n > 0 \right) - \Phi(x) \right] \leq c_n \right.$$

and

$$\sup_x \left| P \left[ \left( \sigma_r^{-1} v_{Z_n}^{-1} Z_n^{-\frac{1}{2}} \left( Z_{n+r} - m^r Z_n \right) \leq x \mid Z_n > 0 \right) - \Phi(x) \right] \leq d_n \right.,$$

where  $\{c_n\}$ ,  $\{d_n\}$  are certain sequences of positive constants satisfying

$$\sum_{n=1}^{\infty} c_n < \infty \quad \text{and} \quad \sum_{n=1}^{\infty} d_n < \infty. \quad \text{Here}$$

$$\sigma_r^2 = \text{var}Z_r = \sigma^2 m^r (m^r - 1) (m^2 - m)^{-1},$$

$r$  any fixed integer,

$$u_n = \int_{|x| < \sqrt{n}} x^2 dP \left( \sigma^{-1} (m^2 - m)^{\frac{1}{2}(W-1)} \leq x \right)$$

$$v_n = \int_{|x| < \sqrt{n}} x^2 dP \left( \sigma_r^{-1} \left( Z_r - m^r \right) \leq x \right)$$

and  $\Phi(x)$  is the distribution function of  $N(0, 1)$ .

Explicit forms for  $c_n$  and  $d_n$  can be found by applying the lemma below. We note also that  $u_n \uparrow 1$  and  $v_n \uparrow 1$  as  $n \rightarrow \infty$ .

**THEOREM 2.** Suppose that  $1 < m = EZ_1$  and  $0 < \text{var}Z_1 = \sigma^2 < \infty$ . Then, on the non-extinction set  $\{W > 0\}$  we have almost surely

$$\limsup_{n \rightarrow \infty} \frac{Z_{n+r} - m^r Z_n}{\left( 2\sigma_r^2 Z_n \log n \right)^{\frac{1}{2}}} = 1, \quad \liminf_{n \rightarrow \infty} \frac{Z_{n+r} - m^r Z_n}{\left( 2\sigma_r^2 Z_n \log n \right)^{\frac{1}{2}}} = -1,$$

and

$$\limsup_{n \rightarrow \infty} \frac{m^{nW} - Z_n}{\left( 2\sigma^2 (m^2 - m)^{-1} Z_n \log n \right)^{\frac{1}{2}}} = 1, \quad \liminf_{n \rightarrow \infty} \frac{m^{nW} - Z_n}{\left( 2\sigma^2 (m^2 - m)^{-1} Z_n \log n \right)^{\frac{1}{2}}} = -1,$$

where  $r$  is any fixed positive integer.

Theorems 1 and 2 extend the scope of results given in Heyde and Brown [7] and in Heyde [6] respectively under the additional condition that  $EZ_1^3 < \infty$ . The form of the bounds obtained in Theorem 1 is however, of necessity, much more complicated in the general case. An explanation of this is not difficult to deduce from results given in [3] and [4]. Our Theorem 2 preserves exactly the form of the Theorem of [6] under the more general conditions.

In order to establish the above results we need the following key lemma. The result of the lemma is given in two parts; the first is needed in the present section and the second to obtain corresponding results for the process with immigration in Section 3.

**LEMMA.** Let  $\xi_i$ ,  $i = 1, 2, 3, \dots$  be independent and identically

distributed random variables with  $E(\xi_1) = 0$  and  $\text{var}\xi_1 = \alpha^2 < \infty$ . Let  $N_n$  be a positive integer valued random variable which is independent of the  $\{\xi_i\}$ . Then,

$$(1) \quad \sup_x \left| P\left(\alpha^{-1} d_{N_n}^{-1} N_n^{-\frac{1}{2}} (\xi_1 + \dots + \xi_{N_n}) \leq x\right) - \Phi(x) \right| \leq AE\left(N_n^{-\frac{1}{2}} a_{N_n}\right) + BE\left(N_n^{\frac{1}{2}} b_{N_n}\right) + E\left(N_n c_{N_n}\right)$$

where  $A, B$  are positive constants and

$$a_n = \int_{|x| < \sqrt{n}} |x|^3 dP\left(\alpha^{-1} \xi_1 \leq x\right), \quad b_n = \int_{|x| \geq \sqrt{n}} |x| dP\left(\alpha^{-1} \xi_1 \leq x\right),$$

$$c_n = P\left(\alpha^{-1} |\xi_1| > \sqrt{n}\right), \quad d_n^2 = \int_{|x| < \sqrt{n}} x^2 dP\left(\alpha^{-1} \xi_1 \leq x\right).$$

If  $\eta_n$  with  $E|\eta_n| < \infty$  is a random variable which is independent of the  $\{\xi_i\}$  and of  $N_n$ , then for any sequence  $\{\epsilon_n\}$  of positive constants with  $\epsilon_n \rightarrow 0$  as  $n \rightarrow \infty$ ,

$$(2) \quad \sup_x \left| P\left(\alpha^{-1} d_{N_n}^{-1} N_n^{-\frac{1}{2}} (\xi_1 + \dots + \xi_{N_n} + \eta_n) \leq x\right) - \Phi(x) \right| \leq AE\left(N_n^{-\frac{1}{2}} a_{N_n}\right) + BE\left(N_n^{\frac{1}{2}} b_{N_n}\right) + E\left(N_n c_{N_n}\right) + \alpha^{-1} \epsilon_n^{-1} E|\eta_n| E\left(N_n^{-\frac{1}{2}} d_{N_n}^{-1}\right) + \frac{1}{2} \epsilon_n.$$

**Proof.** Let

$$e_n^2 = \int_{|x| < \sqrt{n}} x^2 dP\left(\alpha^{-1} \xi_1 \leq x\right) - \left( \int_{|x| < \sqrt{n}} x dP\left(\alpha^{-1} \xi_1 \leq x\right) \right)^2.$$

We have

$$(3) \quad \sup_x \left| P\left(\alpha^{-1} d_{N_n}^{-1} N_n^{-\frac{1}{2}} (\xi_1 + \dots + \xi_{N_n}) \leq x \mid N_n = k\right) - \Phi(x) \right| \leq \sup_x \left| P\left(\alpha^{-1} e_{N_n}^{-1} N_n^{-\frac{1}{2}} (\xi_1 + \dots + \xi_{N_n}) \leq x \mid N_n = k\right) - \Phi(x) \right| + \sup_x \left| \Phi(x) - \Phi\left(e_k d_k^{-1} x\right) \right|.$$

Also, using the mean value theorem,



$$\begin{aligned}
(4) \quad \sup_x \left| \Phi(x) - \Phi\left(e_k d_k^{-1} x\right) \right| &\leq c \left(1 - e_k d_k^{-1}\right) \\
&= c d_k^{-1} (e_k + d_k)^{-1} \left( \int_{|x| < \sqrt{k}} x dP\left(\alpha^{-1} \xi_1 \leq x\right) \right)^2 \\
&= c d_k^{-1} (e_k + d_k)^{-1} \left( \int_{|x| \geq \sqrt{k}} x dP\left(\alpha^{-1} \xi_1 \leq x\right) \right)^2 \\
&\leq c_1 b_k^2 \leq c_1 b_k,
\end{aligned}$$

$c, c_1$  being positive constants. Furthermore, from (22) of Heyde [4] we find that

$$\begin{aligned}
(5) \quad \sup_x \left| P\left(\alpha^{-1} e_N^{-1} N^{-\frac{1}{2}} \left(\xi_1 + \dots + \xi_{N_n}\right) \leq x \mid N_n = k\right) - \Phi(x) \right| \\
\leq A k^{-\frac{1}{2}} a_k + B_1 k^{\frac{1}{2}} b_k + k c_k,
\end{aligned}$$

so that using (3), (4) and (5),

$$\begin{aligned}
(6) \quad \sup_x \left| P\left(\alpha^{-1} d_N^{-1} N^{-\frac{1}{2}} \left(\xi_1 + \dots + \xi_{N_n}\right) \leq x \mid N_n = k\right) - \Phi(x) \right| \\
\leq A k^{-\frac{1}{2}} a_k + B k^{\frac{1}{2}} b_k + k c_k.
\end{aligned}$$

The result (1) follows readily from (6) using the argument of the lemma in §4 of Heyde and Brown [7]. (2) is obtained using exactly the method of Lemma 2.1 of Heyde and Seneta [8] with the aid of (1) instead of the Berry-Esseen bound.

**Proof of Theorem 1.** Suppose that  $Z_n^*$  has the distribution of  $Z_n$  conditional on  $Z_n > 0$ . We firstly note that (see [5], [7]), conditional on  $Z_n > 0$ ,  $m^n Z_n^{-\frac{1}{2}} (W - W_n)$  has the same distribution as  $(Z_n^*)^{-\frac{1}{2}} \left( U_1 + \dots + U_{Z_n^*} \right)$ , where the  $U_i$  are independent of  $Z_n^*$  and are independent and identically distributed, each with the distribution of  $W - 1$ . Also, conditional on  $Z_n > 0$ ,  $Z_n^{-\frac{1}{2}} \left( Z_{n+r} - m^r Z_n \right)$  has the same distribution as  $(Z_n^*)^{-\frac{1}{2}} \left( V_1 + \dots + V_{Z_n^*} \right)$  where the  $V_i$  are independent of

$Z_n^*$  and are independent and identically distributed, each with the distribution of  $Z_r - m^r$ . We can thus apply the lemma in both cases and obtain bounds which we call  $c_n, d_n$  respectively. It remains to show that

$\sum_{n=1}^{\infty} c_n < \infty, \sum_{n=1}^{\infty} d_n < \infty$ . We shall indicate the proof for  $\sum_{n=1}^{\infty} c_n$ ; that

for  $\sum_{n=1}^{\infty} d_n$  follows similarly.

What we have to demonstrate is that  $\sum_{n=1}^{\infty} E\left\{(Z_n^*)^{-\frac{1}{2}} a_{Z_n^*}\right\} < \infty$ ,

$\sum_{n=1}^{\infty} E\left\{(Z_n^*)^{\frac{1}{2}} b_{Z_n^*}\right\} < \infty$  and  $\sum_{n=1}^{\infty} E\left\{Z_n^* c_{Z_n^*}\right\} < \infty$  where  $a_n, b_n, c_n$  are defined

in the lemma with  $\xi_i$  having the distribution of  $W - 1$ . The proofs of the convergence of these three series are identical in form. They depend on results of [4] where it is, in essence, shown in the proof of Theorem 4 that under the conditions of the theorem and if  $\{n_k, k = 1, 2, 3, \dots\}$  is a sequence of integers with  $n_k \sim Kc^{2k}$  as  $k \rightarrow \infty$  ( $K > 0, c > 1$ ), then

$$\sum_{k=1}^{\infty} n_k^{-\frac{1}{2}} a_{n_k} \leq K_1 < \infty, \sum_{k=1}^{\infty} n_k^{\frac{1}{2}} b_{n_k} \leq K_2 < \infty, \sum_{k=1}^{\infty} n_k c_{n_k} \leq K_3 < \infty$$

for certain  $K_1, K_2, K_3$  independent of  $K$ .

For  $u > 0$ , let

$$a_u = \int_{|x| < \sqrt{u}} |x|^3 dP\left\{\sigma^{-1}(m^2 - m)^{\frac{1}{2}}(W - 1) \leq x\right\}.$$

We have

$$\begin{aligned}
E\left((Z_n^*)^{-\frac{1}{2}} a_{Z_n^*}\right) &= \int_0^\infty (xm^n)^{-\frac{1}{2}} a_{xm^n} dP\left(m^{-n} Z_n^* \leq x\right) \\
&= \sum_{k=0}^\infty \int_{k \leq x < k+1} (xm^n)^{-\frac{1}{2}} a_{xm^n} dP\left(m^{-n} Z_n^* \leq x\right) \\
&\leq \sum_{k=0}^\infty k^{-\frac{1}{2}} m^{-\frac{1}{2}n} a_{(k+1)m^n} P\left(k \leq m^{-n} Z_n^* < k+1\right) \\
&\leq c_1 \sum_{k=0}^\infty [(k+1)m^{n+1}]^{-\frac{1}{2}} a_{[(k+1)m^{n+1}]} P\left(k \leq m^{-n} Z_n^*\right),
\end{aligned}$$

where  $c_1$  is a suitable positive constant and  $[x]$  denotes the integer part of  $x$ . Then using Chebyshev's inequality,  $P\left(m^{-n} Z_n^* \geq k\right) \leq ck^{-2}$ , and

$$\begin{aligned}
\sum_{n=1}^\infty E\left((Z_n^*)^{-\frac{1}{2}} a_{Z_n^*}\right) &\leq c_1 \sum_{n=1}^\infty \sum_{k=1}^\infty [km^{n+1}]^{-\frac{1}{2}} a_{[km^{n+1}]} P\left(k-1 \leq m^{-n} Z_n^*\right) \\
&\leq c_1 \sum_{n=1}^\infty [m^{n+1}]^{-\frac{1}{2}} a_{[m^{n+1}]} \\
&\quad + c_1 c \sum_{k=2}^\infty (k-1)^{-2} \sum_{n=1}^\infty [km^{n+1}]^{-\frac{1}{2}} a_{[km^{n+1}]} \\
&\leq c_2 + c_3 \sum_{k=2}^\infty (k-1)^{-2} < \infty,
\end{aligned}$$

as required. That  $\sum_{n=1}^\infty E\left((Z_n^*)^{\frac{1}{2}} b_{Z_n^*}\right) < \infty$  and  $\sum_{n=1}^\infty E\left(Z_n^* c_{Z_n^*}\right) < \infty$  follow in the same fashion.

**Proof of Theorem 2.** This follows the same lines as the proof of the theorem of [6]. We just make use of Theorem 1 and the inequality (6) instead of the results based on the Berry-Esseen inequality employed in [6]. We then obtain

$$\limsup_{n \rightarrow \infty} \frac{Z_{n+r}^{-m^n} Z_n}{\left(2\sigma_r^2 v_r^2 Z_n \log n\right)^{\frac{1}{2}}} = 1, \quad \liminf_{n \rightarrow \infty} \frac{Z_{n+r}^{-m^n} Z_n}{\left(2\sigma_r^2 v_r^2 Z_n \log n\right)^{\frac{1}{2}}} = -1$$

almost surely on  $\{W > 0\}$ . The required results for

$\left(Z_{n+r}^{-m^n} Z_n\right) \left(2\sigma_r^2 Z_n \log n\right)^{-\frac{1}{2}}$  then follow since  $v_n \uparrow 1$  as  $n \rightarrow \infty$ . We have, for example, since  $Z_n \rightarrow \infty$  almost surely on  $\{W > 0\}$ ,

$$\begin{aligned} \limsup_{n \rightarrow \infty} \frac{Z_{n+r} - m^r Z_n}{\left(2\sigma_r^2 Z_n \log n\right)^{\frac{1}{2}}} &\leq \limsup_{n \rightarrow \infty} \frac{Z_{n+r} - m^r Z_n}{\left(2\sigma_r^2 v_{Z_n}^2 Z_n \log n\right)^{\frac{1}{2}}} \\ &\leq \limsup_{n \rightarrow \infty} \frac{Z_{n+r} - m^r Z_n}{\left(2\sigma_r^2 Z_n \log n\right)^{\frac{1}{2}}} \limsup_{n \rightarrow \infty} \frac{1}{v_{Z_n}} \\ &= \limsup_{n \rightarrow \infty} \frac{Z_{n+r} - m^r Z_n}{\left(2\sigma_r^2 Z_n \log n\right)^{\frac{1}{2}}} \end{aligned}$$

on  $\{W > 0\}$ . The remainder of the proof goes through exactly as in [6]. We show

$$\limsup_{n \rightarrow \infty} \frac{|Z_n - m^n W|}{\left(2\sigma^2 (m^2 - m)^{-1} u_{Z_n}^2 Z_n \log n\right)^{\frac{1}{2}}} \leq 1 ,$$

from which it follows that

$$\limsup_{n \rightarrow \infty} \frac{|Z_n - m^n W|}{\left(2\sigma^2 (m^2 - m)^{-1} Z_n \log n\right)^{\frac{1}{2}}} \leq 1 ,$$

and the remainder of the argument of [6] can be repeated word for word.

### 3. The process with immigration

Let  $X_0 = 1, X_1, X_2, \dots$  denote a Galton-Watson process with immigration whose offspring distribution has the distribution of  $Z_1$  with  $1 < EZ_1 = m$  and  $0 < \text{var}Z_1 = \sigma^2 < \infty$ . We shall also suppose that the immigration distribution has a finite mean. We refer the reader to Seneta [9] for a detailed description of the process. Under the present conditions, the theorem of [9] ensures that  $m^{-n} X_n$  converges almost surely to a proper random variable  $V$  with finite mean  $EV$  and such that  $P(V = 0) = 0$ . We shall here obtain the following theorems which extend results presented in Heyde and Seneta [8].

**THEOREM 3.** *Let  $1 < m = EZ_1$  and  $0 < \text{var}Z_1 = \sigma^2 < \infty$ . Then*

$$\sup_x \left| P \left( (m^2 - m)^{\frac{1}{2}} \sigma^{-1} u_n^{-1} X_n^{-\frac{1}{2}} (m^n V - X_n) \leq x \mid X_n > 0 \right) - \Phi(x) \right| \leq \alpha_n$$

and

$$\sup_x \left| P \left( \sigma_r^{-1} v_n^{-1} X_n^{-\frac{1}{2}} (X_{n+r} - m^r X_n) \leq x \mid X_n > 0 \right) - \Phi(x) \right| \leq \beta_n,$$

where  $\{\alpha_n\}$ ,  $\{\beta_n\}$  are certain sequences of positive constants satisfying

$\sum_{n=1}^{\infty} \alpha_n < \infty$  and  $\sum_{n=1}^{\infty} \beta_n < \infty$ . Here  $\sigma_r$ ,  $u_n$ ,  $v_n$  are as defined in the statement of Theorem 1.

Explicit forms for  $\alpha_n$  and  $\beta_n$  can be found by applying the lemma.

**THEOREM 4.** Suppose that  $1 < m = EZ_1$  and  $0 < \text{var}Z_1 = \sigma^2 < \infty$ .

Then, with probability one,

$$\limsup_{n \rightarrow \infty} \frac{X_{n+r} - m^r X_n}{\left( 2\sigma_r^2 X_n \log n \right)^{\frac{1}{2}}} = 1, \quad \liminf_{n \rightarrow \infty} \frac{X_{n+r} - m^r X_n}{\left( 2\sigma_r^2 X_n \log n \right)^{\frac{1}{2}}} = -1,$$

$$\limsup_{n \rightarrow \infty} \frac{m^n V - X_n}{\left( 2\sigma^2 (m^2 - m)^{-1} X_n \log n \right)^{\frac{1}{2}}} = 1, \quad \liminf_{n \rightarrow \infty} \frac{m^n V - X_n}{\left( 2\sigma^2 (m^2 - m)^{-1} X_n \log n \right)^{\frac{1}{2}}} = -1,$$

where  $r$  is any fixed positive integer.

Theorems 3 and 4 extend the scope of results given in Theorems 2 and 3 of [8] under the additional condition that  $EZ_1^3 < \infty$ . The form of the bounds obtained in the present Theorem 3 is however, of necessity, much more complicated in the more general case. The present Theorem 4 preserves exactly the form of Theorem 3 of [8] under the more general conditions.

The proofs of Theorems 2 and 3 follow the same lines as those of Theorems 1 and 2, again using the lemma and (6). We make use of the representations

$$m^n V - X_n = (W^{(1)} - 1) + \dots + \left( W^{(X_n)} - 1 \right) + I^{(n)} \quad \text{a.s.}$$

and

$$X_{n+r} - m^r X_n = \left( Z_r^{(1)} - m^r \right) + \dots + \left( Z_r^{(X_n)} - m^r \right) + Y_{r,n}$$

of [8] and the points noted in the proof of Theorem 3 of [8]. The only real point of difference in the proofs involves showing that we can choose

a sequence  $\{\epsilon_n\}$  with  $\epsilon_n \rightarrow 0$  as  $n \rightarrow \infty$  such that  $\sum_{n=1}^{\infty} \epsilon_n < \infty$  and

$$\sum_{n=1}^{\infty} \epsilon_n^{-1} E \left[ X_n^{-\frac{1}{2} w_{X_n}^{-1}} \mid X_n > 0 \right] < \infty \text{ where } w_n \text{ is either } u_n \text{ or } v_n.$$

We know that  $u_n \uparrow 1$ ,  $v_n \uparrow 1$  so that  $w_n \uparrow 1$  as  $n \rightarrow \infty$ . Thus, conditional on  $X_n > 0$ ,  $w_{X_n} \geq w_1$  and hence

$$\begin{aligned} \sum_{n=1}^{\infty} \epsilon_n^{-1} E \left[ X_n^{-\frac{1}{2} w_{X_n}^{-1}} \mid X_n > 0 \right] &\leq w_1^{-1} \sum_{n=1}^{\infty} \epsilon_n^{-1} E \left[ X_n^{-\frac{1}{2}} \mid X_n > 0 \right] \\ &\leq w_1^{-1} \sum_{n=1}^{\infty} \epsilon_n^{-1} \left( E \left[ X_n^{-1} \mid X_n > 0 \right] \right)^{\frac{1}{2}}. \end{aligned}$$

Now, using Lemma 2.3 of [8] we have  $\epsilon_n = \theta^n$  with  $0 < \gamma < \theta < 1$ ,

$$\sum_{n=1}^{\infty} \epsilon_n^{-1} \left( E \left[ X_n^{-1} \mid X_n > 0 \right] \right)^{\frac{1}{2}} \leq \sum_{n=1}^{\infty} \theta^{-n} \gamma^n < \infty.$$

Thus, with this choice of  $\epsilon_n$ ,  $\sum_{n=1}^{\infty} \epsilon_n < \infty$  and

$$\sum_{n=1}^{\infty} \epsilon_n^{-1} E \left[ X_n^{-\frac{1}{2} w_{X_n}^{-1}} \mid X_n > 0 \right] < \infty, \text{ as required.}$$

### References

- [1] Wolfgang J. Bühler, "Ein zentraler Grenzwertsatz für Verzweigungsprozesse", *Z. Wahrscheinlichkeitstheorie verw. Gebiete* 11 (1969), 139-141.

- [2] Theodore E. Harris, *The theory of branching processes* (Die Grundlehren der Mathematischen Wissenschaften in Einzeldarstellungen, Band 119. Springer-Verlag, Berlin, Göttingen, Heidelberg, 1963; Prentice-Hall, Englewood Cliffs, New Jersey, 1963).
- [3] C.C. Heyde, "On the influence of moments on the rate of convergence to the normal distribution", *Z. Wahrscheinlichkeitstheorie verw. Gebiete* 8 (1967), 12-18.
- [4] C.C. Heyde, "Some properties of metrics in a study on convergence to normality", *Z. Wahrscheinlichkeitstheorie verw. Gebiete* 11 (1969), 181-192.
- [5] C.C. Heyde, "Some central limit analogues for super-critical Galton-Watson processes", *J. Appl. Probability* 8 (1971), 52-59.
- [6] C.C. Heyde, "Some almost sure convergence theorems for branching processes", *Z. Wahrscheinlichkeitstheorie verw. Gebiete* (to appear).
- [7] C.C. Heyde and B.M. Brown, "An invariance principle and some convergence rate results for branching processes", *Z. Wahrscheinlichkeitstheorie verw. Gebiete* (to appear).
- [8] C.C. Heyde and E. Seneta, "Analogues of classical limit theorems for the supercritical Galton-Watson process with immigration", *Math. Biosci.* (to appear).
- [9] E. Seneta, "A note on the supercritical Galton-Watson process with immigration", *Math. Biosci.* 6 (1970), 305-312.

Department of Statistics,  
School of General Studies,  
Australian National University,  
Canberra, ACT.