
Dennstaedtiaceae

Pteridium aquilinum (L.) Kuhn subsp. ***aquilinum*** (*Pteris aquilina* L.) - clearings and abandoned cultivated land - CC

Dioscoreaceae

Dioscorea communis (L.) Caddick & Wilkin (*Tamus communis* L.) - woods - C

Dryopteridaceae

Dryopteris filix-mas (L.) Schott (*Polypodium filix-mas* L.; *Aspidium filix-mas* (L.) Sw.; *Nephrodium filix-mas* (L.) Rich.; *Lastrea filix-mas* (L.) C. Presl) - woods - C

Dryopteris pallida (Bory) Maire & Petitm. subsp. ***pallida*** (*D. villarii* (Bellardi) Woynar subsp. *pallida* (Bory) Heywood; *Nephrodium pallidum* Bory) - shady cliffs, usually to 1,000 m - PC

Dryopteris submontana (Fraser-Jenk. & Jermy) Fraser-Jenk. (*D. villarii* Bellardi subsp. *submontana* Fraser-Jenk. & Jermy) - shady stony slopes - RR - Cava di Rena! (Conti and Soldati 2010).

Dryopteris villarii (Bellardi) Woyn. ex Schinz & Thell. subsp. ***villarii*** (*Polypodium villarii* Bellardi) - screes, cool montane stony slopes - PC

Polystichum aculeatum (L.) Roth (*Polypodium aculeatum* L.; *Dryopteris lobata* (Huds.) Schinz & Thell.; *D. aculeata* Kuntze subsp. *lobata* (Huds.) Briq.; *Aspidium aculeatum* (L.) Sw.) - woods - C

Polystichum lonchitis (L.) Roth (*Polypodium lonchitis* L.) - cool *Fagus sylvatica* woods and shady cliffs, usually over 1,500 m - C

Polystichum setiferum (Forssk.) Woyn. (*Polypodium setiferum* Forssk.) - woods - CC