



Hanly et al., [1] conducted a cohort study using a validated questionnaire—EPIC [1] (n = 21). There is significant loss to follow-up thus limiting the strength of the findings. It is overall a paper of moderate quality, based on sample size, loss to follow-up and strength of conclusions drawn. It is evident that is psychosexual care impacts on quality of life and needs to be addressed by clinicians (Level 2b, recommendation A). The requirement for psychosexual care is confirmed by [1, 2, 3], demonstrating the need for psychosexual care. Although these are qualitative papers of moderate quality [4], this theme of psychosexual care is an important one to address with this study.

---

## References

1. Hanly N, Mireskandari S, Juraskova I. The struggle towards ‘the New Normal’: a qualitative insight into psychosexual adjustment to prostate cancer. *BMC Urol.* 2014;14:56.
2. O’Brien R, Rose P, Campbell C, Weller D, Neal RD, Wilkinson C, McIntosh H, Watson E. “I wish I’d told them”: a qualitative study examining the unmet psychosexual needs of prostate cancer patients during follow-up after treatment. *Patient Educ Couns.* 2011;84:200–7.
3. O’Brien R, Rose PW, Campbell C, Weller D, Neal RD, Wilkinson C, Watson EK. Experiences of follow-up after treatment in patients with prostate cancer: a qualitative study. *BJU Int.* 2010;106:998–1003.
4. Mays N, Pope C, Popay J. Systematically reviewing qualitative and quantitative evidence to inform management and policy-making in the health field. *J Health Serv Res Policy.* 2005;10:6–20.