

# ERRATUM

W. Powell & P. D. S. Caligari, **The use of hormonal and osmotic growth retardants in media used for the storage of potato germplasm in-vitro.** *Potato Research* 32 (1989) 57 – 64

Table 4, in the 'Results', was omitted. This table is given below.

Table 4. The phenotypic correlation between the characters for the 23 cultivars. The four numbers in each group refer, from the top down, to media A, B, C and D.

	Vigour	Green	Number of shoots	Number of nodes	Usable nodes	Number of tubers	Root length	Fresh weight	Internode length
Green	0.95								
	0.89								
	0.84								
	0.84								
Number of shoots	0.39	0.36							
	0.33	0.42							
	0.73	0.62							
	-0.04	-0.04							
Number of nodes	0.89	0.87	0.15						
	0.83	0.89	0.42						
	0.64	0.53	0.58						
	0.72	0.55	-0.00						
Usable nodes	0.87	0.85	0.09	0.94					
	0.88	0.90	0.47	0.85					
	0.70	0.79	0.57	0.42					
	0.90	0.81	-0.08	0.82					
Number of tubers	-0.13	-0.16	-0.17	-0.13	-0.21				
	0.08	0.09	0.28	0.04	0.13				
	0.55	0.32	0.66	0.53	0.19				
	0.13	0.13	0.20	0.08	-0.00				
Root length	0.63	0.62	0.35	0.57	0.52	0.10			
	0.82	0.85	0.39	0.84	0.80	0.15			
	0.61	0.57	0.52	0.55	0.40	0.53			
	0.60	0.76	0.30	0.40	0.52	0.38			
Fresh weight	0.70	0.70	0.54	0.59	0.47	0.18	0.80		
	0.72	0.68	0.36	0.76	0.64	-0.11	0.74		
	0.62	0.56	0.79	0.48	0.58	0.69	0.68		
	0.74	0.61	0.07	0.71	0.68	0.52	0.65		
Internode length	0.68	0.69	0.55	0.60	0.64	-0.18	0.51	0.56	
	0.45	0.35	0.32	0.30	0.41	-0.04	0.42	0.60	
	-0.14	-0.20	0.11	0.03	-0.15	0.22	0.07	0.27	
	0.06	-0.02	0.45	0.02	0.14	-0.06	0.11	0.14	
Sq. SL	0.90	0.88	0.32	0.95	0.91	-0.19	0.61	0.67	0.79
	0.77	0.77	0.26	0.88	0.73	-0.08	0.77	0.90	0.57
	0.38	0.29	0.60	0.60	0.27	0.54	0.45	0.64	0.78
	0.58	0.37	-0.00	0.85	0.74	-0.10	0.27	0.63	0.42