

## Erratum to: Effects of pruning on knotty core taper and form of *Pinus radiata* and *Pinus pinaster*

Andrea Hevia<sup>1</sup> · Juan Gabriel Álvarez-González<sup>2</sup> · Juan Majada<sup>1</sup>

Published online: 7 April 2016  
© Springer-Verlag Berlin Heidelberg 2016

### Erratum to: Eur. J. Wood Prod. DOI 10.1007/s00107-016-1019-0

Due to an error during conversion of Geographic Units (Latitude, Longitude) from UTM coordinates, information given for location of the experimental plots in Table 1 of the manuscript is not correct.

The data of location of the plots should be replaced by:

Experimental plot	Location
<i>Pinus radiata</i>	
La Campa	43°24'N, 5°28'W
Cabada	43°25'N, 6°33'W
Santa Catalina	43°31'N, 6°6'W
Fabal	43°12'N, 6°53'W
<i>Pinus pinaster</i>	
Valseira	43°33'N, 6°12'W
Barcia	43°31'N, 6°28'W
Monteagudo	43°31'N, 6°6'W
Uria	43°4'N, 6°51'W

The online version of the original article can be found under doi:[10.1007/s00107-016-1019-0](https://doi.org/10.1007/s00107-016-1019-0).

✉ Andrea Hevia  
ahevia@cetemas.es

<sup>1</sup> Sustainable Forest Management Area, Forest and Wood Technology Research Centre (CETEMAS), Finca Experimental “La Mata” s/n, 33820 Grado, Spain

<sup>2</sup> Sustainable Forest Management Unit, Agroforestry Engineering Department, High Polytechnic School, University of Santiago de Compostela, Campus Universitario s/n, 27002 Lugo, Spain

Site Index has been recalculated at the time of establishment of the pruning plots. Both changes are not substantial for the study described and do not affect the results and final conclusions presented in this work.

On the next page, the whole Table is presented.

**Table 1** Details of the network of plots and data used in this study

Experimental plot	Location	Site Index* (m)	At time of installation		At time of inventory									
			Age (years)	Stocking (trees ha <sup>-1</sup> )	$H^*$ (m)		$D^*$ (cm)			$dos_j^*$ (cm)			$h_{DOS_j^*}$ (m)	
					Mean	Mean	Mean	Max	Min	Mean	Max	Min		
<i>Pinus radiata</i>														
La Campa	43°24'N, 5°28'W	22.2	9	1187	11.9	16.8	18.0	28.0	9.0	1.7	5.7	0.3		
Cabada	43°25'N, 6°33'W	26.1	7	1261	11.0	14.4	16.5	27.2	8.7	1.4	4.1	0.3		
Santa Catalina	43°31'N, 6°6'W	21.3	8	1440	10.8	15.4	16.7	26.9	9	1.2	3.6	0.3		
Fabal	43°12'N, 6°53'W	23.6	10	1356	11.4	14.8	16.9	28.3	8.1	1.3	4.0	0.3		
<i>Pinus pinaster</i>														
Valsera	43°33'N, 6°12'W	17.9	8	1352	9.99	14.8	16.4	24	8.9	1.3	3.7	0.2		
Barcia	43°31'N, 6°28'W	13.5	11	1379	7.9	13.2	15.9	22.5	9.0	1.3	2.8	0.3		
Monteagudo	43°31'N, 6°6'W	13.2	8	1670	6.7	12.9	15.0	22.5	10.1	0.96	2.3	0.3		
Uria	43°4'N, 6°51'W	15.9	7	1802	6.5	9.7	13.2	14.5	10.6	0.8	0.9	0.6		

\*  $H$  total tree height, m;  $D$  diameter outside bark at breast height (1.30 m above ground), cm;  $dos_j$  diameter over stubs at height of pruned whorl  $j$ , cm;  $h_{DOS_j}$  height above ground at specific point  $j$  on the stem, m. Site Index estimated using the equations defined by Canga (2008) and Álvarez-Álvarez et al. (2011) for *P. radiata* and *P. pinaster*, respectively