

## Erratum to: Risk of psychiatric treatment for mood disorders and psychotic disorders among migrants and Dutch nationals in Utrecht, The Netherlands

J. P. Selten · W. Laan · R. Kupka · H. M. Smeets ·  
J. van Os

Published online: 24 September 2014  
© Springer-Verlag Berlin Heidelberg 2014

**Erratum to: Soc Psychiatry Psychiatr Epidemiol (2012)**  
**47:271–278**  
**DOI 10.1007/s00127-010-0335-7**

In the original publication of this article, there were some errors in Tables 1 and 2. The corrected versions of the tables are shown here.

---

The online version of the original article can be found under  
doi:[10.1007/s00127-010-0335-7](https://doi.org/10.1007/s00127-010-0335-7).

---

J. P. Selten  
Rivierduinen Institute for Mental Health Care, Oegstgeest,  
The Netherlands

J. P. Selten (✉)  
De School, Endegeesterstraatweg 2, 2342 AJ Oegstgeest,  
The Netherlands  
e-mail: [j.selten@ggzleiden.nl](mailto:j.selten@ggzleiden.nl)

W. Laan · H. M. Smeets  
Julius Center for Health Sciences and Primary Care,  
University Medical Centre Utrecht, Utrecht, The Netherlands

R. Kupka  
Altrecht Institute for Mental Health Care, Utrecht,  
The Netherlands

J. P. Selten · J. van Os  
Department of Psychiatry and Psychology, South Limburg  
Mental Health Research and Teaching Network, EURON,  
Maastricht University Medical Centre, Maastricht, The  
Netherlands

**Table 1** Relative risks for ethnic minorities in the town of Utrecht, adjusted for age and sex, of treatment at a mental health institute for unipolar depressive disorder or bipolar disorder, 1 January 2002 to 31 December 2006

Ethnic group	Person-years M/F	Unipolar depressive disorder <sup>a</sup>			Bipolar disorder <sup>b</sup>		
		M/F	RR	(95 % CI)	M/F	RR	(95 % CI)
First generation, 15–65 years							
Dutch nationals <sup>c</sup>	334,624/357,508	563/976	1.0	Ref	142/163	1.0	Ref
Turkish	15,669/14,887	131/214	4.6	(4.1–5.2)	4/5	0.6	(0.3–1.2)
Moroccan	31,270/27,621	179/278	3.6	(3.2–3.9)	7/5	0.5	(0.3–0.9)
Surinamese	8,628/9,310	4/57	1.7	(1.3–2.1)	2/4	0.6	(0.3–1.3)
Western-European <sup>d</sup>	6,364/6,852	4/16	0.7	(0.4–1.1)	4/3	1.4	(0.6–2.9)
Second generation, 15–65 years							
Dutch nationals <sup>c</sup>	334,624/357,508	563/976	1.0	Ref	142/163	1.0	Ref
Turkish	5,445/5,125	16/48	5.0	(3.8–6.5)	1/3	2.7	(1.0–7.3)
Moroccan	6,919/6,674	8/34	2.7	(1.9–3.7)	2/0	1.1	(0.3–4.5)
Surinamese	4,160/4,247	5/21	2.1	(1.4–3.1)	1/3	2.3	(0.9–6.3)
Western-European <sup>e</sup>	8,432/8,927	15/27	1.1	(0.8–1.4)	4/3	0.8	(0.4–1.7)
First and second generation combined, 15–65 years							
Dutch nationals <sup>c</sup>	334,624/357,508	563/976	1.0	Ref	142/163	1.0	Ref
Turkish	21,114/20,112	147/262	4.9	(4.4–5.5)	5/8	0.8	(0.5–1.5)
Moroccan	38,189/34,295	187/312	3.6	(3.3–4.0)	9/5	0.5	(0.3–0.9)
Surinamese	12,788/13,557	29/78	1.8	(1.5–2.2)	3/7	0.8	(0.4–1.6)
Western-European	14,796/15,779	19/43	0.9	(0.7–1.2)	8/6	1.0	(0.6–1.7)

<sup>a</sup> The DSM-IV diagnoses major depressive disorder, single or recurrent episode (codes 296.2× or 296.3×)

<sup>b</sup> The DSM-IV diagnoses for bipolar I or II disorder or for bipolar disorder not otherwise specified (codes 296.0×, 296.4×, 296.5×, 296.6×, 296.7, 296.89, 296.80)

<sup>c</sup> Born in the Netherlands and both parents born in the Netherlands

<sup>d</sup> Born in France, UK, Germany or Belgium

<sup>e</sup> Born in the Netherlands and one or both parents born in France, UK, Germany or Belgium. If parents were born in different countries, country of birth of mother determines ethnicity

**Table 2** Relative risks for ethnic minorities in the town of Utrecht, adjusted for age and sex, of treatment at a mental health institute for non-affective psychotic disorder, 1 January 2002 to 31 December 2006

Ethnic group <sup>a</sup>	M/F	RR	(95 % CI)
First generation, 15–65 years			
Dutch nationals <sup>b</sup>	364/283	1.0	Ref
Turkish	28/14	1.3	(1.0–1.8)
Moroccan	106/22	2.3	(1.9–2.8)
Surinamese	46/24	3.4	(2.6–4.3)
Western European <sup>c</sup>	4/4	0.7	(0.3–1.4)
Second generation, 15–65 years			
Dutch nationals <sup>b</sup>	364/283	1.0	Ref
Turkish	18/4	3.6	(2.4–5.6)
Moroccan	17/4	2.8	(1.8–4.3)
Surinamese	13/11	4.3	(2.9–6.5)
Western European <sup>d</sup>	13/4	0.9	(0.6–1.5)
First and second generation combined, 15–65 years			
Dutch nationals <sup>b</sup>	364/283	1.0	Ref
Turkish	46/18	1.8	(1.4–2.4)
Moroccan	123/26	2.5	(2.1–3.0)
Surinamese	59/35	3.8	(3.0–4.7)
Western European	17/8	0.9	(0.6–1.3)

The DSM-IV diagnoses schizophrenia, schizophreniform disorder, schizoaffective disorder, delusional disorder, brief psychotic disorder or psychotic disorder not otherwise specified

<sup>a</sup> For person-years at risk: see Table 1

<sup>b</sup> Born in the Netherlands and both parents born in the Netherlands

<sup>c</sup> Born in France, UK, Germany or Belgium

<sup>d</sup> Born in the Netherlands and one or both parents born in France, UK, Germany or Belgium. If parents were born in different countries, country of birth of mother determines ethnicity