



Correction to: Impact of socioeconomic position and distance on mental health care utilization: a nationwide Danish follow-up study

Aake Packness¹  · Frans Boch Waldorff¹ · René dePont Christensen¹ · Lene Halling Hastrup² · Erik Simonsen^{2,5} · Mogens Vestergaard³ · Anders Halling⁴

Published online: 10 July 2018
© The Author(s) 2018

Correction to:
Soc Psychiatry Psychiatr Epidemiol (2017) 52:1405–1413
<https://doi.org/10.1007/s00127-017-1437-2>

In the original publication of this article, Table 3 was published incorrectly. The corrected table is shown below.

Open Access This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made.

The original article can be found online at <https://doi.org/10.1007/s00127-017-1437-2>.

✉ Aake Packness
apackness@health.sdu.dk

- ¹ Research Unit of General Practice, Institute of Public Health, University of Southern Denmark, J.B. Winsløvs Vej 9A, 5000 Odense C, Denmark
- ² Psychiatric Research Unit, Region of Zealand, Slagelse, Denmark
- ³ Department of Public Health, Institute of General Medical Practice, Aarhus University, Aarhus, Denmark
- ⁴ Centre for Primary Health Care Research, Lund University, Lund, Sweden
- ⁵ Faculty of Health and Medical Sciences, Institute of Clinical Medicine, University of Copenhagen, Copenhagen, Denmark

Table 3 Separate analyses of correlation of income or education with mental health service used, in odds ratios and incidence rate ratios of contact

Odds ratio of contact to services ^a																	
Logit	Contact out-patient psychiatrist			Contact psychologist			Contact GP-MHS			Psychiatric emergency clinic			Admission MH > 1 day				
	OR	P	CI	N	OR	P	CI	OR	P	CI	OR	P	CI	OR	P	CI	
<i>Family equivalent income^b</i>																	
Highest third	1			50,373	1			1			1			1			
Middle third	1.03	0.414	0.97	1.09	0.71	< 0.001	0.67	0.75	0.90	< 0.001	0.86	0.95	0.94	0.344	0.82	1.07	
Lower third	1.25	< 0.001	1.17	1.34	0.49	< 0.001	0.46	0.53	0.81	< 0.001	0.77	0.86	0.82	0.007	0.71	0.95	
<i>Education (years)^b</i>																	
12+	1			48,183	1			1			1			1			
10–12	0.92	0.005	0.86	0.97	0.67	< 0.001	0.63	0.71	0.89	< 0.001	0.85	0.94	0.99	0.888	0.86	1.13	
< 10	0.95	0.103	0.89	1.01	0.37	< 0.001	0.35	0.40	0.71	< 0.001	0.67	0.75	0.90	0.147	0.78	1.04	
Incidence rate ratios of contact among patients who had one visit or more ^a																	
Poisson	Out-patient psychiatrist			Psychologist			GP-MHS			Psych. emergency clinic			Admission MH > 1 day				
	IRR	P	CI	N	IRR	P	CI	N	IRR	P	CI	IRR	P	CI	IRR	P	CI
<i>Family equivalent income^b</i>																	
Highest third	1			11,113	1			9033	1			17,638	1			1752	1
Middle third	0.90	< 0.001	0.88	0.91	0.93	< 0.001	0.91	0.95	1.00	0.639	0.98	1.02	1.02	0.734	0.91	1.15	1.03
Lower third	0.83	< 0.001	0.81	0.84	0.94	< 0.001	0.91	0.96	0.94	< 0.001	0.92	0.97	1.06	0.364	0.94	1.14	0.95
<i>Education (years)^b</i>																	
12+	1			10,659	1			8869	1			17,038	1			1677	1
10–12	0.92	< 0.001	0.90	0.93	0.92	< 0.001	0.90	0.93	0.99	0.395	0.97	1.01	0.98	0.729	0.87	1.10	1.00
< 10	0.75	< 0.001	0.74	0.76	0.80	< 0.001	0.79	0.82	0.93	< 0.001	0.91	0.96	1.06	0.370	0.94	1.19	1.04

Psychiatric emergency clinic includes emergency contacts and admissions up to 1 day

Results significant within a 95% confidence interval are marked in bold

GP-MHS mental health services by general practitioner (talk therapy), Admission MH admission to mental hospital, IRR incidence rate ratio, CI confidence interval, P 0.05

^aTwo separate analyses of correlation of income or education with mental health service used

^bAdjusted for age, gender, country of origin, cohabitating status, access to vehicle, comorbidity psychiatric, comorbidity somatic