



Correction to: Measurements and Distribution of Atmospheric Particulate-Bound Mercury: A Review

Hui Zhang¹ · Xuewu Fu^{1,2} · Xun Wang^{1,4} · Xinbin Feng^{1,2,3}

Published online: 5 July 2019

© Springer Science+Business Media, LLC, part of Springer Nature 2019

Correction to:

Bulletin of Environmental Contamination and Toxicology
<https://doi.org/10.1007/s00128-019-02663-5>

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

The original version of this article unfortunately contained a mistake in units.

In section Global and Regional Distribution of PBM the unit pg m^{-3} was incorrectly written as ng m^{-3} in three occurrences. The revised sentence with correct unit of measurement is given below.

Mean PBM concentrations at urban sites ranged from 2.3 to 1180 pg m^{-3} (mean $225 \pm 174.5 \text{ pg m}^{-3}$), significantly higher than the PBM concentrations measured at remote sites (mean $25.3 \pm 29.7 \text{ pg m}^{-3}$).

The original article can be found online at <https://doi.org/10.1007/s00128-019-02663-5>.

✉ Hui Zhang
zhanghui1@mail.gyig.ac.cn

- ¹ State Key Laboratory of Environmental Geochemistry, Institute of Geochemistry, Chinese Academy of Sciences, Guiyang 550002, China
- ² CAS Center for Excellence in Quaternary Science and Global Change, Xi'an 710061, China
- ³ University of Chinese Academy of Sciences, Beijing 100049, China
- ⁴ College of Resources and Environment, Southwest University, Chongqing 400715, China