

CORRECTION



Correction to: Norepinephrine in septic shock

Glenn Hernández^{1*} , Jean-Louis Teboul^{2,3} and Jan Bakker^{1,4,5,6}

© 2019 Springer-Verlag GmbH Germany, part of Springer Nature

Correction to: *Intensive Care Med*

<https://doi.org/10.1007/s00134-018-5499-8>

The original version of this article unfortunately contained a mistake. There was an error in the sentence “Guinot et al. showed that NE can improve V–A coupling and stroke volume in hypotensive post-cardiac surgery patients, although stroke volume was found to increase only in patients with preserved coupling [12].”

The correct sentence should be “Guinot et al. showed that NE can improve V–A coupling and stroke volume in hypotensive post-cardiac surgery patients, although stroke volume was found to increase only in patients with non-preserved coupling [12].”

We apologize for this mistake.

Author details

¹ Departamento de Medicina Intensiva, Facultad de Medicina, Pontificia Universidad Católica de Chile, Santiago, Chile. ² Service de réanimation médicale, Hôpital Bicêtre, Hôpitaux Universitaires Paris-Sud, Paris, France. ³ Assistance Publique Hôpitaux de Paris, Université Paris-Sud, Paris, France. ⁴ Department of Intensive Care Adults, Erasmus MC University Medical Center, Rotterdam, The Netherlands. ⁵ Department of Pulmonary and Critical Care, New York University, New York, USA. ⁶ Division of Pulmonary, Allergy, and Critical Care Medicine, Columbia University Medical Center, New York, USA.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Published online: 23 January 2019

*Correspondence: glennguru@gmail.com

¹ Departamento de Medicina Intensiva, Facultad de Medicina, Pontificia Universidad Católica de Chile, Santiago, Chile
Full author information is available at the end of the article

The original article can be found online at <https://doi.org/10.1007/s00134-018-5499-8>.