



Correction to: Intraarticular application of autologous conditioned serum (ACS) reduces bone tunnel widening after ACL reconstructive surgery in a randomized controlled trial

Nikica Darabos¹ · Miroslav Haspl² · Carsten Moser³ · Anela Darabos⁴ · Dubravka Bartolek¹ · Dietrich Groenemeyer³

Published online: 22 October 2018

© European Society of Sports Traumatology, Knee Surgery, Arthroscopy (ESSKA) 2018

Correction to:

Knee Surg Sports Traumatol Arthrosc (2011) 19 (Suppl 1):

S36–S46

<https://doi.org/10.1007/s00167-011-1458-4>

The authors wish to declare the following competing interest, which should have been indicated in the research article.

- The Orthokine[®] product used in the trial was donated by the manufacturer.
- Co-author Carsten Moser was a medical manager and clinical director for the manufacturer of Orthokine between 2003 and 2008.

The original article can be found online at <https://doi.org/10.1007/s00167-011-1458-4>.

✉ Nikica Darabos
darabos.dr@vz.t-com.hr

Miroslav Haspl
miroslav.haspl@akromion.hr

Carsten Moser
moser@microtherapy.de

Anela Darabos
anela.darabos@vz.t-com.hr

Dubravka Bartolek
dubravka.bartolek1@zg.t-com.hr

Dietrich Groenemeyer
groenemeyer@microtherapy.de

¹ University Clinic for Traumatology, Medical School, University of Zagreb, Draskoviceva 19, 10000 Zagreb, Croatia

² Special Hospital for Orthopaedics and Traumatology Akromion, Ljudevita Gaja 2, 49 217 Krapinske Toplice, Croatia

³ Groenemeyer Institute for Microtherapy, University Witten-Herdecke, Universitaetsstrasse 142, 44799 Bochum, Germany

⁴ Scientific Unit, General Hospital Varazdin, Mestroviceva 1, 42000 Varazdin, Croatia