



# Correction to: Volumetric absorptive microsampling as an alternative tool for therapeutic drug monitoring of first-generation anti-epileptic drugs

Sofie Velghe<sup>1</sup> · Christophe P. Stove<sup>1</sup>


Published online: 15 February 2018  
© Springer-Verlag GmbH Germany, part of Springer Nature 2018

## Correction to: Anal Bioanal Chem

<https://doi.org/10.1007/s00216-018-0866-4>

We would like to call the reader's attention to the fact that unfortunately in Fig. 2 of the original article the figure headings of both graphs are the same. However, the heading of the upper graph should be 'IS-compensated recovery Low QC' and the one of the lower graph 'IS-compensated recovery High QC'. Please find below the corrected figure.

The original article has been corrected.



**Fig. 2** IS-compensated recovery (%) at Low and High QC level ( $n=6$ ) for VPA, PB, PHT, CBZ and CBZ-E measured in VAMS samples, prepared by pipetting 10  $\mu$ l blood at 4 different Hct levels (target values at 0.21, 0.42, 0.52 and 0.62). Shown are the mean and standard deviation. Asterisk denotes statistical difference ( $p<0.05$ , one-way ANOVA test) from the 0.42 Hct reference sample

The online version of the original article can be found at <https://doi.org/10.1007/s00216-018-0866-4>

✉ Christophe P. Stove  
christophe.stove@ugent.be

<sup>1</sup> Laboratory of Toxicology, Faculty of Pharmaceutical Sciences, Ghent University, Ottergemsesteenweg 460, 9000 Ghent, Belgium