



## Correction to: Two class I genes of the chicken MHC have different functions: BF1 is recognized by NK cells while BF2 is recognized by CTLs

Taejoong Kim<sup>1</sup> · Henry D. Hunt<sup>1</sup> · Mark S. Parcells<sup>2</sup> · Vicky van Santen<sup>3</sup> · Sandra J. Ewald<sup>4</sup>

Published online: 7 July 2018  
© Springer-Verlag GmbH Germany, part of Springer Nature 2018

### Correction to: Immunogenetics

<https://doi.org/10.1007/s00251-018-1066-2>

The Figure 3 in the original version of this article was incorrectly published. In this article the top panel of Figure 3 that describes the amino acid sequence alignment is now presented correctly.

The original article has been corrected.

---

The online version of the original article can be found at <https://doi.org/10.1007/s00251-018-1066-2>

---

✉ Taejoong Kim  
Taejoong.Kim@ars.usda.gov

✉ Henry D. Hunt  
hunthd@gmail.com

<sup>1</sup> United States Department of Agriculture, Agricultural Research Service, U.S. National Poultry Research Center, 934 College Station Rd., Athens, GA 30605, USA

<sup>2</sup> Department of Animal and Food Science, University of Delaware, Newark, DE 19717, USA

<sup>3</sup> Department of Pathobiology, Auburn University, 264 Green Hall, Auburn, AL 36849, USA

<sup>4</sup> Department of Poultry Science, Auburn University, 105 Poultry Annex Building, Auburn, AL 36849, USA

