



## Correction to: Biosynthesis and production of quercitols and their application in the production of pharmaceuticals: current status and prospects

Nobuya Itoh<sup>1</sup>

Published online: 21 September 2018

© Springer-Verlag GmbH Germany, part of Springer Nature 2018

**Correction to: Applied Microbiology and Biotechnology**  
<https://doi.org/10.1007/s00253-018-8972-y>

The published online version contains mistake in the chemical structure of *scyllo*-inosose in Fig. 5 and Fig. 7. The correct configuration of 1-hydroxyl group in *scyllo*-inosose should have been the same to *myo*-inositol. The configuration of 1-hydroxyl group of *scyllo*-inosose was opposite on the original publication.


---

The online version of the original article can be found at <https://doi.org/10.1007/s00253-018-8972-y>


---

✉ Nobuya Itoh  
nbito@pu-toyama.ac.jp; itohnobuya123@gmail.com

<sup>1</sup> Biotechnology Research Center and Department of Biotechnology, Toyama Prefectural University, 5180 Kurokawa, Imizu, Toyama 939-0398, Japan



**Fig. 5** Proposed biotransformation of *myo*-inositol to quercitols by *Salmonella typhimurium*



**Fig. 7** Catabolic pathway of *myo*-inositol in *Bacillus subtilis* for producing *scyllo*-inositol