



CORRECTION

## Correction to: Factors Associated with Unexpected Readmission Following Lung Resection

Shinsuke Uchida<sup>1</sup>  · Yukihiro Yoshida<sup>1</sup> · Masaya Yotsukura<sup>1</sup> · Kazuo Nakagawa<sup>1</sup> · Shun-ichi Watanabe<sup>1</sup>

Published online: 16 February 2021  
© Société Internationale de Chirurgie 2021

**Correction to:** *World J Surg*  
<https://doi.org/10.1007/s00268-020-05942-z>

In the original version of the online article some of the entries appeared in the wrong column. Following is the corrected table. The original article has also been corrected (Table 3).

---

The original article can be found online at <https://doi.org/10.1007/s00268-020-05942-z>.

---

✉ Shun-ichi Watanabe  
syuwatan@ncc.go.jp

<sup>1</sup> Department of Thoracic Surgery, National Cancer Center Hospital, 1-1 Tsukiji 5-chome, Chuo-ku, Tokyo 104-0045, Japan

**Table 3** Characteristics of patients with readmission or without readmission in detail

Variables	Patients without readmission, <i>n</i> (%)	Patients with readmission, <i>n</i> (%)	<i>P</i> value
Patient number	957	43	
Age, median (range)	69 (32–90)	70 (45–83)	0.306
Sex			
Male	505 (53)	33 (77)	0.002
Female	452 (47)	10 (23)	
Pack-year smoking, mean $\pm$ SD	26.3 $\pm$ 32.3	43.7 $\pm$ 41.9	0.001
Comorbidity			
Hypertension	341 (36)	17 (40)	0.602
Diabetes mellitus	160 (17)	13 (30)	0.022
Ischemic heart failure	34 (4)	4 (9)	0.075
IPF	19 (2)	4 (9)	0.014
Previous history of malignancy	230 (24)	15 (35)	0.106
%FVC, mean $\pm$ SD	102.7 $\pm$ 16.5	99.4 $\pm$ 15.6	0.200
FEV1/FVC (%), mean $\pm$ SD	74.4 $\pm$ 9.0	74.7 $\pm$ 8.1	0.838
Predicted FEV1 (%), mean $\pm$ SD	102.6 $\pm$ 19.9	102.7 $\pm$ 18.8	0.990
Maximum tumor size (mm), mean $\pm$ SD	26.7 $\pm$ 15.7	36.6 $\pm$ 21.9	<0.001
Serum CEA level, mean $\pm$ SD	6.2 $\pm$ 32.5	8.8 $\pm$ 17.6	0.605
Clinical nodal status			
N0	853 (89)	33 (77)	0.019
N1	68 (7)	6 (14)	
N2	36 (4)	4 (9)	
Primary site			
Right	585 (61)	27 (63)	0.827
Left	372 (39)	16 (37)	
Preoperative ipsilateral chest surgery or radiation therapy	34 (3.6)	2 (4.7)	0.465
Surgical procedure			
Pneumonectomy	13 (1)	1 (2)	0.462
Lobectomy	652 (68)	35 (81)	0.066
Segmentectomy	187 (20)	6 (6)	0.245
Wedge resection	105 (11)	1 (2)	0.072
Mediastinal lymph node dissection	590 (62)	29 (67)	0.444
Operative time (min), median (range)	119 (29–409)	128 (52–341)	0.004
Blood loss (ml), median (range)	30 (0–5292)	33 (2–650)	0.740
Transfusion	17 (2)	3 (7)	0.051
Postoperative complications			
Refractory air leakage	56 (5.8)	10 (23.2)	<0.001
Pneumonia	14 (1.1)	4 (9.3)	0.006
Chylothorax	8 (0.8)	0 (0)	0.703
AE of IPF during initial hospitalization	2 (0.2)	1 (2.3)	0.124
Empyema	4 (0.4)	1 (2.3)	0.198
Arrhythmia	9 (0.8)	2 (4.7)	0.078
Cerebral infarction	3 (0.3)	0 (0)	0.876
Reoperation	6 (0.6)	2 (4.7)	0.043
Pleurodesis	6 (0.6)	0 (0)	0.768
Postoperative hospitalization length (days), mean $\pm$ SD	5.0 $\pm$ 5.1	8.6 $\pm$ 10.6	<0.001

*SD* standard deviation, *FVC* forced vital capacity, *FEV* forced expiratory volume, *CEA* carcinoembryonic antigen, *AE* acute exacerbation, *IPF* idiopathic pulmonary fibrosis

**Publisher's Note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.