



Correction to: Immediate and delayed hypersensitivity after intra-arterial injection of iodinated contrast media: a prospective study in patients with coronary angiography

Kyoung-Hee Sohn^{1,2} · Gun-Woo Kim^{2,3} · Suh-Young Lee^{2,4,5} · Hyo-Soo Kim⁶ · Sang-Heon Cho^{2,4,5} · Jung-Kyu Han⁶ · Hye-Ryun Kang^{2,4,5}

Published online: 18 February 2020

© European Society of Radiology 2020

Correction to: European Radiology (2019) 29:5314–5321

<https://doi.org/10.1007/s00330-019-06138-3>

The original version of this article, published on 01 April 2019, unfortunately contained a mistake. The presentation of Fig. 1 was incorrect. The corrected figure is given below.

The online version of the original article can be found at <https://doi.org/10.1007/s00330-019-06138-3>

✉ Jung-Kyu Han
hperates@gmail.com

✉ Hye-Ryun Kang
helenmed@snu.ac.kr

¹ Department of Internal Medicine, Kyung Hee University Medical Center, Seoul, South Korea

² Institute of Allergy and Clinical Immunology, Seoul National University Medical Research Center, Seoul, South Korea

³ Department of Internal Medicine, St. Carollo General Hospital, Suncheon, South Korea

⁴ Division of Allergy and Clinical Immunology, Department of Internal Medicine, Seoul National University College of Medicine, 101 Daehak-ro, Chongno-gu, Seoul 110-744, South Korea

⁵ Seoul National University Hospital Regional Pharmacovigilance Center, Seoul, South Korea

⁶ Department of Internal Medicine and Cardiovascular Center, Seoul National University Hospital, 101 Daehak-ro, Chongno-gu, Seoul 110-744, South Korea

Fig. 1 a Rate of hypersensitivity reactions according to previous exposure to intravenous and intra-arterial contrast agents: Previous exposure to ICM via intra-arterial route increased the incidence of immediate HSR. **b** Pie chart showing severity of immediate and delayed hypersensitivity reactions in immediate and delayed HSR. *HSR* hypersensitivity reactions, *ICM* iodinated contrast media

