

Z Gerontol Geriat 2016 · 49:167
DOI 10.1007/s00391-015-1015-8
Published online: 15 January 2016
© Springer-Verlag Berlin Heidelberg 2015



Maria Magdalena Schreier¹ · Ulrike Bauer¹ · Jürgen Osterbrink¹ · Josef Niebauer² · Bernhard Iglseider³ · Jens Reiss³

¹ Institute of Nursing Science and Practice, Paracelsus Medical University, Salzburg, Austria

² Institute of preventive and rehabilitative Sports Medicine, Landeskrankenhaus, Salzburg, Austria

³ Department of Geriatric Medicine, Christian Doppler Klinik, Paracelsus Medical University, Salzburg, Austria

Erratum to: Fitness training for the old and frail

Effectiveness and impact on daily life coping and self-care abilities

Erratum to:

Z Gerontol Geriat (2015)

DOI 10.1007/s00391-015-0966-0

Methods

Training program: collection and analysis of quantitative data

Medical diagnoses, laboratory parameters and data on balance, risk of falls, etc. were collected from medical records. The statistical procedures included descriptive, univariate and bivariate analyses. Significance testing comparing various time points (t0 = baseline, t4 = end of training and t5 = follow-up) for the intervention and control groups were performed using IBM SPSS Statistics 18.

The following information is incorrect: Cronbach's alpha was set a priori at 5%.

The correct statement is: Alpha was set a priori at 5%.

The authors would like to apologise for this error.

Corresponding address

M.M. Schreier

Institute of Nursing Science and Practice,
Paracelsus Medical University
Salzburg
magdalena.schreier@pmu.ac.at

The online version of the original article can be found under doi:10.1007/s00391-015-0966-0