




Correction to: Force–velocity profiling of sprinting athletes: single-run vs. multiple-run methods

Christian Helland¹ · Thomas Haugen¹ · Elvir Rakovic² · Ola Eriksrud³ · Olivier Seynnes³ · Antti A. Mero⁴ · Gøran Paulsen¹ 

Published online: 5 January 2019
© Springer-Verlag GmbH Germany, part of Springer Nature 2019

Correction to: European Journal of Applied Physiology
<https://doi.org/10.1007/s00421-018-4045-2>

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

The author would like to correct the errors in the publication of the original article. The corrected details are given below for your reading.

In the abstract, the fourth sentence of the result section should read as:

No substantial correlation between the changes (from pre- to post-training tests) in S_{FV} calculated with the single-run and the multiple-run methods were observed ($r=0.03$).

The original article can be found online at <https://doi.org/10.1007/s00421-018-4045-2>.

✉ Gøran Paulsen
goran.paulsen@olympiatoppen.no

¹ Norwegian Olympic and Paralympic Committee and Confederation of Sports, Oslo, Norway

² University of Gothenburg, Gothenburg, Sweden

³ Norwegian School of Sport Sciences, Oslo, Norway

⁴ Biology of Physical Activity, University of Jyväskylä, Jyväskylä, Finland