



Correction to: Two-hour post-challenge glucose is a better predictor of adverse outcome after myocardial infarction than fasting or admission glucose in patients without diabetes

Sudipta Chattopadhyay¹ · Anish George² · Joseph John³ · Thozhukat Sathyapalan⁴

Published online: 5 March 2018
© Springer-Verlag Italia S.r.l., part of Springer Nature 2018

Correction to: Acta Diabetologica
<https://doi.org/10.1007/s00592-018-1114-2>

Unfortunately, name of co-author “Thozhukat” was misspelled in the original publication and the same is corrected here. The original article has been corrected.

The original article can be found online at <https://doi.org/10.1007/s00592-018-1114-2>.

✉ Sudipta Chattopadhyay
Sudipta.Chattopadhyay@nhs.net

Anish George
anish26g@yahoo.com

Joseph John
Joseph.John@nhs.net

Thozhukat Sathyapalan
Thozhukat.Sathyapalan@hyms.ac.uk

- ¹ Department of Cardiology, Milton Keynes University Hospital, Standing Way, Milton Keynes MK6 5LD, UK
- ² Department of Cardiology, Scunthorpe General Hospital, Cliff Gardens, Scunthorpe, UK
- ³ Department of Cardiology, Castle Hill Hospital, Kingston upon Hull, UK
- ⁴ Department of Academic Endocrinology, Diabetes and Metabolism, Hull York Medical School, University of Hull, Kingston upon Hull, UK