




Correction to: Association of ischemic stroke with ankylosing spondylitis: a nationwide longitudinal cohort study

Dong Hyun Lee¹ · Yoon Jin Choi² · In-bo Han¹ · Je Beom Hong¹ · Kyung Do Han³ · Jung Min Choi⁴ · Seil Sohn¹ 

Published online: 24 March 2018
© Springer-Verlag GmbH Austria, part of Springer Nature 2018

Correction to: Acta Neurochirurgica

<https://doi.org/10.1007/s00701-018-3499-7>

The authors regret to state that an incorrect image was uploaded in Fig. 2. Please see a newly updated Fig. 2. All statements including Figure Legends are correct.

The online version of the original article can be found at <https://doi.org/10.1007/s00701-018-3499-7>

✉ Seil Sohn
sisohn@cha.ac.kr

¹ Department of Neurosurgery, CHA Bundang Medical Center, CHA University College of Medicine, 59, Yatap-ro, Bundang-gu, Seongnam-si, Gyeonggi-do 13496, South Korea

² Department of Internal Medicine, Seoul National University Bundang Hospital, Gyeonggi-do, Seongnam, South Korea

³ Department of Biostatistics, College of Medicine, Catholic University of Korea, Seoul, South Korea

⁴ Department of Medical Device Management and Research, SAIHST, Sungkyunkwan University, Seoul, South Korea

Fig. 2 Comparison of the cumulative incidence of ischemic stroke in the AS and control groups. The Kaplan-Meier curves with cumulative hazards of ischemic stroke were compared in the AS and control groups. The incidence of ischemic stroke in the AS group was significantly higher than that in the control group ($p = 0.0041$)

