

Assessing groundwater contamination risk using the DASTI/IDRISI GIS method: coastal system of western Mamora, Morocco

Brahim Ben Kabbour · Lahcen Zouhri ·
Jacky Mania · Jean-Pierre Colbeaux

Published online: 29 August 2007
© Springer-Verlag 2007

Erratum to: Bull Eng Geol Environ (2006) 65:463–470
DOI 10.1007/s10064-004-0265-2

This article contains two errors in Figs. 7 and 8. The correct figures are given below:




Fig. 7 Vulnerability map according to the DRASTIC method [I. Vulnerability ranges (Aller et al. 1987), II. intervals of Id values]

The online version of the original article can be found under doi:[10.1007/s10064-004-0265-2](https://doi.org/10.1007/s10064-004-0265-2).

B. B. Kabbour
FST Beni Mellal, Département de Géologie,
Laboratoire d'Aménagement du Territoire,
Université Cadi Ayyad, Marrakech, Morocco

L. Zouhri (✉)
UMR CNRS 8110, Processus et Bilans des Domaines
Sédimentaires, UFR des Sciences de la Terre,
Université des Sciences et Technologies de Lille (USTL),
59655 Villeneuve d'Ascq Cedex, France
e-mail: lahcen.zouhri@laposte.net

J. Mania
UMR CNRS 8107, LML, Ecole Polytechnique de Lille,
Avenue Paul Langevin, 59655 Villeneuve d'Ascq Cedex, France

J.-P. Colbeaux
UFR de Géologie, Bât SN5, Université de Lille I,
59655 Villeneuve d'Ascq Cedex, France




Fig. 8 Sensibility map according to the DRASTIC method and land use analysis