## **CORRECTION**



## Correction to: Role of low-level laser therapy added to facial expression exercises in patients with idiopathic facial (Bell's) palsy

Banu Ordahan 1 · Ali yavuz Karahan 1

Published online: 17 June 2019 © Springer-Verlag London Ltd., part of Springer Nature 2019

Correction to: Lasers in Medical Science https://doi.org/10.1007/s10103-017-2195-9

The online version of the original article can be found at https://doi.org/ 10.1007/s10103-017-2195-9



Department of Physical Medicine and Rehabilitation, Konya Education and Training Hospital, Yazırmah, Selçuklu, Konya, Turkey

The published version contains a mistake in Figure 1. Below is the correct figure.

Fig. 1 Flow diagram Enrollment Assessed for eligibility (n = 48) Excluded (n =2) • The history of recurrent Bell's Palsy Randomized: (n = 46)Received LLLT (830 nm and 100 Mw, 1 Received facial exercises intervention alone KHz) with facial exercise treatment Allocation (n = 23) (n = 23)Follow-up parameters 3<sup>th</sup> week follow-up 3<sup>th</sup> week follow-up PFDI No drop-out No drop-out SFDI (n = 23)(n = 23)Follow-up 6<sup>th</sup> week follow-up 6<sup>th</sup> week follow-up Follow-up parameters PFDI No drop-out No drop-out SFDI (n = 23)(n = 23)Analysis Analyzed Analyzed (n = 23)(n = 23)

LLLT: Low-level laser treatment, PFDI: Facial disability index physical; SFDI: Facial disability index social

**Publisher's note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

