## **CORRECTION**





## Correction to: Contribution of thrips to seed production in *Habenaria* radiata, an orchid morphologically adapted to hawkmoths

Koji Shigeta<sup>1</sup> · Kenji Suetsugu<sup>2</sup>

Published online: 6 June 2020 © The Botanical Society of Japan 2020

Correction to: Journal of Plant Research https://doi.org/10.1007/s10265-020-01205-z

In the original publication of the article, Fig. 2c was erroneously identical with Fig. 2a and correct version is provided below.

The original article can be found online at https://doi.org/10.1007/s10265-020-01205-z.

- ⊠ Koji Shigeta koji.shiget@gmail.com
- ⊠ Kenji Suetsugu kenji.suetsugu@gmail.com
- Food Technology Research Center, Hiroshima Prefectural Technology Research Institute, 12-70 Hijiyama-honmachi, Minami-ku, Hiroshima 732-0816, Japan
- Department of Biology, Graduate School of Science, Kobe University, 1-1 Rokkodai, Nada-ku, Kobe 657-8501, Japan






Fig. 2 Histograms of the seed set of *Habenaria radiata* from manually cross-pollinated flowers (a), open-pollinated flowers (b), and from flowers bagged with mesh bags (c)

**Publisher's Note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

