

## Erratum to: Peroxisome proliferator-activated receptor gamma activates fas ligand gene promoter inducing apoptosis in human breast cancer cells

Daniela Bonofiglio<sup>1</sup> · Sabrina Gabriele<sup>1</sup> · Saveria Aquila<sup>1,2</sup> · Hongyan Qi<sup>1</sup> ·  
Maria Belmonte<sup>1</sup> · Stefania Catalano<sup>1,2</sup> · Sebastiano Andò<sup>2,3,4</sup>

Published online: 18 August 2017  
© Springer Science+Business Media, LLC 2017

**Erratum to: Breast Cancer Res Treat (2009)**  
**113:423–434**  
**DOI 10.1007/s10549-008-9944-1**

and the new Fig. 5a–c were provided in this erratum. The authors apologize for the inconvenience caused.

Unfortunately, the authors could not find the blots of the original figures of 5a–c. As suggested by the Editor, the authors have repeated the experiments relative to Fig. 5a–c

---

The online version of the original article can be found under  
doi:[10.1007/s10549-008-9944-1](https://doi.org/10.1007/s10549-008-9944-1).

---

✉ Sebastiano Andò  
sebastiano.ando@unical.it

- <sup>1</sup> Department of Pharmaco-Biology, University of Calabria, Arcavacata di Rende, Cosenza 87036, Italy
- <sup>2</sup> Centro Sanitario, University of Calabria, Arcavacata di Rende, Cosenza 87036, Italy
- <sup>3</sup> Faculty of Pharmacy, University of Calabria, Arcavacata di Rende, Cosenza 87036, Italy
- <sup>4</sup> Department of Cellular Biology, University of Calabria, Arcavacata di Rende, Cosenza 87036, Italy

**Fig. 5** BRL induces the extrinsic apoptotic pathway in MCF7 cells. PPAR $\gamma$  (a) and FasL (b) protein expression (evaluated by WB) in MCF7 cells transfected with a 25-nucleotide of RNA interference (RNAi) targeted human PPAR $\gamma$  or FasL mRNA sequence respectively, or with control RNAi as reported in Materials and Methods or not transfected and treated for 48 h as indicated.  $\beta$ -actin was used as loading control. **c** MCF7 cells were treated for 48 h as indicated, or transfected with PPAR $\gamma$ , FasL or control RNAis. Positions of procaspase 8 (P) and its active cleavage product (S) are indicated by arrowheads on the right. One of three similar experiments is presented.  $\beta$ -actin was used as loading control on the same stripped blot. **d** DNA laddering was performed in MCF7 cells treated for 72-h as indicated, or transfected with PPAR $\gamma$  or control RNAis

