

Erratum to: CXCL12 / CXCR4 / CXCR7 chemokine axis and cancer progression

Xueqing Sun · Guangcun Cheng · Mingang Hao ·
Jianghua Zheng · Xiaoming Zhou · Jian Zhang ·
Russell S. Taichman · Kenneth J. Pienta · Jianhua Wang

Published online: 12 November 2010
© Springer Science+Business Media, LLC 2010

Erratum to: Cancer Metastasis Rev (2010) 29:709–722
DOI 10.1007/s10555-010-9256-x

The original version of this article unfortunately contained a mistake. An error has been found in Figure 1. The correct version is given below

The online version of the original article can be found at
<http://dx.doi.org/10.1007/s10555-010-9256-x>.

X. Sun · G. Cheng · M. Hao · J. Zheng · X. Zhou · J. Wang (✉)
Department of Biochemistry and Molecular & Cell Biology,
Key Laboratory of Cell Differentiation and Apoptosis of Chinese,
Ministry of Education, Institute of Medical Science,
Shanghai Jiao Tong University School of Medicine,
Shanghai 200025, People's Republic of China
e-mail: jianhuaw2007@gmail.com

J. Zhang · K. J. Pienta
Department of Internal Medicine and Urology,
University of Michigan,
Ann Arbor, MI 48109, USA

R. S. Taichman
Department of Periodontics & Oral Medicine,
University of Michigan School of Dentistry,
Ann Arbor, MI 48109, USA




Fig. 1 Chemokine family and their cognate receptors. Most chemokines can bind multiple receptors, and a single receptor can bind multiple chemokines, as shown in this case for most CC (blue) and

CXC (green) chemokines. Decoy receptors (black) can also interact with multiple chemokines. By contrast, a minority of receptors (red) have only one ligand