

N-glycan trimming by glucosidase II is essential for *Arabidopsis* development

Pravina Soussilane · Cecilia D'Alessio ·
Thomas Paccalet · Anne-Catherine Fitchette ·
Armando J. Parodi · Richard Williamson ·
Carole Plasson · Loïc Faye · Véronique Gomord

Published online: 13 January 2009
© Springer Science + Business Media, LLC 2009

Erratum to: Glycoconj J
DOI 10.1007/s10719-008-9201-1

The original version of this article unfortunately contained a mistake. The name of the first author was not presented correctly. “Pravina Soussilane” should be “Pravina Soussilane.”

The online version of the original article can be found at <http://dx.doi.org/10.1007/s10719-008-9201-1> of the article with published errors.

P. Soussilane · T. Paccalet · A.-C. Fitchette · C. Plasson · L. Faye ·
V. Gomord (✉)
CNRS, UMR 6037, IFRMP 23, Bâtiment Biologie Extension,
Faculté des Sciences,
Mont-Saint-Aignan, France
e-mail: vgomord@crihan.fr

C. D'Alessio · A. J. Parodi
Instituto Leloir,
Av Patricias Argentinas 435,
Buenos Aires, Argentina

R. Williamson
Plant Cell Biology Group,
Research School of Biological Sciences,
The Australian National University,
G.P.O. Box 475, Canberra ACT 2601, Australia