

## Erratum to: Comparison of the prevalence of convulsions associated with the use of cefepime and meropenem

Akihiro Tanaka · Kenshi Takechi · Shinichi Watanabe ·  
Mamoru Tanaka · Katsuya Suemaru ·  
Hiroaki Araki

Published online: 7 February 2015  
© Koninklijke Nederlandse Maatschappij ter bevordering der Pharmacie 2015

**Erratum to: Int J Clin Pharm (2013) 35:683–687**  
**DOI 10.1007/s11096-013-9799-3**

In the original publication, an incorrect Table 2 (Table 1 duplicated) has been published inadvertently. The correct Table 2 is given in this erratum.

---

The online version of the original article can be found under  
doi:[10.1007/s11096-013-9799-3](https://doi.org/10.1007/s11096-013-9799-3).

---

A. Tanaka (✉) · K. Takechi · S. Watanabe · M. Tanaka ·  
H. Araki  
Division of Pharmacy, Ehime University Hospital,  
454 Shitsukawa, Toon, Ehime 791-0295, Japan  
e-mail: [akiki@m.ehime-u.ac.jp](mailto:akiki@m.ehime-u.ac.jp)

K. Suemaru  
School of Pharmacy, Shujitu University, 1-6-1 Nishigawara,  
Okayama 703-8516, Japan

**Table 2** Clinical characteristics of patients with meropenem-associated convulsions

No.	Patient characteristics (sex, age, eGFR)	Daily dosing schedule	Time of symptoms (days from start of cefepime)	Symptoms and signs	Baseline disease	Anticonvulsant agent	Final assessment
A	F, 42, 95 mL/min	0.5 g every 8 h	2	Clonic convulsion	Cerebral infarction	Phenytoin (iv), valproic acid (po)	Administer thiamylal, and then complete resolution
B	M, 59, 106 mL/min	0.5 g every 8 h	2	Clonic convulsion	Encephaloma	None	Administer diazepam intravenously, and then complete resolution
C	F, 74, 69 mL/min	1 g every 12 h	2	Myoclonus	Multiple cerebral infarction	Clonazepam (po)	Administer diazepam intravenously, but uncontrollable convulsion, and then complete resolution
D	F, 39, 76 mL/min	1 g every 12 h	2	Clonic convulsion and decreased level of consciousness	B- cell lymphoma (infiltration brain), febrile neutropenia	Zonisamide (po)	Administer diazepam intravenously, and then complete resolution after the discontinuation of meropenem