

Erratum to: Acromegaly: surgical results in 548 patients

Cecilia Fernández Mateos¹ · Maria García-Uria² · Tomás Lucas Morante³ · José García-Uria¹

Published online: 5 July 2017
© Springer Science+Business Media, LLC 2017

Erratum to: Pituitary DOI 10.1007/s11102-017-0813-y

The original version of this article unfortunately contained a mistake in text and table header. The corrections are given below. The original article has been corrected.

In Preoperative hormonal values section, the PRL must be >100 ng/ml, not >00 ng/ml as published. The correct paragraph is given below:

PRL evaluation was realized in 453 patients (82.6%). Hyperprolactinemia >5 ng/ml was associated with acromegaly in 95 patients (20.9%), (64 women and 31 men.). PRL >100 ng/ml was found in twelve patients (2.64%), (7 women and 5 men).

The column headers of Table 3 are not in order. The corrected Table 3 is given below:

The online version of the original article can be found under doi:[10.1007/s11102-017-0813-y](https://doi.org/10.1007/s11102-017-0813-y).

✉ Cecilia Fernández Mateos
valdelagua@hotmail.com

Maria García-Uria
mariagarciauria@hotmail.com

Tomás Lucas Morante
tlucasmorante@gmail.com

José García-Uria
neuromic@telefonica.net

- ¹ Neurosurgery Department, Puerta de Hierro Hospital, Madrid, Spain
- ² Emergency Medicine Department, Puerta de Hierro Hospital, Madrid, Spain
- ³ Endocrinology Department, Puerta de Hierro Hospital, Madrid, Spain

Table 3 Postoperative GH-levels (548 patients)

Period	GH >5 ng/ml	GH 5–2.5 ng/ml	GH 2.4–2 ng/ml	GH <2 ng/ml
1975–84	40 (51%)	7 (9%)	2 (3%)	29 (37%)
1985–94	39 (25%)	9 (6%)	6 (4%)	101 (65%)
1995–04	35 (17%)	11 (5.4%)	11 (5.4%)	148 (72.2%)
2005–15	19 (17%)	7 (6%)	6 (5%)	78 (70%)
Total patients	133 (24.2%)	34 (6.2%)	25 (4.5%)	356 (65%)

*Surgical results (1975-2015) Postoperative GH 2-21 days after surgery