



# Correction to: When Marijuana Is Used before Cigarettes or Alcohol: Demographic Predictors and Associations with Heavy Use, Cannabis Use Disorder, and Other Drug-related Outcomes

Brian J. Fairman<sup>1</sup> · C. Debra Furr-Holden<sup>2</sup> · Renee M. Johnson<sup>3</sup>

Published online: 31 May 2018  
© Society for Prevention Research 2018

**Correction to: Prevention Science (2018)**  
<https://doi.org/10.1007/s11121-018-0908-3>

The original version of this article contained a mistake: The first sentence in the Results section of the Abstract should read as follows: “The proportion who had initiated marijuana before other substances increased from 4.4% to 8.0%”.

The authors regret the oversight.

---

The online version of the original article can be found at <https://doi.org/10.1007/s11121-018-0908-3>

---

✉ Brian J. Fairman  
brian.fairman@nih.gov

- <sup>1</sup> Health Behavior Branch, Division of Intramural Population Health Research, Eunice Kennedy Shriver National Institute of Child Health and Human Development, 6710B Rockledge Dr., Bethesda, MD 20817, USA
- <sup>2</sup> Department of Epidemiology and Biostatistics, Michigan State University, East Lansing, MI, USA
- <sup>3</sup> Department of Mental Health, John Hopkins Bloomberg School of Public Health, Baltimore, MD, USA