


# Magnetic Field Amplification in Galaxy Clusters and Its Simulation

J. Donnert<sup>1,2</sup>  · F. Vazza<sup>3</sup> · M. Brüggen<sup>4</sup> · J. ZuHone<sup>5</sup>

Received: 3 July 2018 / Accepted: 30 October 2018 / Published online: 13 November 2018  
© The Author(s) 2018

**Abstract** We review the present theoretical and numerical understanding of magnetic field amplification in cosmic large-scale structure, on length scales of galaxy clusters and beyond. Structure formation drives compression and turbulence, which amplify tiny magnetic seed fields to the microGauss values that are observed in the intracluster medium. This process is intimately connected to the properties of turbulence and the microphysics of the intra-cluster medium. Additional roles are played by merger induced shocks that sweep through the intra-cluster medium and motions induced by sloshing cool cores. The accurate simulation of magnetic field amplification in clusters still poses a serious challenge for simulations of cosmological structure formation. We review the current literature on cosmological simulations that include magnetic fields and outline theoretical as well as numerical challenges.

**Keywords** Galaxy clusters · Magnetic fields · Simulations · Magnetic dynamo

---

Clusters of Galaxies: Physics and Cosmology

Edited by Andrei Bykov, Jelle Kaastra, Marcus Brüggen, Maxim Markevitch, Maurizio Falanga and Frederik Bernard Stefan Paerels

---

✉ J. Donnert  
[donnert@ira.inaf.it](mailto:donnert@ira.inaf.it); [jdonnert@strw.leidenuniv.nl](mailto:jdonnert@strw.leidenuniv.nl)

F. Vazza  
[franco.vazza2@unibo.it](mailto:franco.vazza2@unibo.it)

M. Brüggen  
[mbrueggen@hs.uni-hamburg.de](mailto:mbrueggen@hs.uni-hamburg.de)

J. ZuHone  
[john.zuhone@cfa.harvard.edu](mailto:john.zuhone@cfa.harvard.edu)

<sup>1</sup> via P. Gobetti 101, 40129 Bologna, Italy

<sup>2</sup> Present address: Leiden Observatory, PO Box 9513, 2300 RA Leiden, The Netherlands

<sup>3</sup> Dipartimento di Fisica e Astronomia, via Gobetti 93/2, 40129 Bologna, Italy

<sup>4</sup> Hamburg Observatory, Gojenbergsweg 112, 21029 Hamburg, Germany

<sup>5</sup> Smithsonian Astrophysical Observatory, 60 Garden St., Cambridge, MA 02138, USA

## 1 Introduction

Magnetic fields permeate our Universe, which is filled with ionized gas from the scales of our solar system up to filaments and voids in the large-scale structure (Klein and Fletcher 2015). While magnetic fields are usually not dynamically important, their presence shapes the physical properties of the Baryonic medium (Schekochihin and Cowley 2007). On the largest scales, radio observations remain our most important tool to estimate magnetic fields today (see e.g. van Weeren, this volume). Recent and upcoming advances in instrumentation enable the observation of radio emission on scales of a few kpc at the cluster outskirts and will soon provide three-dimensional magnetic field distributions in the inter-cluster-medium (ICM) through Faraday tomography (Govoni et al. 2014).

Connecting these new observations to theoretical expectations is a major challenge for the community, due to the complexity of the non-thermal physics in the cosmological context. In the framework of cold Dark Matter, structure formation is dominated by gravitational forces and proceeds from the bottom up: smaller DM halos form first (Planelles et al. 2016), and baryons flow into the resulting potential well. Through cooling, stars and galaxies form and evolve into larger structures (groups, clusters, filaments), by infall and merging (Mo et al. 2010). These processes drive turbulent gas motions and a magnetic dynamo that amplifies some form of seed field to  $\mu\text{G}$  values in the center of galaxy clusters. Galaxy feedback injects magnetic fields and relativistic particles (cosmic-ray protons and electrons) into the large-scale structure that interact with shocks and turbulence, get (re-)accelerated and finally become observable at radio frequencies and potentially in the  $\gamma$ -ray regime (Schlickeiser 2002; Lazarian et al. 2012; Brunetti and Jones 2014).

In the past decade significant progress has been made in the simulation of galaxy formation, with an emphasis on physical models for feedback (e.g. Naab and Ostriker 2017). Unfortunately, the same is not true for the simulation of turbulence, magnetic fields and cosmic-ray evolution—nearly every step in the chain of non-thermal processes remains open today:

What is the origin of the magnetic seed fields and the contributions of various astrophysical sources? What are the properties of turbulence and the magnetic dynamo in the ICM, filaments, and voids? What is the distribution and topology of magnetic fields? What is the spatial distribution of radio dark cosmic-ray electrons in clusters? Where are the cosmic-ray protons? What are their sources? What physics governs particle acceleration in shocks that leads to radio relics? How does turbulence couple to cosmic-rays in radio halos? What are the physical properties (viscosity, resistivity, effective collisional scales) of the diffuse plasma in the ICM, filaments and voids?

Answers have proven themselves difficult to obtain, in part because turbulence is a demanding numerical problem, but also because the physics is different enough from galaxy formation to make some powerful numerical approaches like density adaptivity rather ineffective. Today, JVLA and LOFAR observations have achieved an unprecedented spatial and spectral detail in the observation of magnetic phenomena in cluster outskirts (e.g. Owen et al. 2014; Hoang et al. 2017; Rajpurohit et al. 2018), thereby challenging simulations to increase their level of spatial and physical detail. The gap will likely widen in the next years as SKA precursors like ASKAP see first light (Gaensler et al. 2010) and results from the LOFAR survey key science project become available (Shimwell et al. 2017).

Here we review the current status on astrophysical and cosmological simulations of magnetic field amplification in structure formation through compression, shocks, turbulence and cosmic-rays. Such a review will naturally emphasize galaxy clusters, simply because there is only weak observational evidence for magnetic fields in filaments and voids. We will

also touch on ideal MHD as a model for intergalactic plasmas and introduce fundamental concepts of turbulence and the MHD dynamo. We are putting an emphasis on numerical simulations because they are our most powerful tool to study the interplay of non-thermal physics. This must also include some details on common algorithms for MHD and their limitations. Today, these algorithms and their implementation limit our ability to model shocks, turbulence and the MHD dynamo in a cosmological framework.

We exclude from this review topics that are not directly related to simulations of the cosmic magnetic dynamo. While we shortly introduce turbulence and dynamo theory, we do not attempt to go into detail, several reviews are available (e.g. Schekochihin and Cowley 2007, for an introduction). We also do not review models for particle acceleration in clusters (Brunetti and Jones 2014) or observations (see Ferrari et al. 2008, and van Weeren et al., this volume). We also do not discuss in detail the seeding of magnetic fields (see Widrow et al. 2012; Ryu et al. 2012; Subramanian 2016, for recent exhaustive reviews on the topic), nor the amplification of magnetic fields in the interstellar medium (see Federrath 2016, for a recent review) or in galaxies (e.g. Schleicher et al. 2010; Beck et al. 2012; Martin-Alvarez et al. 2018, for theoretical reviews).

## 1.1 Overview

Galaxy clusters form through the gravitational collapse and subsequent merging of virialized structures into haloes, containing about 80% Dark Matter and 20% Baryons (Sarazin 2002; Voit 2005; Planelles et al. 2015). From X-ray observations we know that the diffuse thermal gas in the center of haloes with masses  $> 10^{14} M_{\odot}$  is completely ionized, with temperatures of  $T = 10^8$  K and number densities of  $n_{\text{th}} \approx 10^{-3} \text{ cm}^{-3}$ , (e.g. Sarazin 1988; Borgani et al. 2008). The speed of sound is then  $c_s = \sqrt{\gamma P/\rho} \approx 1200 \text{ km/s}$ , where  $\gamma = 5/3$  is the adiabatic index at density  $\rho$  and pressure  $P$ .

The ideal equation of state for a monoatomic gas is applicable in such a hot under-dense medium, even though the ICM contains  $\approx 25\%$  helium and heavier elements as well (e.g. Böhringer and Werner 2010). In fact, the intracluster medium is one of the most ideal plasmas known, with a plasma parameter of  $g \approx 10^{-15}$  and a Debye length of  $\lambda_D \approx 10^5 \text{ cm}$  that still contains  $\approx 10^{12}$  protons and electrons. In contrast, the mean free path for Coulomb collisions is in the kpc regime (Eq. (6)). Clearly, electromagnetic particle interactions dominate over two-body Coulomb collisions and plasma waves shape the properties of the medium on small scales (e.g. Schlickeiser 2002, Table 8.1).

Cluster magnetic fields of  $1 \mu\text{G}$  were first estimated from upper limits on the diffuse synchrotron emission of intergalactic material in a  $1 \text{ Mpc}^3$  volume by Burbidge (1958). With the discovery of the Coma radio halo by Willson (1970), this was confirmed using equipartition arguments between the cosmic-ray electron energy density and magnetic energy density (e.g. Beck and Krause 2005). Later estimates based on the rotation measure of background sources to the Coma cluster obtain central magnetic fields of 3–7  $\mu\text{G}$  scaling with ICM thermal density with an exponent of 0.5–1 (e.g. Bonafede et al. 2010). Hence the ICM is a high  $\beta = n_{\text{th}} k_B T / B^2 \approx 100$  plasma, where thermal pressure dominates magnetic pressure.

Based on above estimates, one may hope that magneto-hydrodynamics (MHD) is applicable on large enough scales in clusters (Sects. 3.2.1 and 3.2.2). Then the magnetic field  $\mathbf{B}$  evolves with the flow velocity  $\mathbf{v}$  according to the induction equation (Landau et al. 1961):

$$\frac{\partial \mathbf{B}}{\partial t} = -\mathbf{v} \cdot \nabla \mathbf{B} + \mathbf{B} \cdot \nabla \mathbf{v} - \mathbf{B} \nabla \cdot \mathbf{v} + \eta \Delta \mathbf{B}, \quad (1)$$

where the first term accounts for the advection of field lines, the second one for stretching, the third term for compression and the fourth term for the magnetic field dissipation with the

diffusivity  $\eta = c_s/4\pi\sigma$  and the conductivity  $\sigma$ . Because the ICM is a nearly perfect plasma ( $\beta_{\text{pl}} \gg 1$ ), conductivity is very high, diffusivity likely very low ( $\eta \approx 0$ ). Then the induction equation (1) predicts that magnetic fields are frozen into the plasma and advected with the bulk motions of the medium (Kulsrud and Ostriker 2006). Because Eq. (1) is a conservation equation for magnetic flux, magnetic fields cannot be created in the MHD framework, but have to be seeded by some mechanism, also at high redshift (Sect. 2). However, current upper limits on large-scale magnetic fields exclude large-scale seed fields above  $\sim$  nG (Ade et al. 2016), and back-of-the-envelope calculations show that pure compression cannot produce  $\mu\text{G}$  in clusters from such initial values (Sect. 3.1). X-ray observations have revealed substantial turbulent velocities in a few clusters (Schuecker et al. 2004; Zhuravleva et al. 2014; Aharonian et al. 2016). These are in agreement with estimates from rotation measurements (Vogt and Enßlin 2003; Kuchar and Enßlin 2011) that can also be used to constrain magnetic field power spectra (Vacca et al. 2012, 2016; Govoni et al. 2017).

It is reasonable to assume some form of turbulent dynamo in the clusters and possibly filaments (Jaffe 1980; Roland 1981; Ruzmaikin et al. 1989; De Young 1992; Goldshmidt and Rephaeli 1993; Kulsrud et al. 1997; Sánchez-Salcedo et al. 1998; Subramanian et al. 2006; Enßlin and Vogt 2006), but it is necessary to consider plasma-physical arguments to understand the fast growth of seed fields by many orders of magnitude (Schekochihin et al. 2005b; Schekochihin and Cowley 2007). There are clear theoretical predictions for idealized MHD dynamos (e.g. Schekochihin et al. 2004; Porter et al. 2015), which show that magnetic fields are amplified through an inverse cascade at the growing *Alfvén scale*, where the field starts back-reacting on the flow. This is called the *small-scale dynamo* (Sect. 3.3). However, the astrophysical situation differs significantly from these idealized models: structure formation drives turbulence localized, episodic and multi-scale in the presence of a strong gravitational potential in galaxy clusters (Sect. 3.5), and the magneto-hydrodynamical properties of the medium are far from clear (Schekochihin et al. 2009). Shocks and cosmic-rays amplify magnetic fields as well and are very difficult to model (Sect. 5).

With JVLA, LOFAR, ASKAP and the SKA, the Alfvén scale comes within the range of radio observations: radio relics are now spatially resolved to a few kpc in polarization; low-frequency surveys are expected to find hundreds of radio halos and mini-halos; Faraday tomography will allow to map magnetic field structure also along the line of sight (see van Weeren et al., this volume). Future X-ray missions will put stringent bounds on turbulent velocities in clusters and constrain magnetic field amplification by draping and sloshing in cold fronts (Sect. 6).

## 2 Magnetic Seeding Processes

Let us begin with a short overview of proposed seeding mechanisms; a detailed review can be found e.g. in Subramanian (2016). It is very likely that more than one of these mechanisms contributes to the magnetization of the large-scale structure. Hence an important question for simulations of magnetic field amplification is the influence of these seeding mechanisms on the final magnetic field.

### 2.1 Primordial Mechanisms

Several mechanisms for the initial seed field have been suggested to start the dynamo amplification process within galaxies and galaxy clusters. Some of the proposed scenarios involve the generation of currents during inflation, phase transitions and baryogenesis (e.g. Harrison



1973; Kahniashvili et al. 2010, 2011, 2016; Widrow et al. 2012; Durrer and Neronov 2013; Subramanian 2016). These primordial seed fields may either produce small ( $\leq$  Mpc, e.g. Chernin 1967) or large (e.g. Zel'dovich 1970; Turner and Widrow 1988) coherence lengths, whose structure may still persist until today (e.g. Hutschenreuter et al. 2018), in the emptiest cosmic regions, possibly also carrying information on the generation of primordial helicity (e.g. Semikoz and Sokoloff 2005; Campanelli 2009; Kahniashvili et al. 2016).

Owing to uncertainties in the physics of high energy regimes in the early Universe, the uncertainty in the outcome of most of the above scenarios is rather large and fields in the range of  $\sim 10^{-34}$ – $10^{-10}$  G are still possible.

The presence of magnetic fields with rms values larger than a few co-moving  $\sim$  nG on  $\leq$  Mpc scales at  $z \approx 1100$  is presently excluded by the analysis of the CMB angular power spectrum by Planck (Ade et al. 2016; Trivedi et al. 2014), while higher limits are derived for primordial fields with much larger coherence length (Barrow et al. 1997). Conversely, the lack of detected Inverse Compton cascade around high redshift blazars was used to set lower limits<sup>1</sup> on cosmological seed fields of  $\geq 10^{-16}$  G on  $\sim$  Mpc (Dolag et al. 2009, 2011; Neronov and Vovk 2010; Arlen et al. 2014; Caprini and Gabici 2015; Chen et al. 2015).

## 2.2 Seeding from Galactic Outflows

At lower redshift ( $z \leq 6$ ) galactic feedback can transport magnetic fields from galactic to more rarefied scales such as galaxy clusters. In lower mass haloes, star formation drives winds of magnetized plasma into the circum-galactic medium (e.g. Kronberg et al. 1999; Völk and Atoyan 2000; Donnert et al. 2009; Bertone et al. 2006; Samui et al. 2017) and into voids (Beck et al. 2013b). At the high mass end, active galactic nuclei (AGN) can magnetize the central volume of clusters through jets (e.g. Dubois and Teyssier 2008; Xu et al. 2009; Donnert et al. 2009) and even the intergalactic medium during their violent quasar phase (Furlanetto and Loeb 2001). Just taking into account the magnetization from dwarf galaxies in voids, a lower limit of the magnetic field in voids has been derived as  $\sim 10^{-15}$  G (Beck et al. 2013b; Samui et al. 2017).

If magnetic fields have been released by processes triggered during galaxy formation, they might have affected the transport of heat, entropy, metals and cosmic rays in forming cosmic structures (e.g. Planelles et al. 2016; Schekochihin et al. 2008).

Additional processes such as the “Biermann-battery” mechanism (Kulsrud et al. 1997), aperiodic plasma fluctuations in the inter-galactic plasma (Schlickeiser et al. 2012), resistive mechanisms (Miniati and Bell 2011) or ionization fronts around the first stars (Langer et al. 2005) might provide additional amplification to the primordial fields starting from  $z \leq 10^3$ , i.e. after recombination.


## 3 Magnetic Field Amplification in the Intra-cluster Medium

### 3.1 Amplification by Compression

From the third term in the induction equation (Eq. (1)) we find that a positive divergence of the velocity field  $\nabla \cdot \mathbf{v}$ , i.e. a net inflow, results in the growth of the magnetic field (Sur et al.

<sup>1</sup>See however Broderick et al. (2012) for a different interpretation.

**Fig. 1** Magnetic field strength as a function of over density in cosmological SPH simulations. Starting from 3 different cosmological seed field strengths:  $2 \times 10^{-13}$  G (dark green),  $2 \times 10^{-12}$  G (black),  $8 \times 10^{-12}$  G (green) (Dolag et al. 2008, 2005a). Adiabatic evolution solely by compression in grey. Runs with galactic seeds are in red and blue



2012). Indeed, it is a basic result of MHD that magnetic flux  $\Phi$  is conserved (e.g. Kulsrud and Ostriker 2006) leading to the scaling of the magnetic field with density:

$$B(\rho) \propto B(z_*) \left( \frac{\rho}{\langle \rho \rangle} \right)^{2/3}. \tag{2}$$

For a galaxy cluster with an average over-density of  $\Delta = \rho/\langle \rho \rangle \approx 100$  this means that adiabatic compression can amplify the seed field by up to a factor of  $\sim 20$  within the virial radius (or  $\sim 180$  within the cluster core, where the density can be  $\approx 2500$  the mean density). This refers to the average magnetic field inside a radius of the cluster. The peak density and magnetic field can be much higher. However, depending on the redshift and environment of the seed fields, the expectation from adiabatic amplification can be lower. Nonetheless, observations find a scaling exponent of magnetic field strength with cluster density of 0.5–1, which is compatible with amplification by compression.

In Fig. 1 we reproduce a central result from early cosmological SPMHD (smooth particle magneto-hydrodynamics) simulations (Dolag et al. 2005a, 2008). They show the magnetic field strength over density in a cosmological simulation with cosmological seed fields of  $B(z_*) = 2 \times 10^{-13}$  G (dark green),  $B(z_*) = 2 \times 10^{-12}$  G (black),  $B(z_*) = 8 \times 10^{-12}$  G (dark green) co-moving, seeded at  $z_* = 20$  alongside the analytical expectation from Eq. (2). Runs with galactic seeding in blue and red. At central cluster over-densities ( $\rho/\langle \rho \rangle > 1000$ ), all but one simulations reach  $\mu\text{G}$  field strengths. Thus different seeding models are indistinguishable here. Differences to galactic field seeding appear only at lower densities.

In runs with a cosmological seed field, amplification is mostly caused by compression below over-densities of 1000. At larger over-densities, a dynamo caused by velocity gradients along the field lines in the first term of the induction equation (1) operates and leads to much higher field strengths. This is characteristic for turbulence in structure formation, which we will discuss next. Simulations of the cosmic dynamo and their limitations will be covered later in Sect. 4.

### 3.2 A Brief Introduction to Turbulence

Let us first introduce a few key concepts of turbulence used throughout the review. For a more detailed exposure, we refer the reader to the vast literature available on Astrophysical turbulence (e.g. Landau and Lifshitz 1966; Kulsrud and Ostriker 2006; Lazarian et al. 2009).

A key idea of the Kolmogorov picture of turbulence is that random fluid motions with *velocity dispersion*<sup>2</sup>  $v$  of size or scale  $l$  (“eddies”) break up into two eddies of half the size due to the convective  $\mathbf{v} \cdot \nabla \mathbf{v}$  term in the fluid equations. This process constitutes a local energy transfer from large to small scales at a rate  $kv$ , where  $k = 2\pi/l$  is the wave vector. This process continues at each smaller length scale which leads to a cascade of velocity fluctuations down to smaller scales with decreasing kinetic energy. At an *inner scale*  $k_v$ , the local kinetic energy becomes comparable to viscous forces, which dissipate the motion into thermal energy or, in case of a dynamo, also magnetic energy via the Lorentz force. At each scale, the cascading time scale is the *eddy turnover time*  $\tau_l = l/v_l$  and for continuous injection of velocity fluctuations at the *outer scale* a steady state is reached. If the kinetic energy density of these fluctuations is  $1/2\rho v^2 = \rho/2 \int I(k) dk$  (assuming isotropy), then it can be shown that the velocity power spectrum  $I(k)$  is (Kolmogorov 1941, 1991):

$$I(k) \propto v_0^2 \frac{k_0^{2/3}}{k^{5/3}}, \quad (3)$$

where  $v_0$  is the velocity dispersion of the largest eddy at scale  $k_0$ . We note that  $v_0$  is a velocity fluctuation on top of the mean. This dispersion of the associated random velocity field then scales as  $v^2 \propto l^{2/3}$ . It follows that the energy of turbulence is dominated by the largest scales and that viscous forces are important close to the dissipative inner scale, where motions are slowest. The range of scales where Eq. (3) is valid is called the *inertial range*, and the *Reynolds number* is defined as:

$$R_e = \frac{v_0}{k_0 \nu} \quad (4)$$

$$\propto \left( \frac{l_0}{l_\nu} \right)^{4/3} \quad (5)$$

with the *kinematic viscosity*  $\nu$ . The role of small scales is universal in the sense that the cascading does not depend on the driving scale or velocity (assuming homogeneity, scale invariance, isotropy and locality of interactions) (Schekochihin and Cowley 2007). We note that turbulence is not limited to velocity fluctuations around a mean caused by a superposition of velocity eddies. The velocity field causes density and pressure fluctuations as well, because these are coupled via the fluid equations. For sub-sonic turbulence the fluctuations will be adiabatic. This has been used to place an upper limit on the kinematic viscosity in the Coma cluster of  $\nu < 3 \times 10^{29}$  cm<sup>2</sup>/s on scales of 90 kpc using X-ray data (Schuecker et al. 2004).

Whether or not the stage of the dynamo amplification is reached in an astrophysical system ultimately depends on the magnetic Reynolds number (Eq. (15)) and on the nature of the turbulent forcing in the ICM (Federrath et al. 2014; Beresnyak and Miniati 2016). The magnetic Reynolds number is set by the outer scale and the dissipation scale, so it is worth

<sup>2</sup>Note that velocity and velocity dispersion (i.e. root-mean-square of the power-spectrum at scale  $k$ ) are used somewhat interchangeably in the literature. Similarly we denote the velocity dispersion with  $v$  as the distinction is usually clear by context.

discussing the latter next. For galaxy clusters, these scales are connected to the physics of the ICM plasma.

### 3.2.1 The Spitzer Model for the ICM

As noted in the introduction, most theoretical and numerical studies approximate the ICM plasma as a fluid. However, the MHD equations as a statistical description of the many-body plasma are applicable only, if equilibration processes between ionized particles act on length and time scales much smaller than “the scales of interest” of the fluid problem, i.e. if collisional equilibrium among particles (protons, electron, metal ions) is maintained so local particle distributions become Maxwellian and temperature and pressure are well defined (Landau and Lifshitz 1966).

In the “classic” physical picture of the ICM, this arises from ion-ion Coulomb scattering, with a viscosity  $\nu_{ii}$ , over a mean free path  $l_{\text{mfp}}$  which is given by the Spitzer model for fully ionized plasmas (Spitzer 1956). It can be shown that a *whole cluster* is then “collisional” in the sense that  $r_{\text{vir}} \gg l_{\text{mfp}}$  (Sarazin 1986), with:

$$l_{\text{mfp}} \approx 23 \left( \frac{n_{\text{th}}}{10^{-3} \text{ cm}^{-3}} \right)^{-1} \left( \frac{T}{10^8 \text{ K}} \right)^2 \text{ kpc.} \quad (6)$$

Under these conditions, the Reynolds number (Eq. (5)) of the ICM in a cluster during e.g. a major merger is (e.g. Brunetti and Lazarian 2007):

$$R_e = \frac{Lv_L}{\nu_{ii}} \quad (7)$$

$$\approx 52 \frac{v_L}{10^3 \text{ km/s}} \cdot \frac{L}{300 \text{ kpc}} \cdot \frac{n}{10^{-3} \text{ cm}^{-3}} \cdot \left( \frac{T}{8 \text{ keV}} \right)^{-5/2} \cdot \left( \frac{\log \Lambda}{40} \right) \quad (8)$$

where  $L$  is a typical eddy size (ideally the injection scale of turbulence),  $\log \Lambda$  is the Coulomb logarithm (Longair 2011) and  $v_L$  is the rms velocity within the scale  $L$ . Thus based on typical values of the ICM, the Reynolds number would hardly reach  $R_e \sim 10^2$  in most conditions.

In contrast, rotation measures inferred from observations of radio galaxies have demonstrated field reversals on kpc scales, implying much larger Reynolds numbers (Laing et al. 2008; Govoni et al. 2010; Bonafede et al. 2013; Kuchar and Enßlin 2011; Vacca et al. 2012). Turbulent gas motions from AGN feedback have been observed directly with the Hitomi satellite in the Perseus cluster (Aharonian et al. 2016) showing velocity dispersions of  $\sim 200$  km/s on scales of  $< 60$  kpc. This is not compatible with a medium based solely on Coulomb collisions.

Thus it is unavoidable to consider a more complex prescription of the ICM plasma. In the future, stronger constraints on the velocity structure of gas motions in galaxy clusters will be provided by the XIFU instrument on the Athena satellite (Ettori et al. 2013; Roncarelli et al. 2018).

We note that modern numerical simulations of galaxy clusters reach and exceed spatial resolutions of the Spitzer collisional mean free path. It follows that other processes than Coulomb scattering have to maintain collisionality on smaller scales for these simulations to be valid at all. Just adding a magnetic field to the Spitzer model, i.e. Coulomb scattering plus a Lorentz force, does not suffice to make the ICM collisional on kpc scales. In a micro-physical sense the MHD magnetic field is a mean magnetic field that arises *after* averaging over micro-physical quantities (adiabatic invariants Schlickeiser 2002).

### 3.2.2 Turbulence and the Weakly-Collisional ICM

In the MHD limit, turbulence can excite three MHD waves, of which two have compressive nature (fast and slow modes, similar to sound waves) and one is solenoidal (Alfvén mode). The Alfvén speed is given by Alfvén (1942):

$$v_A = \frac{B}{\sqrt{4\pi\rho}} \quad (9)$$

$$= 69 \frac{B}{1 \mu\text{G}} \left( \frac{n_{\text{th}}}{10^{-3} \text{ cm}^{-3}} \right)^{-1/2} \text{ km/s}, \quad (10)$$

with the number density of (thermal) ions  $n_{\text{th}}$ .

Numerical simulations of cluster formation find turbulent velocities at the outer scale of several hundred km/s (Miniati 2014), which means that ICM turbulence starts off *super-Alfvénic* on the largest scales. Thus the magnetic field is dynamically not important near the outer scale and field topology is shaped by fluid motion.

Integrating Eq. (3) over  $k$ , we find that  $v_l \propto l^{1/3}$  and with Eq. (9) the *Alfvén scale*, where the magnetic field back-reacts on turbulent motions (Brunetti and Lazarian 2007):

$$l_A \approx 100 \left( \frac{B}{\mu\text{G}} \right)^3 \left( \frac{L_0}{300 \text{ kpc}} \right) \left( \frac{V_L}{10^3 \text{ km/s}} \right)^{-3} \left( \frac{n_{\text{th}}}{10^{-3} \text{ cm}^{-3}} \right)^{\frac{3}{2}} \text{ pc}, \quad (11)$$

which is already smaller than the classical mean free path derived before and leads to a Reynolds number of a few 1000. As we will see, this scale is crucial to numerically resolve magnetic field growth by turbulence.


In principle, one has to consider three separate turbulent cascades, whose interplay changes close around Alfvén scale (see Brunetti and Lazarian 2011b, and ref. therein). Here the character of turbulence dramatically changes. The Lorentz force introduces strong anisotropy to fluid motions, viscosity and turbulent eddies become anisotropic and non-local interactions between modes in the turbulent cascade start to be important. See Goldreich and Sridhar (1997), Schekochihin and Cowley (2007) for a more detailed picture of these processes.

That leaves us to ask, what is it that keeps the ICM collisional on scales much smaller than the Alfvén scale, so MHD is applicable at all? Schekochihin et al. (2005b), Beresnyak and Lazarian (2006), Schekochihin and Cowley (2007), Schekochihin et al. (2008) propose that due to the large Spitzer mean free path, the non-ideal MHD equations are not sufficient to estimate viscosity and obtain a Reynolds number for the ICM. Kinetic calculations reveal that particle motions perpendicular to the magnetic field are suppressed and motions parallel to the field can exist and excite firehose and mirror instabilities. The instabilities inject MHD waves, which act as scattering agents (magnetic mirrors). Scattering off these self-excited modes isotropizes particle motions on very small length and time scales. This picture is confirmed also by hybrid-kinetic simulations (Kunz et al. 2014).

Under these conditions, a lower limit of the viscous scale of the ICM is given by the mobility of thermal protons in a magnetic field, which is the Larmor radius (e.g. Schekochihin et al. 2005b; Beresnyak and Miniati 2016; Brunetti and Lazarian 2011b):

$$l_{\text{mfp}} = r_{\text{Larmor}} \quad (12)$$

$$\approx 3 \cdot 10^{-12} \text{ kpc} \left( \frac{T}{10 \text{ keV}} \right) \left( \frac{B}{\mu\text{G}} \right)^{-1}. \quad (13)$$



**Fig. 2** Left: Cartoon illustrating the stretching and folding for magnetic field lines on small scales from Schekochihin et al. (2002b). Right: Cartoon from Cho et al. (2009) depicting the growth of magnetic energy in driven turbulence simulations with very weak initial magnetic field. The initial seed field sets the timescale for the end of the kinematic dynamo and the beginning of the non-linear dynamo

In this case, the effective Reynolds number of the ICM becomes:

$$R_{e,eff} = \left(\frac{l_0}{l_v}\right)^{4/3} \sim 10^{19}. \tag{14}$$

This estimate predicts a highly turbulent ICM down to non-astrophysical scales and establishes collisionality on scales of tens of thousands of kilometers. This is good news for simulators, because the fluid approximation is well motivated in galaxy clusters and probably valid down to scales forever out of reach of simulations (Santos-Lima et al. 2014, 2017).

The bad news is that the physics of the medium is complicated, so that e.g. transport properties of the ICM are dominated by scales out of reach for simulations and observations. One example is heat conduction, where some estimates from kinetic theory predict no conduction in the weakly-collisional limit (Schekochihin et al. 2008; Kunz 2011). Indeed, only an upper limit was found by comparing observations with simulations (ZuHone et al. 2015b). Thus, the properties of the medium cannot be constrained any further. Additionally, the likely presence of cosmic-ray protons makes the picture of generation and damping of compressive and Alfvén modes/turbulence even more involved (Fig. 5) (Schlickeiser 2002; Brunetti and Lazarian 2011c; Brunetti et al. 2004).

Now that we have established that MHD is very likely applicable down to sub-pc scales, we can discuss how (large-scale) magnetic fields can be amplified by turbulence in the MHD limit.

### 3.3 The Small-Scale Dynamo

If a magnetic field is present in a turbulent flow, the properties of turbulence can change significantly due to the back-reaction of the field on the turbulent motions (Kraichnan and Nagarajan 1967; Goldreich and Sridhar 1997). In a magnetic dynamo, the kinetic energy of turbulence is transformed into magnetic energy, which is a non-trivial theoretical problem. The dissipation of magnetic energy into heat occurs at the resistive scale  $l_\eta$  and the magnetic Reynolds number is defined as:

$$R_m = \frac{v_0}{k_0 \eta} \tag{15}$$

$$\propto \left(\frac{l_\eta}{l_0}\right)^{4/3} \tag{16}$$

The magnetic Prandtl number relates resistive with diffusive scales Eq. (14).

$$P_m = \frac{\nu}{\eta} \quad (17)$$

$$= \frac{R_m}{R_e} = \left(\frac{l_\nu}{l_\eta}\right)^{4/3}$$

For a theoretical framework of the gyrokinetics on small scales, including a discussion on cluster turbulence, we refer the reader to Schekochihin et al. (2009).

In a simplified picture, magnetic field amplification by turbulence is a consequence of the stretching and folding of pre-existing field lines by the random velocity field of turbulence, which amplifies the field locally due to flux conservation (Fig. 2, left) (Batchelor 1950; Biermann and Schlüter 1951). If a flux tube of radius  $r_1$  and length  $l_1$  with magnetic field strength  $B_1$  is stretched to length  $l_2$  and radius  $r_2$ , mass conservation leads to:

$$\frac{r_2}{r_1} = \sqrt{\frac{l_1}{l_2}}. \quad (18)$$

The magnetic flux  $S_1 = \pi r_1^2 B$  is conserved in the high- $\beta$  regime, so for an incompressible fluid:


$$B_2 = B_1 \frac{l_2}{l_1} \quad (19)$$

By e.g. folding or shear (Fig. 2, left) the field can be efficiently amplified (Vainshtein and Zel'dovich 1972; Schekochihin et al. 2002a). Repeating this process leads to an exponential increase in magnetic energy, if the field does not back-react on the fluid motion (Fig. 2, right). In a turbulent flow the folding occurs on a time scale of the smallest eddy turnover time, i.e. close to the viscous scale. Flux tubes are tangled and merged, and their geometry/curvature is set by the resistive and viscous scales of the flow. The energy available for magnetic field growth is the rate of strain  $\delta v/l$  (Schekochihin et al. 2005a). We note that due to the universality of scales in turbulence, the dynamo process does not depend on the actual magnetic field strength and time scale of the system. As long as the conditions for a small scale dynamo are satisfied, field amplification will proceed as shown in Fig. 2, right.

For a small ( $10^{-13}$  G) initial seed field in galaxy environments or proto-clusters (see Sect. 2), back-reaction is negligible,  $P_m$  is very large and a *small-scale dynamo* (SSD) operates in the *kinematic regime* of exponential amplification without back-reaction (Kulsrud and Anderson 1992). The SSD proceeds from small to large scales in an inverse cascade starting at the resistive scale. A rigorous treatment of this process based on Gaussian random fields in the absence of helicity was first presented by Kazantsev (1968), for an instructive application to proto-clusters see e.g. Federrath et al. (2011b), Schober et al. (2013), and Latif et al. (2013). For a unique experimental perspective on the kinematic dynamo see Meinecke et al. (2015). In Fig. 3, we reproduce the time evolution of magnetic energy (left) and of the magnetic and kinetic power spectra (right) from an idealized simulation of the MHD dynamo (Cho et al. 2009). Here  $k_\nu = 1/l_\nu \approx 100$ , and the kinematic dynamo proceeds until  $t = 15$ . An instructive numerical presentation can be also found in Porter et al. (2015), a detailed exposure is presented in Schekochihin et al. (2004).

The exponential growth of the kinematic dynamo is stifled quickly (Brandenburg 2011), once the magnetic field starts to back-react on the turbulent flow. The dynamo then enters





**Fig. 3** Left: Evolution of magnetic energy over time in a simulation of driven turbulence. The transition from kinematic to non-linear dynamo occurs at  $t = 15$ . Right: Magnetic field energy spectra over wave number for different times of the same run. Both figures by Cho et al. (2009)

the *non-linear regime* and turbulence grows a steep inverse cascade with an outer magnetic scale  $l_B$ . In Fig. 3, this occurs for  $t > 15$  and  $k_B = 1/l_B \approx 10$  at  $t = 40$ . In principle, growth will continue until equipartition with the turbulent kinetic energy is attained (Haugen and Brandenburg 2004; Brandenburg and Subramanian 2005; Cho et al. 2009; Porter et al. 2015; Beresnyak and Miniati 2016).


What does this mean for galaxy clusters? Above we had motivated a lower limit for the viscous scale in proto-clusters of around 1000 km (Eq. (13)) and Reynolds numbers of up to  $10^{19}$ . The resistive scale is highly uncertain, but likely small enough for an SSD to occur. The large Reynolds number leads to a growth timescale of the kinematic dynamo of  $\tau \approx 1000$  yrs (Schekochihin et al. 2002b, 2004; Beresnyak and Miniati 2016). It is clear that this exponential growth is so fast that it will complete in large haloes before galaxy clusters start forming at redshifts 2–1. The kinematic dynamo efficiently amplifies even smallest seed fields until back-reaction plays a role, i.e. the Alfvén scale approaches the viscous scale.

Depending on the physics of the seeding mechanism, the kinematic phase will take place in the environment of high redshift galaxies that is polluted by jets and outflows, in proto-clusters or, in the case of a cosmological seed field, in all collapsing over-dense environments at high redshift (Zeldovich et al. 1983; Kulsrud and Anderson 1992; Kulsrud et al. 1997; Latif et al. 2013).

However, contrary to the idealized turbulence simulations shown in Fig. 3, turbulent driving in clusters occurs highly episodic and at multiple scales at once (Sect. 3.5), so the equipartition regime is never reached. Instead, the magnetic field strength and topology will depend on the driving history of the gas parcel under consideration. It is also immediately clear that as opposed to amplification by isotropic compression, this dynamo erases all imprint of the initial seed field. Thus we cannot hope to constrain seeding processes from magnetic fields in galaxy clusters, but instead have to look to filaments and voids, where the dynamo may not be driven efficiently.

### 3.4 Cosmic-Ray Driven Amplification and Plasma Effects

Magnetic fields can be amplified by a range of effects caused by cosmic rays. Current-driven instabilities, e.g. (Bell 2004), have been shown to amplify magnetic fields by considerable factors (Riquelme and Spitkovsky 2010). The electric current that drives this instability comes from the drift of CRs. The return electric current of the plasma leads to a transverse force that can amplify transverse perturbations in the magnetic field. Bell (2004)



**Fig. 4** Ion number density (top) and magnetic field strength (bottom) for a parallel shock wave with Mach number  $M = 20$  at  $1000 \omega_c^{-1} = mc/eB_0$  from (Caprioli and Spitkovsky 2014b)


pointed out that the fastest instability is caused by the return background plasma current that compensates the current produced by CRs streaming upstream of the shock. It is important to note that this instability is non-resonant and can be treated using ideal MHD. The Bell or non-resonant streaming (NRS) instability has been tested in various numerical studies using a range of methods ranging from pure MHD (Zirakashvili and Ptuskin 2008), full PIC (Riquelme and Spitkovsky 2011), hybrid (Caprioli and Spitkovsky 2014a,b, Fig. 4) to Vlasov or PIC-MHD (Reville and Bell 2013; Reville et al. 2008; Bai et al. 2015); see Marcowith et al. (2016) for a review. In strong SNR shocks, a non-resonant long-wavelength instability can amplify magnetic fields as well (Bykov et al. 2009, 2011), but this has not been confirmed by simulations. A full non-linear calculation is needed to take into account the feedback of the CRs on the shock structure that may lead to a significant modification of the shocks structure (e.g. Malkov and O’C Drury 2001; Vladimirov et al. 2006; Bykov et al. 2014). Recent  $\gamma$ -ray observations of SNR challenge this picture, so CR spectra might be steeper than the test-particle prediction (Caprioli 2012; Slane et al. 2014). All the aforementioned effects operate on length scales comparable to the gyro-radius of protons.

Filamentation instabilities can act on larger scales, as do models where CRs drive a turbulent dynamo (Drury and Downes 2012; Brüggén 2013). In the latter case, the turbulence is caused by the cosmic-ray pressure gradient in the upstream region which exerts a force on the upstream fluid that is not proportional to the gas density. Density fluctuations then lead to fluctuations in the acceleration which, in turn, produce further density fluctuations. CRs are also able to generate strong magnetic fields at shock fronts which is invoked to explain the high magnetic field strengths in several historical supernova remnants. This was first studied in the context of the high magnetic field strengths deduced from X-ray observations of supernova remnants. In fast shocks, the streaming of CRs into the upstream region triggers a class of plasma instabilities that can grow fast enough to produce very strong magnetic fields (Lucek and Bell 2000).

More recently, Reville and Bell (2013) have studied a CR-driven filamentation instability that also results from CR streaming, but contrary to the Bell-instability generates long-wavelength perturbations. Caprioli and Spitkovsky (2014b) have investigated CR-driven filamentation instabilities using a di-hybrid method where electrons are treated as a fluid and protons as kinetic particles. While progress in this field has grown substantially over the past years, very few PIC simulations for weak shocks in high- $\beta$  plasmas have been done (e.g. Guo et al. 2016).

In analytical work (e.g. Melville et al. 2016), it has been shown that microphysical plasma instabilities can produce a more efficient small-scale dynamo than its MHD counterpart de-

**Fig. 5** Cartoon depicting the cascade of only compressive turbulence over length scale in galaxy clusters, considering damping from thermal ions and cosmic-ray protons (Donnert and Brunetti 2014)




scribed above. In this picture, shearing motions drive pressure anisotropies that excite mirror or firehose fluctuations (as seen in direct numerical simulations of collisionless dynamo; see Rincon et al. 2016). These fluctuations lead to anomalous particle scattering that lead to field growth. As shown in Mogavero and Schekochihin (2014), these scatterings can decrease the effective viscosity of the plasma thereby allowing the turbulence to cascade down to smaller scales and thus develop greater rates of strain and amplify the field faster. Within a number of large eddy turn-over times, this process can result in magnetic fields that saturate near equipartition with the kinetic energy of the ICM.

While the total budget of cosmic ray protons stored in clusters is now constrained to  $\leq 1\%$  (on average) for the thermal gas energy by the latest collection of Fermi-LAT data (Ackermann et al. 2014), it cannot be excluded that a larger fraction of cosmic rays may exist close to shocks in the intra-cluster medium. At present, the limits that can be derived from  $\gamma$ -rays are of  $\leq 15\%$ , at least in the case of the (nearby) relics in Coma (Zandanel and Ando 2014).

### 3.5 Processes that Drive Turbulence in Clusters

The accretion of gas and Dark Matter subunits is a main driver of turbulence in clusters. During infall, gas gets shock-heated around the virial radius (Mach numbers  $\sim 10$ ). In major mergers, the displacement of the ICM creates an eddy on the scale of the cluster core radii (e.g. Donnert and Brunetti 2014). Shear flows generated by in-falling substructure inject turbulence through Kelvin-Helmholtz (K-H) and Rayleigh-Taylor (R-T) instabilities (e.g. Subramanian et al. 2006; Su et al. 2017; Khatri and Gaspari 2016). Feedback from central AGN activity, radio galaxies and galactic winds inject turbulence on even smaller scales (e.g. Churazov et al. 2004; Brüggen et al. 2005a; Gaspari et al. 2011). As a result of this complex interplay of episodic driving motions on scales of half a Mpc to less than a kpc, the intra-cluster medium is expected to include weak-to-moderately-strong shocks ( $\mathcal{M} \leq 5$ ) and hydrodynamic shear, leading to a turbulent cascade down to the dissipation scale.

The solenoidal component (Alfvén waves) of the cascade will drive a turbulent dynamo, while the compressive component (fast & slow modes) produces weak shocks and adiabatic compression waves, which can in turn generate further small-scale solenoidal motions (e.g., Porter et al. 2015; Vazza et al. 2017b). The relative contributions from both components



**Fig. 6** Left: Gas motions in the first Eulerian simulation of a merging cluster (Schindler and Mueller 1993). Central and Right panel: projected enstrophy energy flux for a state-of-the-art Eulerian simulation with ENZO at  $z = 1$  and  $z = 0$ , taken from Wittor et al. (2017b)

will depend on the turbulent forcing and its intensity (Federrath et al. 2011b; Porter et al. 2015). Compressive and solenoidal components of the turbulent energy are also expected to accelerate cosmic-ray protons and electrons via second-order Fermi processes, which again alters the properties of turbulence on small scales (see Brunetti and Jones 2014, for a review). In Fig. 5 we reproduce a cartoon plot of the compressive cascade in clusters from Donnert and Brunetti (2014) that depicts the relevant scales: the classical mean free path (Eq. (6)), the Alfvén scale (Eq. (11)) and the dissipation scales, if the cascade is damped by thermal protons ( $k > 10^{-2}$ ) or CR protons ( $k \approx 1$ ). The graph also includes the sound speed and the Alfvén speed and marks the regions accessible by current cosmological simulations. A more involved graph can be found in Brunetti and Jones (2014).

Note that the simple “Kolmogorov” picture of turbulence (Sect. 3.2) with a single well defined injection scale, an inertial range and a single dissipation scale is oversimplified in galaxy clusters. As argued above, structure formation leads to an increase of the outer/driving scale with time and injection concurrently takes place at many smaller scales and can be highly intermittent. Thus a strictly-defined inertial range does probably not exist and turbulence may be more loosely defined in clusters than in other fields of astrophysics. Cosmological simulations can be used to capture the complexity of these processes.

## 4 Simulations of Turbulence and the Small Scale Dynamo in Clusters

### 4.1 Simulations of Cluster Turbulence

Simulations of merging clusters have been pioneered by Evrard (1990), Thomas and Couchman (1992), who reported a shock traveling outward during a merger. Schindler and Mueller (1993) for the first time used a Eulerian PPM scheme with  $60^3$  zones to follow the gas dynamics in an idealized merger (Fig. 6, left) (see also Roettiger et al. 1993, 1997). Using idealized adaptive mesh refinement Eulerian merger simulations, Ricker and Sarazin (2001) for the first time report ram pressure stripping and turbulence, with eddy sizes of “several hundred kpc” [...] “pumped by DM driven oscillations of the gravitational potential”. Takizawa (2005), Asai et al. (2004) used a TVD scheme to study the driving of shocks and turbulence by substructure in idealized cluster simulations. They focused on the injection of instabilities and gas stripping (see Sect. 6).

In cosmological simulations, turbulence was first studied by Dolag et al. (2005b) using SPH with a low viscosity scheme (for shocks see Miniati et al. 2000). They find subsonic velocity dispersions of 400–800 km/s on scales of 20 to 140 kpc, with turbulent energy fractions of 5–30 per cent and a trend for higher turbulent energies in higher mass clusters. Turbulent energy spectra from their simulations were flatter than the Kolmogorov expectation, but might have been limited by numerics (see Sect. 4.4). Their work was extended to a sample of 21 clusters by Vazza et al. (2006), who provided scaling laws for the turbulent energy over cluster mass, see Valdarnini (2011) for a later study.

In a seminal contribution, Ryu et al. (2008) studied the generation and evolution of turbulence in a Eulerian cosmological cluster simulation. They showed that turbulence is largely solenoidal, not compressive, with subsonic velocities in clusters and trans-sonic velocities in filaments. In agreement with prior SPH simulations, they find a clear trend of rms velocity dispersion with cluster mass and turbulent energy fractions/pressures of 10–30%. They also propose a vorticity based dynamo model, which we will discuss in Sect. 4.2.


The influence of turbulent pressure support on cluster scaling relations was studied by Nagai et al. (2007), Lau et al. (2009), Shaw et al. (2010), Burns et al. (2010), Battaglia et al. (2012), Nelson et al. (2014), Schmidt et al. (2017). Consistently, turbulent pressure increases with radius in simulated clusters, which is related to the increased thermal pressure caused by the central potential of the main DM halo. An analytic model for non-thermal pressure support was presented by Shi and Komatsu (2014), and also validated by numerical simulations (Shi et al. 2015, 2016). First power spectra of turbulence in Eulerian cosmological cluster simulations were presented by Xu et al. (2009), Vazza et al. (2009). Their kinetic spectra roughly follow the Kolmogorov scaling. The simulations reach an “injection region” of turbulence larger than 100 kpc, an inertial range between 100 kpc and 10 kpc and a dissipation scale below 10 kpc. Thus their Reynolds number was 10–100.

The next years saw improvements in resolution of cluster simulations, due to the inevitable growth in computing power. Increasingly higher Reynolds numbers could be reached and/or additional physics could be implemented usually with adaptive mesh refinement (AMR). Vazza et al. (2011) studied a sample of simulated clusters with Reynolds number of up to 1000. They also developed new filtering techniques to estimate turbulent energy locally. They showed that the turbulent energy in relaxed clusters reach only a few percent. Maier et al. (2009), Iapichino et al. (2011) added a subgrid-scale model for unresolved turbulence to their simulations and studied the evolution of turbulent energy. They found that peak turbulent energies are reached at the formation redshift of the underlying halo. Their subgrid model shows that unresolved pressure support is usually not a problem in cluster simulations, and that half of the simulated ICM shows large vorticity. Paul et al. (2011) simulated a sample of merging clusters and found a scaling of turbulent energy with cluster mass as  $\propto M^{5/3}$ , consistent with earlier SPH results (Vazza et al. 2006). The influence of minor mergers on the injection of turbulence in a idealized scenario of a cool core cluster was simulated with anisotropic thermal conduction by Ruszkowski and Oh (2011). They found that long-term galaxy motions excite subsonic turbulence with velocities of 100–200 km/s and give a detailed theoretical model for the connection between vorticity and magnetic fields.

Vazza et al. (2012) used an improved local filter to estimate the turbulent diffusivity in their simulations as  $D_{\text{turb}} \approx 10^{29-30} \text{ cm}^2/\text{s}$  and identify accretion and major mergers as dominant drivers of cluster turbulence.

In a series of papers, Miniati (2014, 2015) introduced static Eulerian mesh refinement simulations to the field. They reach a peak resolution of  $\approx 10$  kpc covering the entire virial radius of a massive galaxy cluster with a PPM method. Consistent with previous studies





**Fig. 7** Vorticity map for the innermost regions of a simulated  $\sim 10^{15} M_{\odot}$  galaxy cluster at high resolution, in the “Matrioska run” by Miniati (2014)


they find that shocks generate 60% of the vorticity in clusters. Their adiabatic simulations show turbulent velocity dispersions above 700 km/s, regardless of merger state. The analysis using structure functions reveals that solenoidal/incompressible turbulence with a Kolmogorov spectrum dominates the cluster, while compressive turbulence with a Burgers slope (Burgers 1939) become more important towards the outskirts. They propose that a hierarchy of energy components exists in clusters, where gravitational energy is mostly dissipated into thermal energy, then turbulent energy and finally magnetic energy with a constant efficiency (Miniati and Beresnyak 2015). Vorticity maps from their approach are reproduced in Fig. 7.

In the most recent studies, the resolution has been improved to simulate the first early baroclinic injection of vorticity in cluster outskirts (e.g. Vazza et al. 2017b; Iapichino et al. 2017) as well as its later amplification via compression/stretching during mergers (Wittor et al. 2017a). Using the Hodge-Helmholtz decomposition, high resolution Eulerian simulations measure a very large fraction of turbulence being dissipated into solenoidal motions (Miniati 2014; Vazza et al. 2017b; Wittor et al. 2017a). Baroclinic motions inject enstrophy on large scales, while dissipation and stretching terms govern its evolution.


Recent simulations using Lagrangian methods focus on including more subgrid physics in the setup to study the influence of magnetic fields on galaxy formation. Marinacci et al. (2015) show that the redshift evolution of the rms velocity fluctuations in the “Illustris TNG” galaxy formation simulations is independent of seed magnetic fields.

## 4.2 Cosmological Simulations of Magnetic Fields in Galaxy Clusters

Pioneering studies of magnetic fields in simulated large-scale structures were conducted by De Young (1992), Kulsrud et al. (1997), Roettiger et al. (1999). First full MHD simulations of cluster magnetic fields from nG cosmological seeds have been presented by Dolag et al. (1999, 2002), Bonafede et al. (2011). They found a correlation of the magnetic field strength the ICM gas density with an exponent of 0.9, using smooth particle hydrodynamics (SPH) (Dolag and Stasyszyn 2009; Beck et al. 2016). This is close to the theoretical expectation for spherical collapse (Fig. 1, Eq. (2)) and it is in-line with observations from Faraday rotation measures. In the center of clusters, their simulations obtain a magnetic field strength



**Fig. 8** Projection of magnetic field strength in three cosmological simulations using different MHD approaches and solvers. Left: non-radiative GADGET SPH simulations with galactic seeding by Donnert et al. (2009), based on the MHD method by Dolag and Stasyszyn (2009). Middle: non-radiative ENZO MHD simulation on a fixed grid by Vazza et al. (2014) using the Dender cleaning (Dedner et al. 2002); Right: Simulation with full “Illustris TNG” galaxy formation model using a Lagrangian finite volume method (Marinacci et al. 2018b)




**Fig. 9** 3-dimensional kinetic and magnetic power spectra in ENZO MHD simulations by Xu et al. (2009) (assuming a seeding of magnetic fields by AGN) and by Vazza et al. (2018), assuming a primordial magnetic field of 0.1 nG (comoving), as a function of resolution

of 3–6  $\mu\text{G}$ , over a wide range of cluster masses. Subsequently the simulations were used to model giant radio haloes (Dolag and Enßlin 2000; Donnert et al. 2010), the influence of the field on cluster mass estimates (Dolag and Schindler 2000; Dolag et al. 2001), the propagation of ultra high energy cosmic-rays (Dolag et al. 2005a) and the distribution of fast radio bursts (Dolag et al. 2015). Donnert et al. (2009), Beck et al. (2013a) presented models for cluster magnetic fields seeded by galaxy feedback, and established that different seeding models can lead to the same cluster magnetic field. Beck et al. (2012) showed theoretical and numerical models for magnetic field seeding and amplification in galactic haloes. We reproduce projected magnetic field strengths in cosmological simulations from three different methods, GADGET (SPH), ENZO (Eulerian finite volume) and AREPO (Lagrangian finite volume) in Fig. 8.

Ryu et al. (2008) established the connection between shock driven vorticity during merger events and magnetic field amplification in clusters using Eulerian cosmological sim-






**Fig. 10** Phase diagrams for different cosmological simulations, like Fig. 1. Top left: RAMSES CT simulation of a cooling-flow galaxy cluster (Dubois and Teyssier 2008; Dubois et al. 2009). Top right: ENZO CT simulation of a major merger cluster (fields injected by AGN activity) (Skillman et al. 2013). Bottom left, right: AREPO (Powell scheme) simulation without and with Illustris galaxy formation model, respectively (Marinacci et al. 2015)

ulations. They applied a semi-analytic model of the small scale dynamo coupled to the turbulent energy to derive  $\mu\text{G}$  fields in clusters (see also Beresnyak and Miniati 2016).

Xu et al. (2009, 2011) used AGN seeding in the first direct Eulerian MHD cluster simulations to obtain magnetic field strengths of 1–2  $\mu\text{G}$  in clusters with a second order TVD method and constrained transport (Li et al. 2008). We reproduce power spectra from this simulation in Fig. 9, left. Considering cosmological seed fields, Vazza et al. (2014) used large uniform grids to simulate magnetic field amplification in a massive cluster. Ruszkowski et al. (2011) presented a simulation of cluster magnetic fields with anisotropic thermal conduction. They find that conduction eliminates the radial bias in turbulent velocity and magnetic fields that they observe without conduction.

Within the limit of available numerical approaches, modern simulations find that adiabatic compression/rarefaction of magnetic field lines is the dominant mechanism across most of the cosmic volume (see Fig. 10), with increasing departures at high density,  $\rho \geq 10^2 \langle \rho \rangle$ , when dynamo amplification sets in. Additional scatter in this relation is also found in presence of additional sources of magnetization or dynamo amplification, such as e.g. feedback from AGN, as shown by the comparison between non-radiative and “full physics” runs. Using a Lagrangian finite volume method, Marinacci et al. (2015, 2018a,b) showed magnetic



**Fig. 11** Map of projected mean magnetic field strength for re-simulations of a cluster with increasing resolution, for regions of  $8.1 \times 8.1 \text{ Mpc}^2$  around the cluster center at  $z = 0$ . Each panel shows the mass-weighted magnetic field strength (in units of  $\log_{10}[\mu\text{G}]$  for a slice of  $\approx 250 \text{ kpc}$  along the line of sight). Adapted from Vazza et al. (2018)

field seeding and evolution with the “Illustris” subgrid model for galaxy formation, also including explicit diffusivity. They obtained  $\mu\text{G}$  magnetic fields in clusters when they included seeding from galaxy feedback (Fig. 10, bottom).

Recently, Vazza et al. (2018) simulated the growth of magnetic field as low as  $0.03 \text{ nG}$  up to  $\sim 1\text{--}2 \mu\text{G}$  using AMR with a piece-wise linear finite volume method. By increasing the maximum spatial resolution in a simulated  $\sim 10^{15} M_{\odot}$  cluster, they observed the onset of significant small-scale dynamo for resolutions  $\leq 16 \text{ kpc}$ , with near-equipartition magnetic fields on  $\leq 100 \text{ kpc}$  scales for the best resolved run ( $\approx 4 \text{ kpc}$ ), see Fig. 11. They estimated that  $\sim 4\%$  turbulent kinetic energy was converted into magnetic energy. The amplified 3D fields show clear spectral, topological and dynamical signatures of the small-scale dynamo in action, with mock Faraday Rotation roughly in-line with observations of the Coma cluster (Bonafede et al. 2013). A significant non-Gaussian distribution of field components is consistently found in the final cluster, resulting from the superposition of different amplification patches mixing in the ICM.

### 4.3 Cosmological Simulations of Magnetic Fields Outside of Galaxy Clusters


The peripheral regions of simulated galaxy clusters mark the abrupt transition from supersonic to subsonic accretion flows, and the onset of the virialization process of the infalling gas. The accreted gas moves supersonically with respect to the warm-hot intergalactic medium in the cluster periphery, which triggers  $\mathcal{M} \sim 10\text{--}100$  strong shocks in the outer regions of clusters and in the filaments attached to them (e.g. Ryu et al. 2003;

Pfrommer et al. 2006). Downstream of such strong shocks, supersonic turbulence is injected towards structures, together with a first inject of vorticity by oblique shocks (e.g. Kang et al. 2007; Ryu et al. 2008; Wittor et al. 2017b). In these physical conditions, the SSD is predicted to be less efficient, because of the predominance of compressive forcing of turbulent motions (Ryu et al. 2008; Federrath et al. 2011a; Jones et al. 2011; Schleicher et al. 2013; Porter et al. 2015). In this case, the maximum magnetic field arising from SSD amplification in the  $10^5 \text{ K} \leq T \leq 10^7 \text{ K}$  medium of filaments would be  $\sim 0.01\text{--}0.1 \mu\text{G}$  (e.g. Ryu et al. 2008; Vazza et al. 2014). Direct numerical simulations investigated the small-scale dynamo amplification of primordial fields in cosmic filaments, so far reporting no evidence for dynamo amplification, unlike for galaxy clusters simulated with the same method and at a similar level of spatial detail (Vazza et al. 2014). This trend is explained by the observed predominance of compressive turbulence at all resolutions (unlike in clusters, where turbulence gets increasingly solenoidal as resolution is increased), as well as by the limited amount of turnover times that infalling gas experiences before being accreted onto clusters (e.g. Ryu et al. 2008; Vazza et al. 2014). If these results will be confirmed by simulations with even larger resolutions, it has the important implication that the present-day magnetization of filaments should be anchored to the seeding events of cosmic magnetic fields, posing a strong case for future radio observations (e.g. Gheller et al. 2016; Vazza et al. 2017a). In this scenario the outer regions of galaxy clusters and filaments are expected to retain information also on the *topology* of initial seed fields even today, as shown in numerical simulations at high resolution (e.g. Brüggem et al. 2005b; Marinacci et al. 2015, see also Fig. 12), in case the magnetic fields have a primordial origin. Conversely, if the fields we observe in galaxy clusters are mostly the result of seeding from active galactic nuclei and galactic activities, the magnetization at the scale of filaments and cluster outskirts is predicted to be low (e.g. Donnert et al. 2009; Xu et al. 2009; Marinacci et al. 2015). Future surveys in polarization should have the sensitivity to investigate the outer regions of galaxy clusters down to  $\sim 1\text{--}10 \text{ rad/m}^2$  (e.g. Taylor et al. 2015; Bonafede et al. 2015; Vacca et al. 2016), which is enough to discriminate among most extreme alternatives in cluster outskirts (e.g. Vazza et al. 2017a).

#### 4.4 Discussion

Simulations of magnetic field amplification in clusters have reproduced key observations for two decades now. Most of the early progress has been achieved with Lagrangian methods originally developed in the galaxy formation context, most notably SPH (Dolag et al. 2002). These simulations reproduce the magnetic field strength inferred from rotation measures in clusters and have been used extensively to model related astrophysical questions. However, the adaptivity of Lagrangian methods and the particle noise in SPH limits their ability to resolve the structure of the magnetic field, especially in low density environments (cluster outskirts, filaments).

Clear theoretical expectations for the small scale dynamo in clusters have been established (Ryu et al. 2008; Beresnyak and Miniati 2016), also from idealized simulations (e.g. Schekochihin et al. 2004; Cho et al. 2009; Porter et al. 2015). Some of these expectations have been tested in cosmological simulations using Eulerian codes (e.g. Vazza et al. 2018). Recent Eulerian simulations approach observed field strengths in clusters, but do not reach field strengths obtained from Lagrangian approaches. Beresnyak and Miniati (2016) argued that due to numerical diffusion, Eulerian approaches spend too much time in the exponential/kinetic growth phase, thus the non-linear growth phase is severely truncated. Following



**Fig. 12** Left: magnetic field vectors for a cosmic filaments simulated with AMR using FLASH (Brüggen et al. 2005b). Right: magnetic field vectors around a massive galaxy cluster (top) and a filament (bottom) simulated with AREPO (Marinacci et al. 2015) and for two different topologies of uniform seed magnetic fields

Schekochihin et al. (2004), a clear indicator for the presence of a dynamo that cannot be produced via compression is the anti-correlation of magnetic field strength and its curvature  $\mathbf{K}$ :

$$\mathbf{K} = \frac{(\mathbf{B} \cdot \nabla)\mathbf{B}}{\mathbf{B}^2}, \tag{20}$$

so that  $\mathbf{BK}^{\frac{1}{2}} = \text{const}$ , where the exponent has to be obtained from the magnetic field distribution. In cluster simulations, only Vazza et al. (2018) have demonstrated consistent curvature correlations. We note that in galactic dynamos, consistent results have recently been achieved with Eulerian and Lagrangian codes (Butsky et al. 2017; Rieder and Teyssier 2016; Pakmor et al. 2017; Steinwandel et al. 2018), but only Steinwandel et al. (2018) showed a curvature relation.

In clusters, all simulations show an exponential increase in magnetic field strength followed by a non-linear growth phase (e.g. Beck et al. 2012). However, the timescale of exponential growth is set by the velocity power/rate of strain at the resolution scale, which in turn is determined by the MHD algorithm (resolution, dissipation/noise). The real kinematic dynamo in primordial haloes is far below the resolution scale of every numerical scheme (Beresnyak and Miniati 2016) and needs to be treated with an large eddy approach (Yakhot and Sreenivasan 2005; Cho et al. 2009).

As we have motivated above, dynamo theory predicts that the final structure of cluster magnetic fields is shaped by turbulence near the Alfvén scale, because this is where the eddy turnover time is smallest (Eq. (11), a few kpc in a massive cluster merger). Thus an accurate simulation of field topology has to faithfully follow the velocity field and the magnetic field near this scale in the non-linear growth phase, i.e. at least achieve Reynolds numbers (Eq. (5)) of 300–500 at redshifts  $z < 1$  during a major merger (Haugen et al. 2004; Beresnyak and Miniati 2016). For an outer scale of 300 kpc, this implies evolution of turbulence velocity and magnetic field growth at about 1 kpc, including numeric effects.

This makes the small scale dynamo in clusters is a very hard problem, because it combines the large dynamical range of scales in cosmological clustering with the evolution of

two coupled vector fields (turbulence and magnetic fields) near the resolution scale. Additionally, seeding on smaller scales by galactic outflows may play an important role. Hence, it is likely the numerical dissipation scale that shapes the outcome of MHD simulations in a cosmological context. We now provide a short discussion of effective Reynolds numbers and numerical limitations in current approaches.

#### 4.4.1 Effective Reynolds Numbers

From the numerical viewpoint, the Reynolds number of a flow increases with the effective dynamic range reached inside a given volume. Its upper limit is set by the driving scale and the spatial resolution in the volume of interest following Eq. (5). However, in any numerical scheme the effective dynamic range and Reynolds number of the flow are reduced by the cut-off of velocity and magnetic field power near the numerical dissipation scale in Fourier space (e.g. Dobler et al. 2003). Simply put, numerical error takes away velocity and magnetic field power close the resolution scale in most schemes. The shape of the velocity power spectrum on small scales determines how much velocity power (rate of strain  $\delta u/l$ , see Sect. 3.3) is available to fold the magnetic field and drive the small-scale dynamo. Thus a less diffusive (finite volume) code reaches higher effective Reynolds numbers, faster amplification and a more tangled field structure at the same resolution.

We can quantify this behavior by introducing an *effective Reynolds number* of an MHD simulation of turbulence as:


$$R_{e,\min} \approx \left( \frac{L}{\varepsilon \Delta x} \right)^{4/3}, \quad (21)$$

where  $\Delta x$  is the resolution element,  $\varepsilon$  is a factor depending on the diffusivity of the numerical method, and  $L$  is the outer scale (in clusters 300–500 kpc, Sect. 3.5). As a conservative estimate, one may assume in modern SPH codes  $\varepsilon \geq 10$  (Price 2012a, Fig. 13), in hybrid codes  $\varepsilon \approx 10$  (Hopkins (2015)). For second order finite difference/volume codes one often assumes  $\varepsilon \approx 7$  (e.g. Kritsuk et al. 2011; Rieder and Teysier 2016). In Fig. 13 left, we reproduce velocity power spectra from a driven compressible turbulence in a box simulation with  $128^3$  zones using the finite volume (FV) code AREPO and the discontinuous Galerkin (DG) code TENET (Bauer et al. 2016). Second order FV is shown in yellow, while second, third and fourth order DG power spectra are shown in green, blue and purple, respectively. The formal resolution/Nyquist scale remains constant in all runs. However, with increasing order of spatial and time interpolation, viscosity reduces, the effective dissipation scale shrinks, velocity power on small scales increases, the inertial range grows in size, and with it the effective Reynolds number of the simulation (i.e.  $\varepsilon$  decreases). Note that the DG scheme has more power near the dissipation scale than the FV scheme, even at the same order (green vs. yellow). This indicates that formal convergence order is not sufficient to determine effective Reynolds numbers at a given resolution.  $\varepsilon$  obviously depends on implementation details and has to be determined empirically with driven turbulence “in a box” simulations. For a recent review on high-order finite-volume schemes, see Balsara (2017).

#### 4.4.2 Dynamos in Eulerian Schemes

In non-adaptive Eulerian cluster simulations the effective Reynolds number is set by the resolution of the grid and the diffusivity of the numerical method (e.g. Kritsuk et al. 2011). Federrath et al. (2011b) and Latif et al. (2013) reported that only by resolving the Jeans





**Fig. 13** Left: Velocity power spectra of driven compressible turbulence with a second order finite volume scheme (yellow) and Discontinuous Galerkin schemes (2nd order: green, 3rd order blue, 4th order purple) (Bauer et al. 2016). Right: Velocity power spectra of decaying turbulence simulated with modern SPH (Beck et al. 2016). The spectra were obtained using wavelet kernel binning to remove aliasing above the kernel scale  $k_{\text{hsm1}}$


length of a halo with  $\geq 64$  cells the small-scale dynamo can develop (e.g.  $R_M \sim 32$  setting  $\varepsilon = 2$  in Eq. (21)) in a proto-galactic halo of  $10^6 M_\odot$  at  $z \sim 10$ . However, Vazza et al. (2014) reported that small-scale amplification can begin before  $z = 0$  in  $\sim 10^{14} M_\odot$  galaxy clusters if their virial diameter is resolved with at least  $\geq 100$  cells ( $R_M \sim 50$ ), while in order to approach energy equipartition between turbulence and magnetic fields by  $z = 0$  one needs to resolve the virial diameter with  $\geq 1500$  elements ( $R_M \sim 750$  in the ideal case). These differences likely arise from the shapes of the numerical dissipative and resistive scales. The underlying Eulerian methods were either second or first order accurate and used CT or Dedner cleaning to constrain magnetic field divergence.

Eulerian structure formation simulations produce flows with supersonic velocities relative to the simulation grid. At the same time, the truncation error of Eulerian methods is inherently velocity dependent (Robertson et al. 2010; Bauer et al. 2016). It has been shown that these errors do not pose a problem for the simulation of clusters in a cosmological context (Mitchell et al. 2009), but they may suppress the growth of instabilities close to the dissipation scale (e.g. Springel 2010) and thus further reduce the effective Reynolds number of the simulation. We note that poorly un-split Eulerian schemes may also affect angular momentum conservation close to the resolution scale and further reduce the accuracy of e.g. galaxy formation simulations, where angular momentum conservation is desirable to produce disc galaxies.

These arguments extend also to magnetic fields, whose advection poses a challenging test for all Eulerian schemes. In Fig. 14, right, we reproduce the time evolution of magnetic energy during the advection of a magnetic field loop in 2D with the ATHENA code at different resolutions (Gardiner and Stone 2008). As the size of the field loop approaches the resolution scale, field energy is diffused more quickly. Again, the diffusivity added by the scheme to keep local magnetic field divergence small varies with implementation and has to be determined by empirical tests. There are sizable differences even among CT schemes, which inherently conserve the divergence constraint to machine precision (see e.g. Lee 2013).

#### 4.4.3 Dynamos in Lagrangian Schemes

In adaptive Lagrangian cluster simulations, the resolution is a function of density and thus varies in space and time during the formation of a cluster or filament. Thus the dissipation



**Fig. 14** Left: Growth of magnetic energy in supersonic driven turbulence simulations in SPMHD (dashed lines) and FV (solid lines) at different resolutions (Tricco et al. 2016). Right: Evolution of magnetic energy in the advection of a magnetic field loop in 2 dimensions from Gardiner and Stone (2008)

scale and the Reynolds number are not well defined in Fourier space and turbulence can be strictly defined only on the coarsest resolution element in a given volume. The effect of the adaptivity on the dynamo and especially the resulting field structure is not entirely clear. It seems reasonable to assume additional (magnetic) dissipation, if a magnetized gas parcel moves to a less-dense environment and is adiabatically expanded and divergence cleaned. While turbulent driving is correlated with over-densities in a cosmological context, the turbulent cascade is not. Thus density adaptivity, which is a very powerful approach in galaxy formation simulations, might introduce a density bias to the magnetic field distribution in strongly stratified media. The growth rate of the turbulent dynamo depends on the eddy turnover time, which is smallest in highly resolved regions. Thus Lagrangian schemes might grow magnetic fields faster in high density regions (cluster cores) than in low density regions (cluster outskirts). However, it remains unclear how strongly current results are affected by this issue, simply because no Eulerian simulation with kpc resolution in the cluster outskirts is available.

In cosmological simulations, Dolag et al. (1999, 2002) reported sizeable cluster magnetic fields even with a traditional SPH algorithm and comparably low resolution. As we have shown, theory provides clear predictions for the evolution of a magnetic field in a turbulent dynamo, which have been successfully verified with Eulerian methods. For some Lagrangian methods (e.g. Pakmor et al. 2011), it is reasonable to assume that at fixed resolution the result will be similar to the established dynamo theory, simply because their dissipation scale defaults to a finite volume method. For other new hybrid methods (Hopkins and Raives 2016) the situation is less clear. In general, the idealized magnetic dynamo in Lagrangian schemes is not well researched yet and we would encourage the community to close this gap.

For traditional SPH algorithms, its ability to accurately model hydrodynamic turbulence was heavily debated (Bauer and Springel 2012; Price 2012a). We note that computing a grid representation from an irregularly sampled vector field to obtain a power spectrum is a diffusive process and prone to aliasing (Beck et al. 2016). Modern SPH schemes have improved significantly, and it has been shown that sub-kernel re-meshing motions are required to maintain sampling accuracy (Price 2012b). The influence of these motions on the magnetic dynamo are not well understood, especially in the subsonic regime that is dominant in clusters.



In the supersonic regime, Tricco et al. (2016) compared simulations with  $M = 10$  using the SPMHD code PHANTOM and the finite volume code FLASH with an HLL3R solver (Waagan et al. 2011) and both with Dedner cleaning. They found that the growth of magnetic energy in the SPMHD dynamo speeds up with increasing resolution. In contrast, the finite volume scheme converged (Fig. 14, left). They found Prandtl numbers of  $P_r = 2$  and  $P_r < 1$ , respectively. They argued that the growth in the SPMHD dynamo is due to the artificial viscosity and resistivity employed, which is negligibly small in the absence of shocks.

We note that these results cannot be simply transferred to galaxy cluster simulations. As mentioned before, cluster turbulence is largely sub-sonic, super-Alfvénic and solenoidal, thus shocks do not play a role for the dissipation of turbulent energy. Cosmological codes usually do not include explicit dissipation terms, in contrast diffusivity is usually minimized. Driven subsonic turbulence simulations with SPMHD are required to characterize the sub-sonic SPMHD dynamo in clusters and clarify the role particle noise could play even in early SPMHD cluster simulations. We note that some numerical amplification has been reported in SPMHD simulations of the galactic dynamo (Staszczyn and Elstner 2015; Dobbs et al. 2016).

## 5 Magnetic Field Amplification at Shocks


Shocks amplify magnetic fields by a number of mechanisms, not all of which are well understood (Brüggen et al. 2012). Compression at the shock interface leads to the amplification of the quasi-perpendicular part of the upstream magnetic field. Compressional amplification has the allure of explaining the large degrees of polarization in radio relics, but suffers from the limitation of small amplification factors. For amplification by pure compression, (Iapichino and Brüggen 2012) find for the ratio of magnetic fields:

$$\frac{B_{\text{dw}}}{B_{\text{uw}}} = \sqrt{\frac{2\sigma^2 + 1}{3}}, \quad (22)$$

with the shock compression ratio  $\sigma$ . Thus, for typical shock strengths in cluster mergers, ( $M \approx 2-3$ ), the amplification factor is limited to around 2.5, which results in inconsistencies of the minimum magnetic field strengths inferred in some radio relics with global magnetic field scalings (Donnert et al. 2017). Similar expressions have been found for SNR (Reynolds 1998).

### 5.1 Shock-Driven Dynamo

Downstream of shocks, magnetic fields can be amplified by a small-scale dynamo that is driven by turbulence created at the shock front (Binney 1974). This has been observed in supernova remnants (SNR) (Parizot et al. 2006). This turbulence could be driven by the baroclinic vorticity that is generated for example by upstream inhomogeneities in gas density. For parameters relevant in SNR, Giacalone and Jokipii (2007) have demonstrated in MHD simulations that density inhomogeneities in the pre-shock fluid cause turbulence and magnetic field amplification in the post-shock fluid. Simulations by Inoue et al. (2009) showed that the maximum amplification is set by the plasma beta parameter. Sano et al. (2012) argued that turbulence is injected by Richtmyer-Meshkov instabilities. Fraschetti (2013) derived an analytical approach for 2D SNR shocks. Guo et al. (2012) studied the interaction of a SNR shock propagating into a turbulent medium upstream. However, the relevant parameters in



**Fig. 15** Left: Magnetic field amplification over Alfvénic Mach number in 2D MHD shock simulations from Ji et al. (2016). Right: Models for magnetic field evolution inferred in the Sausage relic from Donnert et al. (2016)

the shock and the upstream medium in SNR blast waves differ significantly from galaxy cluster shocks. In clusters, Mach numbers are lower ( $< 5$ ) and the plasma beta parameter is larger ( $\beta_{\text{pl}} \geq 100$ ). It is unclear if these results from SNR carry over to the ICM.


Literature on turbulent magnetic field amplification in ICM shocks remains scarce. Iapichino and Brüggén (2012) studied the evolution of vorticity behind the shock. They argue that self-generated vorticity from the shock is not sufficient to drive a turbulent dynamo downstream, but that about 30% of turbulent pressure is required upstream of the shock to explain observed magnetic field lower limits.

Ji et al. (2016) studied magnetic field amplification in idealized MHD simulations of shocks. They found that amplification is independent of plasma beta for Mach numbers of a few, but is linearly dependent on the Alfvénic Mach number in shocks. In Fig. 15, left, we show their results for 2D simulations at different resolutions, with the highest resolution in magenta. Below  $M_A \approx 10$ , compression dominates the amplification and results in magnetic field structures perpendicular to the shock normal. Above  $M_A \approx 10$ , turbulence injected by the shock amplifies magnetic fields to strengths significantly higher than expected by compression. In this limit, the field topology becomes mostly quasi-parallel, because velocity shear is largest in the direction of shock propagation.

Along these lines, in Wittor et al. (2017a), it has been found that the stretching motions dominate the evolution of turbulence in galaxy clusters. However, baroclinic motions are needed to generate turbulence. The enstrophy dissipation rate peaks when the enstrophy is maximal and this is the time when magnetic field amplification by a small-scale dynamo would be the strongest.

These results have important implications for radio relics. In most relics the lower limit for the downstream magnetic field is found to be around 1–3  $\mu\text{G}$  (e.g. Finoguenov et al. 2010). This is consistent with equipartition magnetic field strengths of 4–7  $\mu\text{G}$  (e.g. Nakazawa et al. 2009; Finoguenov et al. 2010; Stroe et al. 2014).

The ordered topology of magnetic fields expected by compressional amplification can explain the large degree of polarization found in some radio relics, thus dis-favouring turbulent amplification. However, given typical Mach numbers (2–3), the lower limits on magnetic field strengths in relics imply upstream fields of about 1–2  $\mu\text{G}$  ahead of some shocks (Eq. (22)). As relics reside in the outskirts of clusters, this is difficult to explain with the common scaling of magnetic field strength with density/radius in the ICM (Bonafede et al. 2010).




**Fig. 16** Amplified magnetic fields produced by a remnant-core cold front in an MHD simulation detailed in Asai et al. (2007). The panels show slices of temperature (left), temperature gradient (middle), and magnetic field strength (right). An amplified magnetic field appears in a draping layer around the cold front

However, a recent model (Donnert et al. 2016) of the Sausage relic motivates Alfvénic Mach numbers of around 100 in the shock and showed that exponential downstream field amplification (Fig. 15, right) can explain the steepening in the radio spectrum above 8 GHz found in the Sausage (Stroe et al. 2013). More discussion will also be found in van Weeren et al. (this volume).

## 6 Magnetic Field Amplification from Cold Front Motions

Aside from turbulence and shocks, many galaxy clusters also possess subsonic bulk flows which can amplify magnetic fields in localized regions. The first evidence of these motions was provided shortly after the launch of the *Chandra* X-ray Observatory. *Chandra*'s sub-arcsecond spatial resolution revealed the presence of surface brightness edges in many clusters. Through spectroscopic analysis most of these edges, which superficially appear as shocks, were identified to be contact discontinuities, where the denser (brighter) side of the edge is colder than the lighter (dimmer) side. These features have been dubbed “cold fronts”, and are believed to be the result of subsonic gas motions driven by cluster mergers and cosmic accretion (for recent reviews see Markevitch and Vikhlinin 2007; Zuhone and Roediger 2016). Cold fronts have been described as forming via at least three processes: “remnant-core” fronts are formed by cool cores of sub-clusters or galaxies falling into or merging with larger, more diffuse structures, “sloshing” cold fronts which are formed in cool-core clusters by the displacement of the central low-entropy gas of the DM-dominated core, and “stream” cold fronts which are formed by collisions between coherent streams of gas (Birnbom et al. 2010; Zuhone and Roediger 2016; Zinger et al. 2018).

The relevance of such bulk motions for the amplification of the cluster magnetic field was first shown by Lyutikov (2006). They demonstrated that the subsonic motion of a dense gas cloud through the ICM would amplify and stretch magnetic fields, regardless of the initial geometry, along the contact discontinuity that forms, producing a thin “magnetic draping layer”. The only condition is that the Alfvénic Mach number  $\mathcal{M}_A > 1$ , a condition readily satisfied in the ICM. The width of the layers is given roughly by  $\Delta r \sim L / \mathcal{M}_A^2$ . For typical conditions in the ICM and a mildly subsonic cloud with  $\mathcal{M} \lesssim 1.0$ ,  $\Delta r \sim 0.01L$ . Lyutikov (2006) also pointed out that such layers should be associated with a depletion of plasma. This is so because the total pressure should remain continuous within and around such a layer, given the subsonic motion of the gas, and thus an increase in magnetic pressure requires a decrease in thermal pressure. This indicates that such layers may be visible in X-ray observations of cold fronts, though in practice there will be large uncertainties given projection effects.



**Fig. 17** Amplified magnetic fields produced by sloshing cold fronts in an MHD simulation. Left panel: Slice through the gas temperature in keV, showing the spiral-shaped cold fronts. Right panel: Slice through the magnetic field strength, showing amplified fields within the cold fronts. Figure reproduced from Zuhone and Roediger (2016)


The first numerical simulations used to examine this effect followed the evolution of a cold, dense core moving subsonically through a hot, magnetized ICM (Dursi 2007; Dursi and Pfrommer 2008; Asai et al. 2004, 2005, 2007), using a variety of field geometries. In all cases they confirmed the basic picture offered by Lyutikov (2006) of magnetic field amplification in a thin layer “draping” the cold front surface which forms at the head of the cool core. Figure 16 shows an example MHD simulation of a remnant-core cold front producing an amplified magnetic field in a draping layer from Asai et al. (2007).

A second type of cold front, the “sloshing” variety (Fig. 17), occurs in more relaxed systems when the cold gas core is perturbed by infalling subclusters and is separated from the DM-dominated potential well. This gas then oscillates back and forth in the cluster center, producing a spiral-shaped pattern. Simulations have shown that sloshing cold fronts are also associated with amplified magnetic fields. The first simulations to demonstrate this were those of Zuhone et al. (2011), who simulated the evolution of initially tangled magnetic fields with a number of initial magnetic field strengths and correlation scales. The sloshing cold fronts are also associated with amplified magnetic layers, but unlike in the scenario envisaged by Lyutikov (2006) the layers are on the *inside* of the front surface rather than outside, due to the fact that for the sloshing cold fronts the shear flow is predominantly inside. This results in increased magnetic fields within the volume bounded by the cold fronts (Fig. 17), an effect important for the generation of radio mini-halos (Sect. 6.2).

## 6.1 Effects of Cold Front Magnetic Fields on the Thermal Plasma

The above considerations indicate that if a highly magnetized layer forms tangential to a cold front or otherwise because of shearing motions that it may produce a dip in X-ray surface brightness at this location. These dips were first noticed in MHD simulations of sloshing cold fronts by Zuhone et al. (2011). In these simulations, the layers reached magnetic field strengths with  $\beta_{pl} \sim 10$  and dips in density and temperature of roughly  $\sim 10$ – $30\%$ , which could produce dips in surface brightness of roughly  $\sim 5$ – $10\%$ , depending on the gas temperature. The evidence for such features in X-ray observations of clusters is so far inconclusive, but there are some tantalizing hints (e.g. Werner et al. 2016) (Fig. 18).

The magnetic tension from a field stretched parallel to a front surface will suppress the growth of K-H instabilities if the field is strong enough (Chandrasekhar 1961). The initial smooth appearance of many cold fronts as seen in *Chandra* observations led readily



**Fig. 18** Evidence for amplified magnetic fields in X-ray observations of the Virgo cluster from Werner et al. (2016). Left panel: residual image of the X-ray surface brightness as seen by *Chandra* near the cold front. Three linear enhancements in surface brightness are apparent. Middle and right panels: Slices of gas density and magnetic field strength near the cold front surface in a MHD simulation of the Virgo cluster. Wide bands of strong magnetic field in between narrow channels of weak field produce linear features in density similar to those seen in Virgo

to the proposal that such suppression was occurring. For example, the apparent smoothness of the merger-remnant cold front in A3667 led Vikhlinin et al. (2001), Vikhlinin and Markevitch (2002) to estimate a magnetic field strength near the front surface between  $6 \mu\text{G} < B < 14 \mu\text{G}$ . More recently, Chen et al. (2017) estimated a magnetic field strength of  $B \sim 20\text{--}30 \mu\text{G}$  at the sloshing cold fronts in A2204 based on the lack of observed KHI.


As deeper *Chandra* observations of nearby clusters with longer exposures have been obtained over the years, some evidence for K-H instabilities has been uncovered. Roediger et al. (2015a,b), Kraft et al. (2017) presented evidence of gas stripping of caused by KHI in the elliptical galaxy M89 using deep X-ray observations and tailored simulations. Ichinohe et al. (2017) showed evidence for KHI in a longer combined exposure of A3667 than was available to Vikhlinin et al. (2001), Vikhlinin and Markevitch (2002), but did not make an updated estimate of the magnetic field strength. Finally, Su et al. (2017) showed evidence for K-H instabilities at the interface of the cold front in NGC 1404, and used their presence to place an upper limit on the magnetic field strength at the front of  $5 \mu\text{G}$ .

However, the presence of *some* degree of KHI in cold fronts is not inconsistent with the picture of magnetic draping layers per se—it is rather likely an indication of the strength of the fields in these layers. A recent series of papers has constrained the magnetic field strength in the Perseus cluster using MHD simulations and X-ray observations. Walker et al. (2017) showed convincing evidence of a giant KHI eddy at one of the cold front edges in the Perseus cluster. They compared the appearance of the cold fronts to the simulations from ZuHone et al. (2011), and suggested that a cluster with an initial  $\beta \sim 200$  before the sloshing began could explain the presence of the KHI eddy—simulations with initially larger or smaller average magnetic field strengths produced results that were inconsistent in terms of having either too few or too many KHI eddies along the interface.

## 6.2 Amplified Magnetic Fields and Cosmic Rays: Radio Mini-Halos

Radio mini-halos are the smaller-scale siblings of the giant radio halos, hosted in cool-core clusters. Their emission is similarly diffuse and has a steep spectrum ( $\alpha \sim 1\text{--}2$ ), but are nearly an order of magnitude smaller than radio halos and are confined to the core region. Mazzotta and Giacintucci (2008) were the first to discover that the radio mini-halos in the clusters RX J1720.1+2638 and MS 1455.0+2232 were confined to the region on the sky





**Fig. 19** Association of radio mini-halos with cold fronts from ZuHone et al. (2013). Left panel: *Chandra* observation of RXJ 1720.1+26, showing sloshing cold fronts, with 610 MHz radio contours overlaid. The radio emission is coincident with the sloshing cold fronts. Middle panel: Simulated 327 MHz radio contours overlaid on projected temperature from an MHD/CRE simulation of sloshing core gas which produces a mini-halo with emission bounded by the cold fronts. Right panel: Profiles of temperature and radio emission taken along the two directions shown in the middle panel, which show clearly that the radio emission drops steeply at the position of the cold fronts

bounded by sloshing cold fronts seen in the X-ray observations. Subsequent investigations of mini-halo emission from a number of cool-core clusters have confirmed the existence of sharp drops in radio emission at the position of the cold front surfaces in many cases (Giacintucci et al. 2014a,b, 2017).

Such radio emission requires a population of CRE with  $\gamma \sim 10^3\text{--}10^4$ , given the typical magnetic field strengths in clusters. Since CRE with such energies cool rapidly via synchrotron and Inverse-Compton losses, the existence of mini-halos requires a mechanism to replenish these electrons, either by reacceleration from a lower-energy population (the “reacceleration” model Brunetti and Lazarian 2007, 2011a,b) or as the byproducts of collisions of CRp with the ICM thermal proton population (the “hadronic” model, Dennison 1980; Pfrommer and EnBlin 2004; Keshet and Loeb 2010), though this model is strongly constrained by the Fermi-LAT upper limits on gamma-ray emission in clusters (Ackermann et al. 2014), which are also produced by the same collisions. A review of these processes and their implications for non-thermal emission in clusters can be found in Brunetti and Jones (2014).

As previously noted, sloshing cold fronts are very common in cool-core clusters, and these motions amplify magnetic fields. A stronger magnetic field within the core would lead to an enhancement of the mini-halo emission. Since this amplified magnetic field is largely confined to the volume bounded by the cold fronts, it may also explain the association of mini-halos with cold fronts and the steep drops in radio emission coincident with the front surfaces, as suggested by Keshet and Loeb (2010).

The only simulations so far to directly test the reacceleration scenario for mini-halos were performed by ZuHone et al. (2013). They used a MHD simulation of gas sloshing in a cool-core cluster coupled with a simulation of the evolution of the CRE spectrum under reacceleration by turbulence and radiative and Coulomb losses along trajectories of passive tracer particles. It was found that reacceleration by turbulence coupled with the magnetic field amplification, both produced by the sloshing motions, could produce mini-halos which have the characteristic diffuse emission, steep spectrum, and spatial relationship to the cold fronts (Fig. 19). They noted that the mini-halo produced in their simulation had two further interesting characteristics: the emission was transient and brightest shortly after the beginning of the sloshing motions, and had a “patchy” appearance due to the intermittent and irregular distribution of turbulent gas motions in the core region. The latter prediction

is perhaps supported by the recent JVLA 230–470 MHz observations of the Perseus mini-halo by Gendron-Marsolais et al. (2017), which revealed a complex appearance of the radio emission.

The coincidence of mini-halos with cold fronts could also be explained by amplified magnetic fields in the hadronic model. Keshet and Loeb (2010) also suggested that these rapidly amplifying magnetic fields may be responsible for the steep spectrum of mini-halos. ZuHone et al. (2015a) tested this possibility using simulations that the fast amplification of magnetic fields by sloshing motions could produce diffuse, core-confined mini-halos with steep spectra by steepening the CRe spectrum. Though the amplification of the magnetic field strength within the sloshing cold fronts reproduced the observed spatial properties of mini-halo emission in this simulation, they found that only a small, observationally insignificant number of tracer particle trajectories experienced sufficiently rapid changes in magnetic field strength to steepen their radio spectra to  $\alpha \sim 2$ , and then only for brief periods of time. More complex mini-halo morphologies and spectral index properties may be produced in the context of hadronic models by taking into account other CRp physics such as diffusion, streaming, and advection by turbulence and/or bulk motions (Enßlin et al. 2011; Pfrommer 2013; Wiener et al. 2013, 2018; Jacob and Pfrommer 2017a,b).

## 7 Concluding Remarks

We have presented the most important mechanisms that are expected to control and drive the amplification of magnetic fields observed in galaxy clusters at radio wavelengths. We gave a short introduction to turbulence, motivated MHD as a model for the intra-cluster-medium and introduced the basic principles of the small-scale MHD turbulent dynamo. We reviewed the outcome of (cosmological) numerical simulations of the growth of magnetic fields, under typical conditions in the intracluster medium. We provided a short discussion of numerical limitations of current approaches and established the demands of upcoming radio surveys of Faraday rotation measures and giant radio haloes. We also introduced magnetic field amplification at shocks and by cosmic-rays, which is evidenced by radio relics at cluster outskirts. Further we discussed magnetic field amplification by cold fronts and internal cluster motions, which are likely connected to radio mini halos.

The observed properties of magnetic fields in the intra-cluster medium require large amplification factors ( $\geq 10^3$ ) even considering the effect of gas compression. Indeed, there is plenty of time and turbulent energy to boost the magnetic field energy up to observed values with a turbulent small-scale dynamo. The theoretical grounds of this small-scale dynamo model for the amplification of weak fields in random flows are robust and are covered in a significant amount of literature. A central outcome of these studies is the importance of the Alfvén scale on the amplification and topology of magnetic fields in turbulent flows.

Given these expectations, the quantitative outcome of the small-scale dynamo in current cosmological simulations is likely not sufficient to robustly predict magnetic fields in the ICM for upcoming radio interferometers. Modern instruments require the robust prediction of magnetic field and shock structures down to a few kpc in the whole volume of a massive galaxy cluster. This means future cosmological simulations will need to resolve the small-scale dynamo down to at least similar resolutions.

A way forward may be higher order MHD methods suitable for cosmology, that resolve motions closer to the grid scale. Another possibility, especially for shocks, could be adaptive techniques to selectively refine the mesh, where the amplification is active. We note that



**Table 1** Reference parameters for the amplification of magnetic fields in various locations of the intra-cluster-medium. 2nd column: typical observed value; 3rd column: plasma beta; 4th column: inferred (minimum) amplification factor of magnetic field energy (also including gas compression); 5th column: amplification factor only considering dynamo amplification; 6th column: minimum amplification time; 7th column: typical factor of field reversals; 8th column: estimated growth factor in magnetic energy

	$B_{\text{obs}}$ [ $\mu\text{G}$ ]	$\beta_{\text{pl}}$	$(\delta B/B_0)_{\text{tot}}^2$	$(\delta B/B_0)_{\text{dyn}}^2$	$t_{\text{dyn}}$ [Gyr]	$l_{\text{B}}$ [kpc]	$\Gamma$ [ $\text{Gyr}^{-1}$ ]
SSD	$\sim 0.1\text{--}5$	$\sim 10^2$	$\geq 10^8$	$\geq 10^3$	$\leq 10$	$\leq 50\text{--}100$	$\sim 1$
Shocks	$\sim 1\text{--}3$	$\sim 10$	$\geq 100$	$\geq 30$	$\leq 0.01$	$\leq 10\text{--}10^2$	$\sim 0.003$
Sloshing	$\sim 10\text{--}20$	$\sim 1$	$\geq 10$	$\geq 10$	$\leq 0.01$	$\leq 1\text{--}10$	$\sim 0.001$

ample computing power is available, as the largest computers approach  $10^{18}$  floating point operations per second in the next years.

Given the large magnetic Reynolds number in the real intra-cluster medium, observed  $\sim \mu\text{G}$  fields can be the result of either primordial  $\sim \text{nG}$  (co-moving) or  $\sim 10^{-6}$  nG fields, as well as from higher and more concentrated magnetic seeds released by galactic winds or active galactic nuclei. Density adaptive techniques are well suited for these kinds of simulations. However, as opposed to galaxy formation, turbulence evolution and the SSD are only weakly correlated with density peaks in the large-scale structure. Thus density adaptive techniques might turn out to be inefficient for this problem, e.g. resolving shocks with a few kpc at the virial radius of a cluster would require exceedingly large particle numbers.

The inferred amplification factors in clusters shocks traced by radio relics and at the contact discontinuities generated by bulk motions in cluster cores are lower than from the SSD ( $\sim 10\text{--}30$ ), yet amplification must again operate on small scales ( $\leq 10^2$  kpc) and on short timescales ( $\sim 10$  Myr). Fully reproducing these trends with simulations may still be a challenge for numerical simulations, again because of resolution. Additionally, more complex interplay of magnetic fields and cosmic-rays are likely at work and plasma conditions across such discontinuities in the fluid may require a consideration of modified viscosity and thermal conduction.

In Table 1 we summarize these general trends in magnetic field amplification found from observations and guided by simulations, which we consider to be robust against numerical issues. We give the typically observed magnetic field, the estimated total magnetic energy growth (considering for each case the magnetic field before the process begins; the magnetic energy growth only due to small-scale dynamo (e.g. after removing for the  $\propto n^{4/3}$  compression factor related to each process); the estimated (minimum) time for amplification; the typical energy containing scale related to each mechanism. The final column gives the estimated growth factor,  $\gamma_{\text{growth}}$  for each amplification mechanism, parameterized by  $(\delta B/B_0)_{\text{dyn}}^2 \approx \exp(\Gamma t_{\text{dyn}})$ .

In this review, we were addressing the successes and limitations of numerical models for predictions of extragalactic magnetic fields. It seems obvious to foresee that the upcoming generation of radio observations (culminating in the SKA data) will pose ever more challenging questions to our theoretical and numerical models of rarefied space plasmas. Only in synergy will numerical techniques and radio observations exploit the next generation of radio telescopes to study plasma physics in the Universe.

**Acknowledgements** The authors would like to thank the referee, D.C. Ellison, for valuable comments that improved the review. The authors thank K. Dolag for discussions on galactic dynamos. JD thanks T. Tricco,

D. Price, D. Ryu and T. Jones for discussions on numerical MHD methods. JAZ thanks C. Pfrommer for useful comments.

F.V. acknowledges financial support from the ERC Starting Grant “MAGCOW”, no. 714196, and the usage of computational resources on the Piz-Daint supercluster at CSCS-ETHZ (Lugano, Switzerland) under project s701 and s805, and of the JURECA supercluster at JFZ (Jülich, Germany) under project hhh42 (HiMAG). JD thanks H. Roettgering and Leiden observatory for the hospitality. This research has received funding from the People Programme (Marie Skłodowska Curie Actions) of the European Unions Eighth Framework Programme H2020 under REA grant agreement no. 658912, “CosmoPlasmas”.

**Open Access** This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made.

## References

- M. Ackermann, M. Ajello, A. Albert, A. Allafort, W.B. Atwood, L. Baldini, J. Ballet, G. Barbiellini, D. Bastieri, K. Bechtol, R. Bellazzini, E.D. Bloom, E. Bonamente, E. Bottacini, T.J. Brandt, J. Bregeon, M. Brigida, P. Bruel, R. Buehler, S. Buson, G.A. Caliandro, R.A. Cameron, P.A. Caraveo, E. Cavazzuti, R.C.G. Chaves, J. Chiang, G. Chiaro, S. Ciprini, R. Claus, J. Cohen-Tanugi, J. Conrad, F. D’Ammando, A. de Angelis, F. de Palma, C.D. Dermer, S.W. Digel, P.S. Drell, A. Drlica-Wagner, C. Favuzzi, A. Franckowiak, S. Funk, P. Fusco, F. Gargano, D. Gasparrini, S. Germani, N. Giglietto, F. Giordano, M. Giroletti, G. Godfrey, G.A. Gomez-Vargas, I.A. Grenier, S. Guiriec, M. Gustafsson, D. Hadasch, M. Hayashida, J. Hewitt, R.E. Hughes, T.E. Jeltema, G. Jóhannesson, A.S. Johnson, T. Kamae, J. Kataoka, J. Knödseder, M. Kuss, J. Lande, S. Larsson, L. Latronico, M. Llana Garde, F. Longo, F. Loparco, M.N. Lovellette, P. Lubrano, M. Mayer, M.N. Mazziotta, J.E. McEnery, P.F. Michelson, W. Mitthumsiri, T. Mizuno, M.E. Monzani, A. Morselli, I.V. Moskalenko, S. Murgia, R. Nemmen, E. Nuss, T. Ohsugi, M. Orienti, E. Orlando, J.F. Ormes, J.S. Perkins, M. Pesce-Rollins, F. Piron, G. Pivato, S. Rainò, R. Rando, M. Razzano, S. Razzaque, A. Reimer, O. Reimer, J. Ruan, M. Sánchez-Conde, A. Schulz, C. Sgrò, E.J. Siskind, G. Spandre, P. Spinelli, E. Storm, A.W. Strong, D.J. Suson, H. Takahashi, J.G. Thayer, J.B. Thayer, D.J. Thompson, L. Tibaldo, M. Tinivella, D.F. Torres, E. Troja, Y. Uchiyama, T.L. Usher, J. Vandenbroucke, G. Vianello, V. Vitale, B.L. Winer, K.S. Wood, S. Zimmer (Fermi-LAT Collaboration), A. Pinzke, C. Pfrommer, Search for cosmic-ray-induced gamma-ray emission in galaxy clusters. *Astrophys. J.* **787**, 18 (2014). <https://doi.org/10.1088/0004-637X/787/1/18>. arXiv:1308.5654
- P.A.R. Ade, N. Aghanim, M. Arnaud, F. Arroja, M. Ashdown, J. Aumont, C. Baccigalupi, M. Ballardini, A.J. Banday et al. (Planck Collaboration), Planck 2015 results. XIX. Constraints on primordial magnetic fields. *Astron. Astrophys.* **594**, A19 (2016). <https://doi.org/10.1051/0004-6361/201525821>. arXiv:1502.01594
- F. Aharonian, H. Akamatsu, F. Akimoto, S.W. Allen, N. Anabuki, L. Angelini, K. Arnaud, M. Audard, H. Awaki, M. Axelsson, A. Bamba, M. Bautz, R. Blandford, L. Brenneman, G.V. Brown, E. Bulbul, E. Cackett, M. Chernyakova, M. Chiao, P. Coppi, E. Costantini, J. de Plaa, J.W. den Herder, C. Done, T. Dotani, K. Ebisawa, M. Eckart, T. Enoto, Y. Ezoe, A.C. Fabian, C. Ferrigno, A. Foster, R. Fujimoto, Y. Fukazawa, A. Furuzawa, M. Galeazzi, L. Gallo, P. Gandhi, M. Giustini, A. Goldwurm, L. Gu, M. Guainazzi, Y. Haba, K. Hagino, K. Hamaguchi, I. Harrus, I. Hatsukade, K. Hayashi, T. Hayashi, K. Hayashida, J. Hiraga, A. Hornschemeier, A. Hoshino, J. Hughes, R. Iizuka, H. Inoue, Y. Inoue, K. Ishibashi, M. Ishida, K. Ishikawa, Y. Ishisaki, M. Itoh, N. Iyamoto, J. Kaastra, T. Kallman, T. Kamae, E. Kara, J. Kataoka, S. Katsuda, J. Katsuta, M. Kawaharada, N. Kawai, R. Kelley, D. Khangulyan, C. Kilbourne, A. King, T. Kitaguchi, S. Kitamoto, T. Kitayama, T. Kohmura, M. Kokubun, S. Koyama, K. Koyama, P. Kretschmar, H. Krimm, A. Kubota, H. Kunieda, P. Laurent, F. Lebrun, S.H. Lee, M. Leutenegger, O. Limousin, M. Loewenstein, K.S. Long, D. Lumb, G. Madejski, Y. Maeda, D. Maier, K. Makishima, M. Markevitch, H. Matsumoto, K. Matsushita, D. McCammon, B. McNamara, M. Mehdipour, E. Miller, J. Miller, S. Mineshige, K. Mitsuda, I. Mitsuishi, T. Miyazawa, T. Mizuno, H. Mori, K. Mori, H. Moseley, K. Mukai, H. Murakami, T. Murakami, R. Mushotzky, R. Nagino, T. Nakagawa, H. Nakajima, T. Nakamori, T. Nakano, S. Nakashima, K. Nakazawa, M. Nobukawa, H. Noda, M. Nomachi, S. O’Dell, H. Odaka, T. Ohashi, M. Ohno, T. Okajima, N. Ota, M. Ozaki, F. Paerels, S. Paltani, A. Parmar, R. Petre, C. Pinto, M. Pohl, F.S. Porter, K. Pottschmidt, B. Ramsey, C. Reynolds, H. Russell, S. Safi-Harb, S. Saito, K. Sakai, H. Sameshima, G. Sato, K. Sato, R. Sato, M. Sawada, N. Schartel, P. Serlemitsos, H. Seta, M. Shidatsu, A. Simionescu, R. Smith, Y. Soong, L. Stawarz, Y.

- Sugawara, S. Sugita, A. Szymkowiak, H. Tajima, H. Takahashi, T. Takahashi, S. Takeda, Y. Takei, T. Tamagawa, K. Tamura, T. Tamura, T. Tanaka, Y. Tanaka, M. Tashiro, Y. Tawara, Y. Terada, Y. Terashima, F. Tombesi, H. Tomida, Y. Tsuboi, M. Tsujimoto, H. Tsunemi, T. Tsuru, H. Uchida, H. Uchiyama, Y. Uchiyama, S. Ueda, Y. Ueda, S. Ueno, S. Uno, M. Urry, E. Ursino, C. de Vries, S. Watanabe, N. Werner, D. Wik, D. Wilkins, B. Williams, S. Yamada, H. Yamaguchi, K. Yamaoka, N.Y. Yamasaki, M. Yamauchi, S. Yamauchi, T. Yaqoob, Y. Yatsu, D. Yonetoku, A. Yoshida, T. Yuasa, I. Zhuravleva, A. Zoghbi (Hitomi Collaboration), The quiescent intracluster medium in the core of the Perseus cluster. *Nature* **535**, 117–121 (2016). <https://doi.org/10.1038/nature18627>. arXiv:1607.04487
- H. Alfvén, Existence of electromagnetic-hydrodynamic waves. *Nature* **150**, 405–406 (1942). <https://doi.org/10.1038/150405d0>
- T.C. Arlen, V.V. Vassilev, T. Weisgarber, S.P. Wakely, S. Yusef Shafi, Intergalactic magnetic fields and gamma-ray observations of extreme TeV blazars. *Astrophys. J.* **796**, 18 (2014). <https://doi.org/10.1088/0004-637X/796/1/18>. arXiv:1210.2802
- N. Asai, N. Fukuda, R. Matsumoto, Magnetohydrodynamic simulations of the formation of cold fronts in clusters of galaxies including heat conduction. *Astrophys. J. Lett.* **606**, L105–L108 (2004). <https://doi.org/10.1086/421440>. arXiv:astro-ph/0404160
- N. Asai, N. Fukuda, R. Matsumoto, Three-dimensional MHD simulations of X-ray emitting subcluster plasmas in cluster of galaxies. *Adv. Space Res.* **36**, 636–642 (2005). <https://doi.org/10.1016/j.asr.2005.04.041>. arXiv:astro-ph/0504227
- N. Asai, N. Fukuda, R. Matsumoto, Three-dimensional magnetohydrodynamic simulations of cold fronts in magnetically turbulent ICM. *Astrophys. J.* **663**, 816–823 (2007). <https://doi.org/10.1086/518235>. arXiv:astro-ph/0703536
- X.N. Bai, D. Caprioli, L. Sironi, A. Spitkovsky, Magnetohydrodynamic-particle-in-cell method for coupling cosmic rays with a thermal plasma: application to non-relativistic shocks. *Astrophys. J.* **809**, 55 (2015). <https://doi.org/10.1088/0004-637X/809/1/55>. arXiv:1412.1087
- D.S. Balsara, Higher-order accurate space-time schemes for computational astrophysics—Part I: Finite volume methods. *Living Rev. Comput. Astrophys.* **3**, 2 (2017). <https://doi.org/10.1007/s41115-017-0002-8>. arXiv:1703.01241
- J.D. Barrow, P.G. Ferreira, J. Silk, Constraints on a primordial magnetic field. *Phys. Rev. Lett.* **78**, 3610–3613 (1997). <https://doi.org/10.1103/PhysRevLett.78.3610>. arXiv:astro-ph/9701063
- G.K. Batchelor, On the spontaneous magnetic field in a conducting liquid in turbulent motion. *Proc. R. Soc. Lond. Ser. A, Math. Phys. Sci.* **201**, 405–416 (1950). <https://doi.org/10.1098/rspa.1950.0069>
- N. Battaglia, J.R. Bond, C. Pfrommer, J.L. Sievers, On the cluster physics of Sunyaev-Zel'dovich and X-ray surveys. I. The influence of feedback, non-thermal pressure, and cluster shapes on Y-M scaling relations. *Astrophys. J.* **758**, 74 (2012). <https://doi.org/10.1088/0004-637X/758/2/74>. arXiv:1109.3709
- A. Bauer, V. Springel, Subsonic turbulence in smoothed particle hydrodynamics and moving-mesh simulations. *Mon. Not. R. Astron. Soc.* **423**, 2558–2578 (2012). <https://doi.org/10.1111/j.1365-2966.2012.21058.x>. arXiv:1109.4413
- A. Bauer, K. Schaal, V. Springel, P. Chandrashekar, R. Pakmor, C. Klingenberg, Simulating turbulence using the astrophysical discontinuous Galerkin code TENET. arXiv:1602.09079 (2016)
- R. Beck, M. Krause, Revised equipartition and minimum energy formula for magnetic field strength estimates from radio synchrotron observations. *Astron. Nachr.* **326**, 414–427 (2005). <https://doi.org/10.1002/asna.200510366>. arXiv:astro-ph/0507367
- A.M. Beck, H. Lesch, K. Dolag, H. Kotarba, A. Geng, F.A. Staszczyn, Origin of strong magnetic fields in Milky Way-like galactic haloes. *Mon. Not. R. Astron. Soc.* **422**, 2152–2163 (2012). <https://doi.org/10.1111/j.1365-2966.2012.20759.x>. arXiv:1202.3349
- A.M. Beck, K. Dolag, H. Lesch, P.P. Kronberg, Strong magnetic fields and large rotation measures in protogalaxies from supernova seeding. *Mon. Not. R. Astron. Soc.* **435**, 3575–3586 (2013a). <https://doi.org/10.1093/mnras/stt1549>. arXiv:1308.3440
- A.M. Beck, M. Hanasz, H. Lesch, R.S. Remus, F.A. Staszczyn, On the magnetic fields in voids. *Mon. Not. R. Astron. Soc.* **429**, L60–L64 (2013b). <https://doi.org/10.1093/mnras/ls026>. arXiv:1210.8360
- A.M. Beck, G. Murante, A. Arth, R.S. Remus, A.F. Teklu, J.M.F. Donnert, S. Planelles, M.C. Beck, P. Förster, M. Imgrund, K. Dolag, S. Borgani, An improved SPH scheme for cosmological simulations. *Mon. Not. R. Astron. Soc.* **455**, 2110–2130 (2016). <https://doi.org/10.1093/mnras/stv2443>. arXiv:1502.07358
- A.R. Bell, Turbulent amplification of magnetic field and diffusive shock acceleration of cosmic rays. *Mon. Not. R. Astron. Soc.* **353**, 550–558 (2004). <https://doi.org/10.1111/j.1365-2966.2004.08097.x>
- A. Beresnyak, A. Lazarian, Polarization intermittency and its influence on MHD turbulence. *Astrophys. J. Lett.* **640**, L175–L178 (2006). <https://doi.org/10.1086/503708>. arXiv:astro-ph/0512315
- A. Beresnyak, F. Miniati, Turbulent amplification and structure of the intracluster magnetic field. *Astrophys. J.* **817**, 127 (2016). <https://doi.org/10.3847/0004-637X/817/2/127>. arXiv:1507.00342

- S. Bertone, C. Vogt, T. Enßlin, Magnetic field seeding by galactic winds. *Mon. Not. R. Astron. Soc.* **370**, 319–330 (2006). <https://doi.org/10.1111/j.1365-2966.2006.10474.x>. arXiv:astro-ph/0604462
- L. Biermann, A. Schlüter, Cosmic radiation and cosmic magnetic fields. II. Origin of cosmic magnetic fields. *Phys. Rev.* **82**, 863–868 (1951). <https://doi.org/10.1103/PhysRev.82.863>
- J. Binney, Galaxy formation without primordial turbulence: mechanisms for generating cosmic vorticity. *Mon. Not. R. Astron. Soc.* **168**, 73–92 (1974). <https://doi.org/10.1093/mnras/168.1.73>
- Y. Birnboim, U. Keshet, L. Hernquist, Cold fronts by merging of shocks. *Mon. Not. R. Astron. Soc.* **408**, 199–212 (2010). <https://doi.org/10.1111/j.1365-2966.2010.17176.x>. arXiv:1006.1892
- H. Böhringer, N. Werner, X-ray spectroscopy of galaxy clusters: studying astrophysical processes in the largest celestial laboratories. *Astron. Astrophys. Rev.* **18**, 127–196 (2010). <https://doi.org/10.1007/s00159-009-0023-3>
- A. Bonafede, L. Feretti, M. Murgia, F. Govoni, G. Giovannini, D. Dallacasa, K. Dolag, G.B. Taylor, The Coma cluster magnetic field from Faraday rotation measures. *Astron. Astrophys.* **513**, A30 (2010). <https://doi.org/10.1051/0004-6361/200913696>. arXiv:1002.0594
- A. Bonafede, K. Dolag, F. Stasyszyn, G. Murante, S. Borgani, A non-ideal magnetohydrodynamic GADGET: simulating massive galaxy clusters. *Mon. Not. R. Astron. Soc.* **418**, 2234–2250 (2011). <https://doi.org/10.1111/j.1365-2966.2011.19523.x>. arXiv:1107.0968
- A. Bonafede, F. Vazza, M. Brüggen, M. Murgia, F. Govoni, L. Feretti, G. Giovannini, G. Ogrean, Measurements and simulation of Faraday rotation across the Coma radio relic. *Mon. Not. R. Astron. Soc.* **433**, 3208–3226 (2013). <https://doi.org/10.1093/mnras/stt960>. arXiv:1305.7228
- A. Bonafede, F. Vazza, M. Brüggen, T. Akahori, E. Carretti, S. Colafrancesco, L. Feretti, C. Ferrari, G. Giovannini, F. Govoni, M. Johnston-Hollitt, M. Murgia, A. Scaife, V. Vacca, F. Govoni, L. Rudnick, A. Scaife, Unravelling the origin of large-scale magnetic fields in galaxy clusters and beyond through Faraday Rotation Measures with the SKA, in *Advancing Astrophysics with the Square Kilometre Array (AASKA14)* (2015), p. 95. arXiv:1501.00321
- S. Borgani, A. Diaferio, K. Dolag, S. Schindler, Thermodynamical properties of the ICM from hydrodynamical simulations. *Space Sci. Rev.* **134**, 269–293 (2008). <https://doi.org/10.1007/s11214-008-9317-4>. arXiv:0801.1032
- A. Brandenburg, Nonlinear small-scale dynamos at low magnetic Prandtl numbers. *Astrophys. J.* **741**, 92 (2011). <https://doi.org/10.1088/0004-637X/741/2/92>. arXiv:1106.5777
- A. Brandenburg, K. Subramanian, Astrophysical magnetic fields and nonlinear dynamo theory. *Phys. Rep.* **417**, 1–209 (2005). <https://doi.org/10.1016/j.physrep.2005.06.005>. arXiv:astro-ph/0405052
- A.E. Broderick, P. Chang, C. Pfommer, The cosmological impact of luminous TeV blazars. I. Implications of plasma instabilities for the intergalactic magnetic field and extragalactic gamma-ray background. *Astrophys. J.* **752**, 22 (2012). <https://doi.org/10.1088/0004-637X/752/1/22>. arXiv:1106.5494
- M. Brüggen, Magnetic field amplification by cosmic ray-driven turbulence—I. Isotropic CR diffusion. *Mon. Not. R. Astron. Soc.* **436**, 294–303 (2013). <https://doi.org/10.1093/mnras/stt1566>. arXiv:1308.5230
- M. Brüggen, M. Hoeft, M. Ruszkowski, X-ray line tomography of AGN-induced motion in clusters of galaxies. *Astrophys. J.* **628**, 153–159 (2005a). <https://doi.org/10.1086/430732>. arXiv:astro-ph/0503656
- M. Brüggen, M. Ruszkowski, A. Simionescu, M. Hoeft, C. Dalla Vecchia, Simulations of magnetic fields in filaments. *Astrophys. J. Lett.* **631**, L21–L24 (2005b). <https://doi.org/10.1086/497004>. arXiv:astro-ph/0508231
- M. Brüggen, A. Bykov, D. Ryu, H. Röttgering, Magnetic fields, relativistic particles, and shock waves in cluster outskirts. *Space Sci. Rev.* **166**, 187–213 (2012). <https://doi.org/10.1007/s11214-011-9785-9>. arXiv:1107.5223
- G. Brunetti, T.W. Jones, Cosmic rays in galaxy clusters and their nonthermal emission. *Int. J. Mod. Phys. B* **23**, 1430007 (2014). <https://doi.org/10.1142/S0218271814300079>. arXiv:1401.7519
- G. Brunetti, A. Lazarian, Compressible turbulence in galaxy clusters: physics and stochastic particle reacceleration. *Mon. Not. R. Astron. Soc.* **378**, 245–275 (2007). <https://doi.org/10.1111/j.1365-2966.2007.11771.x>. arXiv:astro-ph/0703591
- G. Brunetti, A. Lazarian, Acceleration of primary and secondary particles in galaxy clusters by compressible MHD turbulence: from radio haloes to gamma-rays. *Mon. Not. R. Astron. Soc.* **410**, 127–142 (2011a). <https://doi.org/10.1111/j.1365-2966.2010.17457.x>. arXiv:1008.0184
- G. Brunetti, A. Lazarian, Particle reacceleration by compressible turbulence in galaxy clusters: effects of a reduced mean free path. *Mon. Not. R. Astron. Soc.* **412**, 817–824 (2011b). <https://doi.org/10.1111/j.1365-2966.2010.17937.x>. arXiv:1011.1198
- G. Brunetti, A. Lazarian, Particle reacceleration by compressible turbulence in galaxy clusters: effects of a reduced mean free path. *Mon. Not. R. Astron. Soc.* **412**, 817–824 (2011c). <https://doi.org/10.1111/j.1365-2966.2010.17937.x>. arXiv:1011.1198
- G. Brunetti, P. Blasi, R. Cassano, S. Gabici, Alfvénic reacceleration of relativistic particles in galaxy clusters: MHD waves, leptons and hadrons. *Mon. Not. R. Astron. Soc.* **350**, 1174–1194 (2004)

- G.R. Burbidge, Possible sources of radio emission in clusters of galaxies. *Astrophys. J.* **128**, 1 (1958). <https://doi.org/10.1086/146509>
- J.M. Burgers, Some considerations on the fields of stress connected with dislocations in a regular crystal lattice I. *Proc. K. Ned. Akad. Wet.* **42**, 293–378 (1939)
- J.O. Burns, S.W. Skillman, B.W. O’Shea, Galaxy clusters at the edge: temperature, entropy, and gas dynamics near the virial radius. *Astrophys. J.* **721**, 1105–1112 (2010). <https://doi.org/10.1088/0004-637X/721/2/1105>. arXiv:1004.3553
- I. Butsky, J. Zrake, K. Jh, H.I. Yang, T. Abel, Ab initio simulations of a supernova-driven galactic dynamo in an isolated disk galaxy. *Astrophys. J.* **843**, 113 (2017). <https://doi.org/10.3847/1538-4357/aa799f>. arXiv:1610.08528
- A.M. Bykov, S.M. Osipov, I.N. Toptygin, Long-wavelength MHD instability in the prefront of collisionless shocks with accelerated particles. *Astron. Lett.* **35**, 555–563 (2009). <https://doi.org/10.1134/S1063773709080052>
- A.M. Bykov, S.M. Osipov, D.C. Ellison, Cosmic ray current driven turbulence in shocks with efficient particle acceleration: the oblique, long-wavelength mode instability. *Mon. Not. R. Astron. Soc.* **410**, 39–52 (2011). <https://doi.org/10.1111/j.1365-2966.2010.17421.x>. arXiv:1010.0408
- A.M. Bykov, D.C. Ellison, S.M. Osipov, A.E. Vladimirov, Magnetic field amplification in nonlinear diffusive shock acceleration including resonant and non-resonant cosmic-ray driven instabilities. *Astrophys. J.* **789**, 137 (2014). <https://doi.org/10.1088/0004-637X/789/2/137>. arXiv:1406.0084
- L. Campanelli, Helical magnetic fields from inflation. *Int. J. Mod. Phys. B* **18**, 1395–1411 (2009). <https://doi.org/10.1142/S0218271809015175>. arXiv:0805.0575
- C. Caprini, S. Gabici, Gamma-ray observations of blazars and the intergalactic magnetic field spectrum. *Phys. Rev. Lett.* **91**(12), 123514 (2015). <https://doi.org/10.1103/PhysRevD.91.123514>. arXiv:1504.00383
- D. Caprioli, Cosmic-ray acceleration in supernova remnants: non-linear theory revised. *J. Cosmol. Astropart. Phys.* **7**, 038 (2012). <https://doi.org/10.1088/1475-7516/2012/07/038>. arXiv:1206.1360
- D. Caprioli, A. Spitkovsky, Simulations of ion acceleration at non-relativistic shocks. I. Acceleration efficiency. *Astrophys. J.* **783**, 91 (2014a). <https://doi.org/10.1088/0004-637X/783/2/91>. arXiv:1310.2943
- D. Caprioli, A. Spitkovsky, Simulations of ion acceleration at non-relativistic shocks. II. Magnetic field amplification. *Astrophys. J.* **794**, 46 (2014b). <https://doi.org/10.1088/0004-637X/794/1/46>. arXiv:1401.7679
- S. Chandrasekhar, *Hydrodynamic and Hydromagnetic Stability* (1961)
- W. Chen, J.H. Buckley, F. Ferrer, Search for GeV  $\gamma$ -ray pair halos around low redshift blazars. *Phys. Rev. Lett.* **115**(21), 211103 (2015). <https://doi.org/10.1103/PhysRevLett.115.211103>. arXiv:1410.7717
- H. Chen, C. Jones, F. Andrade-Santos, J.A. ZuHone, Z. Li, Gas sloshing in Abell 2204: constraining the properties of the magnetized intracluster medium. *Astrophys. J.* **838**, 38 (2017). <https://doi.org/10.3847/1538-4357/aa64de>. arXiv:1703.01895
- A.D. Chernin, A. Cosmological model with a disordered magnetic field. *Sov. Astron.* **10**, 634 (1967)
- J. Cho, E.T. Vishniac, A. Beresnyak, A. Lazarian, D. Ryu, Growth of magnetic fields induced by turbulent motions. *Astrophys. J.* **693**, 1449–1461 (2009). <https://doi.org/10.1088/0004-637X/693/2/1449>. arXiv:0812.0817
- E. Churazov, W. Forman, C. Jones, R. Sunyaev, H. Böhringer, XMM-Newton observations of the Perseus cluster—II. Evidence for gas motions in the core. *Mon. Not. R. Astron. Soc.* **347**, 29–35 (2004). <https://doi.org/10.1111/j.1365-2966.2004.07201.x>. arXiv:astro-ph/0309427
- D.S. De Young, Galaxy-driven turbulence and the growth of intracluster magnetic fields. *Astrophys. J.* **386**, 464–472 (1992). <https://doi.org/10.1086/171032>
- A. Dedner, F. Kemm, D. Kröner, C.D. Munz, T. Schnitzer, M. Wesenberg, Hyperbolic divergence cleaning for the MHD equations. *J. Comput. Phys.* **175**, 645–673 (2002). <https://doi.org/10.1006/jcph.2001.6961>
- B. Dennison, Formation of radio halos in clusters of galaxies from cosmic-ray protons. *Astrophys. J. Lett.* **239**, L93–L96 (1980). <https://doi.org/10.1086/183300>
- C.L. Dobbs, D.J. Price, A.R. Pettitt, M.R. Bate, T.S. Tricco, Magnetic field evolution and reversals in spiral galaxies. *Mon. Not. R. Astron. Soc.* **461**, 4482–4495 (2016). <https://doi.org/10.1093/mnras/stw1625>. arXiv:1607.05532
- W. Dobler, N.E. Haugen, T.A. Yousef, A. Brandenburg, Bottleneck effect in three-dimensional turbulence simulations. *Phys. Rev. E* **68**(2), 026304 (2003). <https://doi.org/10.1103/PhysRevE.68.026304>. arXiv:astro-ph/0303324
- K. Dolag, T.A. EnBlin, Radio halos of galaxy clusters from hadronic secondary electron injection in realistic magnetic field configurations. *Astron. Astrophys.* **362**, 151–157 (2000)
- K. Dolag, S. Schindler, The effect of magnetic fields on the mass determination of clusters of galaxies. *Astron. Astrophys.* **364**, 491–496 (2000). arXiv:astro-ph/0010296
- K. Dolag, F. Stasyszyn, An MHD GADGET for cosmological simulations. *Mon. Not. R. Astron. Soc.* **398**, 1678–1697 (2009). <https://doi.org/10.1111/j.1365-2966.2009.15181.x>. arXiv:0807.3553



- K. Dolag, M. Bartelmann, H. Lesch, SPH simulations of magnetic fields in galaxy clusters. *Astron. Astrophys.* **348**, 351–363 (1999). [arXiv:astro-ph/0202272](https://arxiv.org/abs/astro-ph/0202272)
- K. Dolag, A. Evrard, M. Bartelmann, The temperature-mass relation in magnetized galaxy clusters. *Astron. Astrophys.* **369**, 36–41 (2001). <https://doi.org/10.1051/0004-6361:20010094>. [arXiv:astro-ph/0101302](https://arxiv.org/abs/astro-ph/0101302)
- K. Dolag, M. Bartelmann, H. Lesch, Evolution and structure of magnetic fields in simulated galaxy clusters. *Astron. Astrophys.* **387**, 383–395 (2002). <https://doi.org/10.1051/0004-6361:20020241>
- K. Dolag, D. Grasso, V. Springel, I. Tkachev, Constrained simulations of the magnetic field in the local Universe and the propagation of ultrahigh energy cosmic rays. *J. Cosmol. Astropart. Phys.* **1**, 009 (2005a). <https://doi.org/10.1088/1475-7516/2005/01/009>. [arXiv:astro-ph/0410419](https://arxiv.org/abs/astro-ph/0410419)
- K. Dolag, F. Vazza, G. Brunetti, G. Tormen, Turbulent gas motions in galaxy cluster simulations: the role of smoothed particle hydrodynamics viscosity. *Mon. Not. R. Astron. Soc.* **364**, 753–772 (2005b). <https://doi.org/10.1111/j.1365-2966.2005.09630.x>. [arXiv:astro-ph/0507480](https://arxiv.org/abs/astro-ph/0507480)
- K. Dolag, A.M. Bykov, A. Diaferio, Non-thermal processes in cosmological simulations. *Space Sci. Rev.* **134**, 311–335 (2008). <https://doi.org/10.1007/s11214-008-9319-2>. [arXiv:0801.1048](https://arxiv.org/abs/0801.1048)
- K. Dolag, M. Kachelrieß, S. Ostapchenko, R. Tomàs, Blazar halos as probe for extragalactic magnetic fields and maximal acceleration energy. *Astrophys. J.* **703**, 1078–1085 (2009). <https://doi.org/10.1088/0004-637X/703/1/1078>. [arXiv:0903.2842](https://arxiv.org/abs/0903.2842)
- K. Dolag, M. Kachelrieß, S. Ostapchenko, R. Tomàs, Lower limit on the strength and filling factor of extragalactic magnetic fields. *Astrophys. J. Lett.* **727**, L4 (2011). <https://doi.org/10.1088/2041-8205/727/1/L4>. [arXiv:1009.1782](https://arxiv.org/abs/1009.1782)
- K. Dolag, B.M. Gaensler, A.M. Beck, M.C. Beck, Constraints on the distribution and energetics of fast radio bursts using cosmological hydrodynamic simulations. *Mon. Not. R. Astron. Soc.* **451**, 4277–4289 (2015). <https://doi.org/10.1093/mnras/stv1190>. [arXiv:1412.4829](https://arxiv.org/abs/1412.4829)
- J. Donnert, G. Brunetti, An efficient Fokker-Planck solver and its application to stochastic particle acceleration in galaxy clusters. *Mon. Not. R. Astron. Soc.* **443**, 3564–3577 (2014). <https://doi.org/10.1093/mnras/stu1417>. [arXiv:1407.2735](https://arxiv.org/abs/1407.2735)
- J. Donnert, K. Dolag, H. Lesch, E. Müller, Cluster magnetic fields from galactic outflows. *Mon. Not. R. Astron. Soc.* **392**, 1008–1021 (2009). <https://doi.org/10.1111/j.1365-2966.2008.14132.x>. [arXiv:0808.0919](https://arxiv.org/abs/0808.0919)
- J. Donnert, K. Dolag, G. Brunetti, R. Cassano, A. Bonafede, Radio haloes from simulations and hadronic models—I. The Coma cluster. *Mon. Not. R. Astron. Soc.* **401**, 47–54 (2010). <https://doi.org/10.1111/j.1365-2966.2009.15655.x>. [arXiv:0905.2418](https://arxiv.org/abs/0905.2418)
- J.M.F. Donnert, A. Stroe, G. Brunetti, D. Hoang, H. Roettgering, Magnetic field evolution in giant radio relics using the example of CIZA J2242.8+5301. *Mon. Not. R. Astron. Soc.* **462**, 2014–2032 (2016). <https://doi.org/10.1093/mnras/stw1792>. [arXiv:1603.06570](https://arxiv.org/abs/1603.06570)
- J.M.F. Donnert, A.M. Beck, K. Dolag, H.J.A. Röttgering, Simulations of the galaxy cluster CIZA J2242.8+5301—I. Thermal model and shock properties. *Mon. Not. R. Astron. Soc.* **471**, 4587–4605 (2017). <https://doi.org/10.1093/mnras/stx1819>. [arXiv:1703.05682](https://arxiv.org/abs/1703.05682)
- L.O. Drury, T.P. Downes, Turbulent magnetic field amplification driven by cosmic ray pressure gradients. *Mon. Not. R. Astron. Soc.* **427**, 2308–2313 (2012). <https://doi.org/10.1111/j.1365-2966.2012.22106.x>. [arXiv:1205.6823](https://arxiv.org/abs/1205.6823)
- Y. Dubois, R. Teyssier, Cosmological MHD simulation of a cooling flow cluster. *Astron. Astrophys.* **482**, L13–L16 (2008). <https://doi.org/10.1051/0004-6361:200809513>. [arXiv:0802.0490](https://arxiv.org/abs/0802.0490)
- Y. Dubois, J. Devriendt, A. Slyz, J. Silk, Influence of AGN jets on the magnetized ICM. *Mon. Not. R. Astron. Soc.* **399**, L49–L53 (2009). <https://doi.org/10.1111/j.1745-3933.2009.00721.x>. [arXiv:0905.3345](https://arxiv.org/abs/0905.3345)
- R. Durrer, A. Neronov, Cosmological magnetic fields: their generation, evolution and observation. *Astron. Astrophys. Rev.* **21**, 62 (2013). <https://doi.org/10.1007/s00159-013-0062-7>. [arXiv:1303.7121](https://arxiv.org/abs/1303.7121)
- L.J. Dursi, Bubble wrap for bullets: the stability imparted by a thin magnetic layer. *Astrophys. J.* **670**, 221–230 (2007). <https://doi.org/10.1086/521997>. [arXiv:0706.3216](https://arxiv.org/abs/0706.3216)
- L.J. Dursi, C. Pfrommer, Draping of cluster magnetic fields over bullets and bubbles—morphology and dynamic effects. *Astrophys. J.* **677**, 993–1018 (2008). <https://doi.org/10.1086/529371>. [arXiv:0711.0213](https://arxiv.org/abs/0711.0213)
- T.A. Enßlin, C. Vogt, Magnetic turbulence in cool cores of galaxy clusters. *Astron. Astrophys.* **453**, 447–458 (2006). <https://doi.org/10.1051/0004-6361:20053518>. [arXiv:astro-ph/0505517](https://arxiv.org/abs/astro-ph/0505517)
- T. Enßlin, C. Pfrommer, F. Miniati, K. Subramanian, Cosmic ray transport in galaxy clusters: implications for radio halos, gamma-ray signatures, and cool core heating. *Astron. Astrophys.* **527**, A99 (2011). <https://doi.org/10.1051/0004-6361/201015652>. [arXiv:1008.4717](https://arxiv.org/abs/1008.4717)
- S. Etori, G.W. Pratt, J. de Plaa, D. Eckert, J. Nevalainen, E.S. Battistelli, S. Borgani, J.H. Croston, A. Finoguenov, J. Kaastra, M. Gaspari, F. Gastaldello, M. Gitti, S. Molendi, E. Pointecouteau, T.J. Ponman, T.H. Reiprich, M. Roncarelli, M. Rossetti, J.S. Sanders, M. Sun, G. Trinchieri, F. Vazza, M. Arnaud, H. Böhringer, F. Brighenti, H. Dahle, S. De Grandi, J.J. Mohr, A. Moretti, S. Schindler, The hot and energetic universe: the astrophysics of galaxy groups and clusters. [arXiv:1306.2322](https://arxiv.org/abs/1306.2322) (2013)

- A.E. Evrard, Formation and evolution of X-ray clusters—a hydrodynamic simulation of the intracluster medium. *Astrophys. J.* **363**, 349–366 (1990). <https://doi.org/10.1086/169350>
- C. Federrath, Magnetic field amplification in turbulent astrophysical plasmas. *J. Plasma Phys.* **82**(6), 535820601 (2016). <https://doi.org/10.1017/S0022377816001069>. arXiv:1610.08132
- C. Federrath, G. Chabrier, J. Schober, R. Banerjee, R.S. Klessen, D.R.G. Schleicher, Mach number dependence of turbulent magnetic field amplification: solenoidal versus compressive flows. *Phys. Rev. Lett.* **107**(11), 114504 (2011a). <https://doi.org/10.1103/PhysRevLett.107.114504>. arXiv:1109.1760
- C. Federrath, S. Sur, D.R.G. Schleicher, R. Banerjee, R.S. Klessen, A new Jeans resolution criterion for (M)HD simulations of self-gravitating gas: application to magnetic field amplification by gravity-driven turbulence. *Astrophys. J.* **731**, 62 (2011b). <https://doi.org/10.1088/0004-637X/731/1/62>. arXiv:1102.0266
- C. Federrath, J. Schober, S. Bovino, D.R.G. Schleicher, The turbulent dynamo in highly compressible supersonic plasmas. *Astrophys. J. Lett.* **797**, L19 (2014). <https://doi.org/10.1088/2041-8205/797/2/L19>. arXiv:1411.4707
- C. Ferrari, F. Govoni, S. Schindler, A.M. Bykov, Y. Rephaeli, Observations of extended radio emission in clusters. *Space Sci. Rev.* **134**, 93–118 (2008). <https://doi.org/10.1007/s11214-008-9311-x>. arXiv:0801.0985
- A. Finoguenov, C.L. Sarazin, K. Nakazawa, D.R. Wik, T.E. Clarke, XMM-Newton observation of the northwest radio relic region in A3667. *Astrophys. J.* **715**, 1143–1151 (2010). <https://doi.org/10.1088/0004-637X/715/2/1143>. arXiv:1004.2331
- F. Fraschetti, Turbulent amplification of a magnetic field driven by the dynamo effect at rippled shocks. *Astrophys. J.* **770**, 84 (2013). <https://doi.org/10.1088/0004-637X/770/2/84>. arXiv:1304.4956
- S.R. Furlanetto, A. Loeb, Intergalactic magnetic fields from quasar outflows. *Astrophys. J.* **556**, 619–634 (2001). <https://doi.org/10.1086/321630>. arXiv:astro-ph/0102076
- B.M. Gaensler, T.L. Landecker, A.R. Taylor, POSSUM Collaboration, Survey science with ASKAP: Polarization Sky Survey of the Universe’s Magnetism (POSSUM), in *American Astronomical Society Meeting Abstracts #215*. Bulletin of the American Astronomical Society, vol. 42 (2010), p. 515.
- T.A. Gardiner, J.M. Stone, An unsplit Godunov method for ideal MHD via constrained transport in three dimensions. *J. Comput. Phys.* **227**, 4123–4141 (2008). <https://doi.org/10.1016/j.jcp.2007.12.017>. arXiv:0712.2634
- M. Gaspari, C. Melioli, F. Brighenti, A. D’Ercole, The dance of heating and cooling in galaxy clusters: three-dimensional simulations of self-regulated active galactic nuclei outflows. *Mon. Not. R. Astron. Soc.* **411**, 349–372 (2011). <https://doi.org/10.1111/j.1365-2966.2010.17688.x>. arXiv:1007.0674
- M. Gendron-Marsolaïs, J. Hlavacek-Larrondo, R.J. van Weeren, T. Clarke, A.C. Fabian, H.T. Intema, G.B. Taylor, K.M. Blundell, J.S. Sanders, Deep 230–470 MHz VLA observations of the mini-halo in the Perseus cluster. *Mon. Not. R. Astron. Soc.* **469**, 3872–3880 (2017). <https://doi.org/10.1093/mnras/stx1042>. arXiv:1701.03791
- C. Gheller, F. Vazza, M. Brüggén, M. Alpaslan, B.W. Holwerda, A.M. Hopkins, J. Liske, Evolution of cosmic filaments and of their galaxy population from MHD cosmological simulations. *Mon. Not. R. Astron. Soc.* **462**, 448–463 (2016). <https://doi.org/10.1093/mnras/stw1595>. arXiv:1607.01406
- J. Giacalone, J.R. Jokipii, Magnetic field amplification by shocks in turbulent fluids. *Astrophys. J. Lett.* **663**, L41–L44 (2007). <https://doi.org/10.1086/519994>
- S. Giacintucci, M. Markevitch, G. Brunetti, J.A. Zuhone, T. Venturi, P. Mazzotta, H. Bourdin, Mapping the particle acceleration in the cool core of the galaxy cluster RX J1720.1+2638. *Astrophys. J.* **795**, 73 (2014a). <https://doi.org/10.1088/0004-637X/795/1/73>. arXiv:1403.2820
- S. Giacintucci, M. Markevitch, T. Venturi, T.E. Clarke, R. Cassano, P. Mazzotta, New detections of radio minihalos in cool cores of galaxy clusters. *Astrophys. J.* **781**, 9 (2014b). <https://doi.org/10.1088/0004-637X/781/1/9>. arXiv:1311.5248
- S. Giacintucci, M. Markevitch, R. Cassano, T. Venturi, T.E. Clarke, G. Brunetti, Occurrence of radio minihalos in a mass-limited sample of galaxy clusters. *Astrophys. J.* **841**, 71 (2017). <https://doi.org/10.3847/1538-4357/aa7069>. arXiv:1701.01364
- P. Goldreich, S. Sridhar, Magnetohydrodynamic turbulence revisited. *Astrophys. J.* **485**, 680–688 (1997). <https://doi.org/10.1086/304442>. arXiv:astro-ph/9612243
- O. Goldshmidt, Y. Rephaeli, Turbulent generation of intracluster magnetic fields and Faraday rotation measurements. *Astrophys. J.* **411**, 518–528 (1993). <https://doi.org/10.1086/172853>
- F. Govoni, K. Dolag, M. Murgia, L. Feretti, S. Schindler, G. Giovannini, W. Boschin, V. Vacca, A. Bonafede, Rotation measures of radio sources in hot galaxy clusters. *Astron. Astrophys.* **522**, A105 (2010). <https://doi.org/10.1051/0004-6361/200913665>. arXiv:1007.5207
- F. Govoni, M. Johnston-Hollitt, I. Agudo, T. Akahori, R. Beck, A. Bonafede, T.D. Carozzi, S. Colafrancesco, L. Feretti, K. Ferriere, B.M. Gaensler, L. Harvey-Smith, M. Haverkorn, G.H. Heald, S.A. Mao, L. Rud-



- nick, D. Schnitzeler, A. Scaife, J.M. Stil, K. Takahashi, A.R. Taylor, O. Wucknitz, Cosmic magnetism science in the SKA1 era. Technical Report (2014)
- F. Govoni, M. Murgia, V. Vacca, F. Loi, M. Girardi, F. Gastaldello, G. Giovannini, L. Feretti, R. Paladino, E. Carretti, R. Concu, A. Melis, S. Poppi, G. Valente, G. Bernardi, A. Bonafede, W. Boschin, M. Brienza, T.E. Clarke, S. Colafrancesco, F. de Gasperin, D. Eckert, T.A. Enßlin, C. Ferrari, L. Gregorini, M. Johnston-Hollitt, H. Junklewitz, E. Orrù, P. Parma, R. Perley, M. Rossetti, G.B. Taylor, F. Vazza, Sardinia Radio Telescope observations of Abell 194. The intra-cluster magnetic field power spectrum. *Astron. Astrophys.* **603**, A122 (2017). <https://doi.org/10.1051/0004-6361/201630349>. arXiv:1703.08688
- F. Guo, S. Li, H. Li, J. Giacalone, J.R. Jokipii, D. Li, On the amplification of magnetic field by a supernova blast shock wave in a turbulent medium. *Astrophys. J.* **747**, 98 (2012). <https://doi.org/10.1088/0004-637X/747/2/98>. arXiv:1112.6373
- F. Guo, X. Li, H. Li, W. Daughton, B. Zhang, N. Lloyd-Ronning, Y.H. Liu, H. Zhang, W. Deng, Efficient production of high-energy nonthermal particles during magnetic reconnection in a magnetically dominated ion-electron plasma. *Astrophys. J. Lett.* **818**, L9 (2016). <https://doi.org/10.3847/2041-8205/818/1/L9>. arXiv:1511.01434
- E.R. Harrison, Magnetic fields in the early Universe. *Mon. Not. R. Astron. Soc.* **165**, 185 (1973). <https://doi.org/10.1093/mnras/165.2.185>
- N.E.L. Haugen, A. Brandenburg, Suppression of small scale dynamo action by an imposed magnetic field. *Phys. Rev. E* **70**(3), 036408 (2004). <https://doi.org/10.1103/PhysRevE.70.036408>. arXiv:astro-ph/0402281
- N.E.L. Haugen, A. Brandenburg, A.J. Mee, Mach number dependence of the onset of dynamo action. *Mon. Not. R. Astron. Soc.* **353**, 947–952 (2004). <https://doi.org/10.1111/j.1365-2966.2004.08127.x>. arXiv:astro-ph/0405453
- D.N. Hoang, T.W. Shimwell, A. Stroe, H. Akamatsu, G. Brunetti, J.M.F. Donnert, H.T. Intema, D.D. Mulcahy, H.J.A. Röttgering, R.J. van Weeren, A. Bonafede, M. Brüggen, R. Cassano, K.T. Chyzy, T. Enßlin, C. Ferrari, F. de Gasperin, L. Gu, M. Hoeft, G.K. Miley, E. Orrù, R. Pizzo, G.J. White, Deep LOFAR observations of the merging galaxy cluster CIZA J2242.8+5301. *Mon. Not. R. Astron. Soc.* **471**, 1107–1125 (2017). <https://doi.org/10.1093/mnras/stx1645>. arXiv:1706.09903
- P.F. Hopkins, A new class of accurate, mesh-free hydrodynamic simulation methods. *Mon. Not. R. Astron. Soc.* **450**, 53–110 (2015). <https://doi.org/10.1093/mnras/stv195>. arXiv:1409.7395
- P.F. Hopkins, M.J. Raives, Accurate, meshless methods for magnetohydrodynamics. *Mon. Not. R. Astron. Soc.* **455**, 51–88 (2016). <https://doi.org/10.1093/mnras/stv2180>. arXiv:1505.02783
- S. Hutschenreuter, S. Dorn, J. Jasche, F. Vazza, D. Paoletti, G. Lavaux, E. TA, The primordial magnetic field in our cosmic backyard. arXiv:1803.02629 (2018)
- L. Iapichino, M. Brüggen, Magnetic field amplification by shocks in galaxy clusters: application to radio relics. *Mon. Not. R. Astron. Soc.* **423**, 2781–2788 (2012). <https://doi.org/10.1111/j.1365-2966.2012.21084.x>. arXiv:1204.2455
- L. Iapichino, W. Schmidt, J.C. Niemeyer, J. Merklein, Turbulence production and turbulent pressure support in the intergalactic medium. *Mon. Not. R. Astron. Soc.* **414**, 2297–2308 (2011). <https://doi.org/10.1111/j.1365-2966.2011.18550.x>. arXiv:1102.3352
- L. Iapichino, C. Federrath, R.S. Klessen, Adaptive mesh refinement simulations of a galaxy cluster merger—I. Resolving and modelling the turbulent flow in the cluster outskirts. *Mon. Not. R. Astron. Soc.* **469**, 3641–3655 (2017). <https://doi.org/10.1093/mnras/stx882>. arXiv:1704.02922
- Y. Ichinohe, A. Simionescu, N. Werner, T. Takahashi, An azimuthally resolved study of the cold front in Abell 3667. *Mon. Not. R. Astron. Soc.* **467**, 3662–3676 (2017). <https://doi.org/10.1093/mnras/stx280>. arXiv:1702.01026
- T. Inoue, R. Yamazaki, S. Inutsuka, Turbulence and magnetic field amplification in supernova remnants: interactions between a strong shock wave and multiphase interstellar medium. *Astrophys. J.* **695**, 825–833 (2009). <https://doi.org/10.1088/0004-637X/695/2/825>. arXiv:0901.0486
- S. Jacob, C. Pfrommer, Cosmic ray heating in cool core clusters—I. Diversity of steady state solutions. *Mon. Not. R. Astron. Soc.* **467**, 1449–1477 (2017a). <https://doi.org/10.1093/mnras/stx131>. arXiv:1609.06321
- S. Jacob, C. Pfrommer, Cosmic ray heating in cool core clusters—II. Self-regulation cycle and non-thermal emission. *Mon. Not. R. Astron. Soc.* **467**, 1478–1495 (2017b). <https://doi.org/10.1093/mnras/stx132>. arXiv:1609.06322
- W. Jaffe, On the morphology of the magnetic field in galaxy clusters. *Astrophys. J.* **241**, 925–927 (1980). <https://doi.org/10.1086/158407>
- S. Ji, S.P. Oh, M. Ruszkowski, M. Markevitch, The efficiency of magnetic field amplification at shocks by turbulence. *Mon. Not. R. Astron. Soc.* **463**, 3989–4003 (2016). <https://doi.org/10.1093/mnras/stw2320>. arXiv:1603.08518
- T.W. Jones, D.H. Porter, D. Ryu, J. Cho, Cluster turbulence: simulation insights. *Mem. Soc. Astron. Ital.* **82**, 588 (2011). arXiv:1101.4050

- T. Kahniashvili, A.G. Tevzadze, S.K. Sethi, K. Pandey, B. Ratna, Primordial magnetic field limits from cosmological data. *Phys. Rev. Lett.* **82**(8), 083005 (2010). <https://doi.org/10.1103/PhysRevD.82.083005>. arXiv:1009.2094
- T. Kahniashvili, A.G. Tevzadze, B. Ratna, Phase transition generated cosmological magnetic field at large scales. *Astrophys. J.* **726**, 78 (2011). <https://doi.org/10.1088/0004-637X/726/2/78>. arXiv:0907.0197
- T. Kahniashvili, A. Brandenburg, A.G. Tevzadze, The evolution of primordial magnetic fields since their generation. *Phys. Scr.* **91**(10), 104008 (2016). <https://doi.org/10.1088/0031-8949/91/10/104008>. arXiv:1507.00510
- H. Kang, D. Ryu, R. Cen, J.P. Ostriker, *Astrophys. J.* **669**, 729–740 (2007). <https://doi.org/10.1086/521717>. arXiv:0704.1521
- A.P. Kazantsev, Enhancement of a magnetic field by a conducting fluid. *Sov. Phys. JETP* **26**, 1031 (1968)
- U. Keshet, A. Loeb, Using radio halos and minihalos to measure the distributions of magnetic fields and cosmic rays in galaxy clusters. *Astrophys. J.* **722**, 737–749 (2010). <https://doi.org/10.1088/0004-637X/722/1/737>. arXiv:1003.1133
- R. Khatri, M. Gaspari, Thermal SZ fluctuations in the ICM: probing turbulence and thermodynamics in Coma cluster with Planck. *Mon. Not. R. Astron. Soc.* **463**, 655–669 (2016). <https://doi.org/10.1093/mnras/stw2027>. arXiv:1604.03106
- U. Klein, A. Fletcher, *Galactic and Intergalactic Magnetic Fields* (2015)
- A. Kolmogorov, The local structure of turbulence in incompressible viscous fluid for very large Reynolds' numbers. *Dokl. Akad. Nauk SSSR* **30**, 301–305 (1941)
- A.N. Kolmogorov, The local structure of turbulence in incompressible viscous fluid for very large Reynolds numbers. *Proc. R. Soc. Lond. Ser. A, Math. Phys. Sci.* **434**, 9–13 (1991). <https://doi.org/10.1098/rspa.1991.0075>
- R.P. Kraft, E. Roediger, M. Machacek, W.R. Forman, P.E.J. Nulsen, C. Jones, E. Churazov, S. Randall, Y. Su, A. Sheardown, Stripped elliptical galaxies as probes of ICM physics. III. Deep Chandra observations of NGC 4552: measuring the viscosity of the intracluster medium. *Astrophys. J.* **848**, 27 (2017). <https://doi.org/10.3847/1538-4357/aa8a6e>. arXiv:1709.02784
- R.H. Kraichnan, S. Nagarajan, Growth of turbulent magnetic fields. *Phys. Fluids* **10**, 859–870 (1967). <https://doi.org/10.1063/1.1762201>
- A.G. Kritsuk, Å Nordlund, D. Collins, P. Padoan, M.L. Norman, T. Abel, R. Banerjee, C. Federrath, M. Flock, D. Lee, P.S. Li, W.C. Müller, R. Teyssier, S.D. Ustyugov, C. Vogel, H. Xu, Comparing numerical methods for isothermal magnetized supersonic turbulence. *Astrophys. J.* **737**, 13 (2011). <https://doi.org/10.1088/0004-637X/737/1/13>. arXiv:1103.5525
- P.P. Kronberg, H. Lesch, U. Hopp, Magnetization of the intergalactic medium by primeval galaxies. *Astrophys. J.* **511**, 56–64 (1999)
- P. Kuchar, T.A. Enßlin, Magnetic power spectra from Faraday rotation maps. REALMAF and its use on Hydra A. *Astron. Astrophys.* **529**, A13 (2011). <https://doi.org/10.1051/0004-6361/200913918>. arXiv:0912.3930
- R.M. Kulsrud, S.W. Anderson, The spectrum of random magnetic fields in the mean field dynamo theory of the Galactic magnetic field. *Astrophys. J.* **396**, 606–630 (1992). <https://doi.org/10.1086/171743>
- R.M. Kulsrud, E.C. Ostriker, Plasma physics for astrophysics. *Phys. Today* **59**(1), 58 (2006). <https://doi.org/10.1063/1.2180179>
- R.M. Kulsrud, R. Cen, J.P. Ostriker, D. Ryu, The protogalactic origin for cosmic magnetic fields. *Astrophys. J.* **480**, 481–491 (1997). <https://doi.org/10.1086/303987>. arXiv:astro-ph/9607141
- M.W. Kunz, Dynamical stability of a thermally stratified intracluster medium with anisotropic momentum and heat transport. *Mon. Not. R. Astron. Soc.* **417**, 602–616 (2011). <https://doi.org/10.1111/j.1365-2966.2011.19303.x>. arXiv:1104.3595
- M.W. Kunz, A.A. Schekochihin, J.M. Stone, Firehose and mirror instabilities in a collisionless shearing plasma. *Phys. Rev. Lett.* **112**(20), 205003 (2014). <https://doi.org/10.1103/PhysRevLett.112.205003>. arXiv:1402.0010
- R.A. Laing, A.H. Bridle, P. Parma, M. Murgia, Structures of the magnetoionic media around the Fanaroff-Riley Class I radio galaxies 3C31 and Hydra A. *Mon. Not. R. Astron. Soc.* **391**, 521–549 (2008). <https://doi.org/10.1111/j.1365-2966.2008.13895.x>. arXiv:0809.2411
- L.D. Landau, E.M. Lifshitz *Hydrodynamik* (1966)
- L.D. Landau, E.M. Lifshitz, A.L. King, Electrodynamics of continuous media. *Am. J. Phys.* **29**, 647–648 (1961). <https://doi.org/10.1119/1.1937882>
- M. Langer, N. Aghanim, J.L. Puget, Magnetic fields from reionisation. *Astron. Astrophys.* **443**, 367–372 (2005). <https://doi.org/10.1051/0004-6361:20053372>. arXiv:astro-ph/0508173
- M.A. Latif, D.R.G. Schleicher, W. Schmidt, J. Niemeyer, The small-scale dynamo and the amplification of magnetic fields in massive primordial haloes. *Mon. Not. R. Astron. Soc.* **432**, 668–678 (2013). <https://doi.org/10.1093/mnras/stt503>. arXiv:1212.1619

- E.T. Lau, A.V. Kravtsov, D. Nagai, Residual gas motions in the intracluster medium and bias in hydrostatic measurements of mass profiles of clusters. *Astrophys. J.* **705**, 1129–1138 (2009). <https://doi.org/10.1088/0004-637X/705/2/1129>. arXiv:0903.4895
- A. Lazarian, A. Beresnyak, H. Yan, M. Opher, Y. Liu, Properties and selected implications of magnetic turbulence for interstellar medium, local bubble and solar wind. *Space Sci. Rev.* **143**, 387–413 (2009). <https://doi.org/10.1007/s11214-008-9452-y>. arXiv:0811.0826
- A. Lazarian, L. Vlahos, G. Kowal, H. Yan, A. Beresnyak, E.M. de Gouveia Dal Pino, Turbulence, magnetic reconnection in turbulent fluids and energetic particle acceleration. *Space Sci. Rev.* **173**, 557–622 (2012). <https://doi.org/10.1007/s11214-012-9936-7>. arXiv:1211.0008
- D. Lee, A solution accurate, efficient and stable unsplit staggered mesh scheme for three dimensional magnetohydrodynamics. *J. Comput. Phys.* **243**, 269–292 (2013). <https://doi.org/10.1016/j.jcp.2013.02.049>. arXiv:1303.6988
- S. Li, H. Li, R. Cen, CosmoMHD: a cosmological magnetohydrodynamics code. *Astrophys. J. Suppl. Ser.* **174**, 1–12 (2008). <https://doi.org/10.1086/521302>. arXiv:astro-ph/0611863
- M.S. Longair, *High Energy Astrophysics* (2011)
- S.G. Lucek, A.R. Bell, Non-linear amplification of a magnetic field driven by cosmic ray streaming. *Mon. Not. R. Astron. Soc.* **314**, 65–74 (2000). <https://doi.org/10.1046/j.1365-8711.2000.03363.x>
- M. Lyutikov, Magnetic draping of merging cores and radio bubbles in clusters of galaxies. *Mon. Not. R. Astron. Soc.* **373**, 73–78 (2006). <https://doi.org/10.1111/j.1365-2966.2006.10835.x>. arXiv:astro-ph/0604178
- A. Maier, L. Iapichino, W. Schmidt, J.C. Niemeyer, Adaptively refined large eddy simulations of a galaxy cluster: turbulence modeling and the physics of the intracluster medium. *Astrophys. J.* **707**, 40–54 (2009). <https://doi.org/10.1088/0004-637X/707/1/40>. arXiv:0909.1800
- M.A. Malkov, L. O’C Drury, Nonlinear theory of diffusive acceleration of particles by shock waves. *Rep. Prog. Phys.* **64**, 429–481 (2001). <https://doi.org/10.1088/0034-4885/64/4/201>
- A. Marcowith, A. Bret, A. Bykov, M.E. Dieckman, L. O’C Drury, B. Lembège, M. Lemoine, G. Morlino, G. Murphy, G. Pelletier, I. Plotnikov, B. Reville, M. Riquelme, L. Sironi, A. Stockem Novo, The microphysics of collisionless shock waves. *Rep. Prog. Phys.* **79**(4), 046901 (2016). <https://doi.org/10.1088/0034-4885/79/4/046901>. arXiv:1604.00318
- F. Marinacci, M. Vogelsberger, P. Mocz, R. Pakmor, The large-scale properties of simulated cosmological magnetic fields. *Mon. Not. R. Astron. Soc.* **453**, 3999–4019 (2015). <https://doi.org/10.1093/mnras/stv1692>. arXiv:1506.00005
- F. Marinacci, M. Vogelsberger, R. Kannan, P. Mocz, R. Pakmor, V. Springel, Non-ideal magnetohydrodynamics on a moving mesh. *Mon. Not. R. Astron. Soc.* **476**, 2476–2492 (2018a). <https://doi.org/10.1093/mnras/sty397>. arXiv:1710.10265
- F. Marinacci, M. Vogelsberger, R. Pakmor, P. Torrey, V. Springel, L. Hernquist, D. Nelson, R. Weinberger, A. Pillepich, J. Naiman, S. Genel, First results from the IllustrisTNG simulations: radio haloes and magnetic fields. *Mon. Not. R. Astron. Soc.* **480**, 5113–5139 (2018b). <https://doi.org/10.1093/mnras/sty2206>. arXiv:1707.03396
- M. Markevitch, A. Vikhlinin, *Phys. Rep.* **443**, 1–53 (2007). <https://doi.org/10.1016/j.physrep.2007.01.001>. arXiv:astro-ph/0701821
- S. Martín-Alvarez, J. Devriendt, A. Slyz, R. Teyssier, A three-phase amplification of the cosmic magnetic field in galaxies. *Mon. Not. R. Astron. Soc.* **479**(3), 3343–3365 (2018). <https://doi.org/10.1093/mnras/sty1623>. arXiv:1806.06866
- P. Mazzotta, S. Giacintucci, Do radio core-halos and cold fronts in non-major-merging clusters originate from the same gas sloshing? *Astrophys. J. Lett.* **675**, L9 (2008). <https://doi.org/10.1086/529433>. arXiv:0801.1905
- J. Meinecke, P. Tzeferacos, A. Bell, R. Bingham, R. Clarke, E. Churazov, R. Crowston, H. Doyle, R.P. Drake, R. Heathcote, M. Koenig, Y. Kuramitsu, C. Kuranz, D. Lee, M. MacDonald, C. Murphy, M. Notley, H.S. Park, A. Pelka, A. Rivasio, B. Reville, Y. Sakawa, W. Wan, N. Woolsey, R. Yurchak, F. Miniati, A. Schekochihin, D. Lamb, G. Gregori, Developed turbulence and nonlinear amplification of magnetic fields in laboratory and astrophysical plasmas. *Proc. Natl. Acad. Sci. USA* **112**, 8211–8215 (2015). <https://doi.org/10.1073/pnas.1502079112>
- S. Melville, A.A. Schekochihin, M.W. Kunz, Pressure-anisotropy-driven microturbulence and magnetic-field evolution in shearing, collisionless plasma. *Mon. Not. R. Astron. Soc.* **459**, 2701–2720 (2016). <https://doi.org/10.1093/mnras/stw793>. arXiv:1512.08131
- F. Miniati, The Matryoshka Run: a Eulerian refinement strategy to study the statistics of turbulence in virialized cosmic structures. *Astrophys. J.* **782**, 21 (2014). <https://doi.org/10.1088/0004-637X/782/1/21>. arXiv:1310.2951

- F. Miniati, The Matryoshka Run. II. Time-dependent turbulence statistics, stochastic particle acceleration, and microphysics impact in a massive galaxy cluster. *Astrophys. J.* **800**, 60 (2015). <https://doi.org/10.1088/0004-637X/800/1/60>. arXiv:1409.3576
- F. Miniati, A.R. Bell, Resistive magnetic field generation at cosmic dawn. *Astrophys. J.* **729**, 73 (2011). <https://doi.org/10.1088/0004-637X/729/1/73>. arXiv:1001.2011
- F. Miniati, A. Beresnyak, Self-similar energetics in large clusters of galaxies. *Nature* **523**, 59–62 (2015). <https://doi.org/10.1038/nature14552>. arXiv:1507.01940
- F. Miniati, D. Ryu, H. Kang, T.W. Jones, R. Cen, J.P. Ostriker, Properties of cosmic shock waves in large-scale structure formation. *Astrophys. J.* **542**, 608–621 (2000). <https://doi.org/10.1086/317027>. arXiv:astro-ph/0005444
- N.L. Mitchell, I.G. McCarthy, R.G. Bower, T. Theuns, R.A. Crain, On the origin of cores in simulated galaxy clusters. *Mon. Not. R. Astron. Soc.* **395**, 180–196 (2009). <https://doi.org/10.1111/j.1365-2966.2009.14550.x>. arXiv:0812.1750
- H. Mo, F.C. van den Bosch, S. White *Galaxy Formation and Evolution* (2010)
- F. Mogavero, A.A. Schekochihin, Models of magnetic field evolution and effective viscosity in weakly collisional extragalactic plasmas. *Mon. Not. R. Astron. Soc.* **440**, 3226–3242 (2014). <https://doi.org/10.1093/mnras/stu433>. arXiv:1312.3672
- T. Naab, J.P. Ostriker, Theoretical challenges in galaxy formation. *Annu. Rev. Astron. Astrophys.* **55**, 59–109 (2017). <https://doi.org/10.1146/annurev-astro-081913-040019>. arXiv:1612.06891
- D. Nagai, A.V. Kravtsov, A. Vikhlinin, Effects of galaxy formation on thermodynamics of the intracluster medium. *Astrophys. J.* **668**, 1–14 (2007). <https://doi.org/10.1086/521328>. arXiv:astro-ph/0703661
- K. Nakazawa, C.L. Sarazin, M. Kawaharada, T. Kitaguchi, S. Okuyama, K. Makishima, N. Kawano, Y. Fukazawa, S. Inoue, M. Takizawa, D.R. Wik, A. Finoguenov, T.E. Clarke, Hard X-ray properties of the merging cluster Abell 3667 as observed with Suzaku. *Publ. Astron. Soc. Jpn.* **61**, 339–355 (2009). <https://doi.org/10.1093/pasj/61.2.339>. arXiv:0812.1438
- K. Nelson, E.T. Lau, D. Nagai, Hydrodynamic simulation of non-thermal pressure profiles of galaxy clusters. *Astrophys. J.* **792**, 25 (2014). <https://doi.org/10.1088/0004-637X/792/1/25>. arXiv:1404.4636
- A. Neronov, I. Vovk, Evidence for strong extragalactic magnetic fields from Fermi observations of TeV blazars. *Science* **328**, 73 (2010). <https://doi.org/10.1126/science.1184192>. arXiv:1006.3504
- F.N. Owen, L. Rudnick, J. Eilek, U. Rau, S. Bhatnagar, L. Kogan, Wideband Very Large Array Observations of A2256. I. Continuum, rotation measure, and spectral imaging. *Astrophys. J.* **794**, 24 (2014). <https://doi.org/10.1088/0004-637X/794/1/24>. arXiv:1408.5931
- R. Pakmor, A. Bauer, V. Springel, Magnetohydrodynamics on an unstructured moving grid. *Mon. Not. R. Astron. Soc.* **418**, 1392–1401 (2011). <https://doi.org/10.1111/j.1365-2966.2011.19591.x>. arXiv:1108.1792
- R. Pakmor, F.A. Gómez, R.J.J. Grand, F. Marinacci, C.M. Simpson, V. Springel, D.J.R. Campbell, C.S. Frenk, T. Guillet, C. Pfrommer, S.D.M. White, Magnetic field formation in the Milky Way like disc galaxies of the Auriga project. *Mon. Not. R. Astron. Soc.* **469**, 3185–3199 (2017). <https://doi.org/10.1093/mnras/stx1074>. arXiv:1701.07028
- E. Parizot, A. Marcowith, J. Ballet, Y.A. Gallant, Observational constraints on energetic particle diffusion in young supernovae remnants: amplified magnetic field and maximum energy. *Astron. Astrophys.* **453**, 387–395 (2006). <https://doi.org/10.1051/0004-6361:20064985>. arXiv:astro-ph/0603723
- S. Paul, L. Iapichino, F. Miniati, J. Bagchi, K. Mannheim, Evolution of shocks and turbulence in major cluster mergers. *Astrophys. J.* **726**, 17 (2011). <https://doi.org/10.1088/0004-637X/726/1/17>. arXiv:1001.1170
- C. Pfrommer, Toward a comprehensive model for feedback by active galactic nuclei: new insights from M87 observations by LOFAR, Fermi, and H.E.S.S. *Astrophys. J.* **779**, 10 (2013). <https://doi.org/10.1088/0004-637X/779/1/10>. arXiv:1303.5443
- C. Pfrommer, T.A. Enßlin, Constraining the population of cosmic ray protons in cooling flow clusters with  $\gamma$ -ray and radio observations: are radio mini-halos of hadronic origin? *Astron. Astrophys.* **413**, 17–36 (2004). <https://doi.org/10.1051/0004-6361:20031464>
- C. Pfrommer, V. Springel, T.A. Enßlin, M. Jubelgas, *Mon. Not. R. Astron. Soc.* **367**, 113–131 (2006). <https://doi.org/10.1111/j.1365-2966.2005.09953.x>. arXiv:astro-ph/0603483
- S. Planelles, D.R.G. Schleicher, A.M. Bykov, Large-scale structure formation: from the first non-linear objects to massive galaxy clusters. *Space Sci. Rev.* **188**, 93–139 (2015). <https://doi.org/10.1007/s11214-014-0045-7>. arXiv:1404.3956
- S. Planelles, D.R.G. Schleicher, A.M. Bykov, Large-scale structure formation: from the first non-linear objects to massive galaxy clusters, in *Multi-scale Structure Formation and Dynamics in Cosmic Plasmas* (2016), pp. 93–139. [https://doi.org/10.1007/978-1-4939-3547-5\\_4](https://doi.org/10.1007/978-1-4939-3547-5_4)
- D.H. Porter, T.W. Jones, D. Ryu, Vorticity, shocks, and magnetic fields in subsonic, ICM-like turbulence. *Astrophys. J.* **810**, 93 (2015). <https://doi.org/10.1088/0004-637X/810/2/93>. arXiv:1507.08737

- D.J. Price, Resolving high Reynolds numbers in smoothed particle hydrodynamics simulations of subsonic turbulence. *Mon. Not. R. Astron. Soc.* **420**, L33–L37 (2012a). <https://doi.org/10.1111/j.1745-3933.2011.01187.x>. arXiv:1111.1255
- D.J. Price, Smoothed particle hydrodynamics and magnetohydrodynamics. *J. Comput. Phys.* **231**, 759–794 (2012b). <https://doi.org/10.1016/j.jcp.2010.12.011>. arXiv:1012.1885
- K. Rajpurohit, M. Hoeft, R.J. van Weeren, L. Rudnick, H.J.A. Röttgering, W.R. Forman, M. Brüggen, J.H. Croston, F. Andrade-Santos, W.A. Dawson, H.T. Intema, R.P. Kraft, C. Jones, M.J. Jee, Deep VLA observations of the cluster 1RXS J0603.3+4214 in the frequency range of 1–2 GHz. *Astrophys. J.* **852**, 65 (2018). <https://doi.org/10.3847/1538-4357/aa9f13>. arXiv:1712.01327
- B. Reville, A.R. Bell, Universal behaviour of shock precursors in the presence of efficient cosmic ray acceleration. *Mon. Not. R. Astron. Soc.* **430**, 2873–2884 (2013). <https://doi.org/10.1093/mnras/stt100>. arXiv:1301.3173
- B. Reville, S. O’Sullivan, P. Duffy, J.G. Kirk, The transport of cosmic rays in self-excited magnetic turbulence. *Mon. Not. R. Astron. Soc.* **386**, 509–515 (2008). <https://doi.org/10.1111/j.1365-2966.2008.13059.x>. arXiv:0802.0109
- S.P. Reynolds, Models of synchrotron X-rays from shell supernova remnants. *Astrophys. J.* **493**, 375–396 (1998). <https://doi.org/10.1086/305103>
- P.M. Ricker, C.L. Sarazin, Off-axis cluster mergers: effects of a strongly peaked dark matter profile. *Astrophys. J.* **561**, 621–644 (2001)
- M. Rieder, R. Teyssier, A small-scale dynamo in feedback-dominated galaxies as the origin of cosmic magnetic fields—I. The kinematic phase. *Mon. Not. R. Astron. Soc.* **457**, 1722–1738 (2016). <https://doi.org/10.1093/mnras/stv2985>. arXiv:1506.00849
- F. Rincon, F. Califano, A.A. Schekochihin, F. Valentini, Turbulent dynamo in a collisionless plasma. *Proc. Natl. Acad. Sci. USA* **113**, 3950–3953 (2016). <https://doi.org/10.1073/pnas.1525194113>. arXiv:1512.06455
- M.A. Riquelme, A. Spitkovsky, Magnetic amplification by magnetized cosmic rays in supernova remnant shocks. *Astrophys. J.* **717**, 1054–1066 (2010). <https://doi.org/10.1088/0004-637X/717/2/1054>. arXiv:0912.4990
- M.A. Riquelme, A. Spitkovsky, Electron injection by whistler waves in non-relativistic shocks. *Astrophys. J.* **733**, 63 (2011). <https://doi.org/10.1088/0004-637X/733/1/63>. arXiv:1009.3319
- B.E. Robertson, A.V. Kravtsov, N.Y. Gnedin, T. Abel, D.H. Rudd, Computational Eulerian hydrodynamics and Galilean invariance. *Mon. Not. R. Astron. Soc.* **401**, 2463–2476 (2010). <https://doi.org/10.1111/j.1365-2966.2009.15823.x>. arXiv:0909.0513
- E. Roediger, R.P. Kraft, P.E.J. Nulsen, W.R. Forman, M. Machacek, S. Randall, C. Jones, E. Churazov, R. Kokotanekova, Stripped elliptical galaxies as probes of ICM physics: I. Tails, wakes, and flow patterns in and around stripped ellipticals. *Astrophys. J.* **806**, 103 (2015a). <https://doi.org/10.1088/0004-637X/806/1/103>. arXiv:1409.6300
- E. Roediger, R.P. Kraft, P.E.J. Nulsen, W.R. Forman, M. Machacek, S. Randall, C. Jones, E. Churazov, R. Kokotanekova, Stripped elliptical galaxies as probes of ICM physics: II. Stirred, but mixed? Viscous and inviscid gas stripping of the Virgo elliptical M89. *Astrophys. J.* **806**, 104 (2015b). <https://doi.org/10.1088/0004-637X/806/1/104>. arXiv:1409.6312
- K. Roettiger, J. Burns, C. Loken, When clusters collide—a numerical hydro/N-body simulation of merging galaxy clusters. *Astrophys. J. Lett.* **407**, L53–L56 (1993). <https://doi.org/10.1086/186804>
- K. Roettiger, C. Loken, J.O. Burns, Numerical simulations of merging clusters of galaxies. *Astrophys. J. Suppl. Ser.* **109**, 307 (1997)
- K. Roettiger, J.M. Stone, J.O. Burns, Magnetic field evolution in merging clusters of galaxies. *Astrophys. J.* **518**, 594–602 (1999)
- J. Roland, On the origin of the intergalactic magnetic field and of the radio halo associated with the Coma cluster of galaxies. *Astron. Astrophys.* **93**, 407–410 (1981)
- M. Roncarelli, M. Gaspari, S. Ettori, V. Biffi, F. Brighenti, E. Bulbul, N. Clerc, E. Cucchetti, E. Pointecouteau, E. Rasia, Measuring turbulence and gas motions in galaxy clusters via synthetic *Athena* X-IFU observations. arXiv:1805.02577 (2018)
- M. Ruszkowski, S.P. Oh, Galaxy motions, turbulence and conduction in clusters of galaxies. *Mon. Not. R. Astron. Soc.* **414**, 1493–1507 (2011). <https://doi.org/10.1111/j.1365-2966.2011.18482.x>. arXiv:1008.5016
- M. Ruszkowski, D. Lee, M. Brüggen, I. Parrish, S.P. Oh, Cosmological magnetohydrodynamic simulations of cluster formation with anisotropic thermal conduction. *Astrophys. J.* **740**, 81 (2011). <https://doi.org/10.1088/0004-637X/740/2/81>. arXiv:1010.2277
- A. Ruzmaikin, D. Sokolov, A. Shukurov, The dynamo origin of magnetic fields in galaxy clusters. *Mon. Not. R. Astron. Soc.* **241**, 1–14 (1989). <https://doi.org/10.1093/mnras/241.1.1>
- D. Ryu, H. Kang, E. Hallman, T.W. Jones, *Astrophys. J.* **593**, 599–610 (2003). <https://doi.org/10.1086/376723>. arXiv:astro-ph/0305164



- D. Ryu, H. Kang, J. Cho, S. Das, Turbulence and magnetic fields in the large-scale structure of the universe. *Science* **320**, 909 (2008). <https://doi.org/10.1126/science.1154923>. arXiv:0805.2466
- D. Ryu, D.R.G. Schleicher, R.A. Treumann, C.G. Tsagas, L.M. Widrow, Magnetic fields in the large-scale structure of the universe. *Space Sci. Rev.* **166**, 1–35 (2012). <https://doi.org/10.1007/s11214-011-9839-z>. arXiv:1109.4055
- S. Samui, K. Subramanian, R. Srianand, Efficient cold outflows driven by cosmic rays in high redshift galaxies and their global effects on the IGM. arXiv:1706.01890 (2017)
- F.J. Sánchez-Salcedo, A. Brandenburg, A. Shukurov, Turbulence and magnetic fields in clusters of galaxies. *Astrophys. Space Sci.* **263**, 87–90 (1998). <https://doi.org/10.1023/A:1002144413095>
- T. Sano, K. Nishihara, C. Matsuoka, T. Inoue, Magnetic field amplification associated with the Richtmyer-Meshkov instability. *Astrophys. J.* **758**, 126 (2012). <https://doi.org/10.1088/0004-637X/758/2/126>. arXiv:1209.0961
- R. Santos-Lima, E.M. de Gouveia Dal Pino, G. Kowal, D. Falceta-Gonçalves, A. Lazarian, M.S. Nakwacki, Magnetic field amplification and evolution in turbulent collisionless magnetohydrodynamics: an application to the intracluster medium. *Astrophys. J.* **781**, 84 (2014). <https://doi.org/10.1088/0004-637X/781/2/84>. arXiv:1305.5654
- R. Santos-Lima, E.M. de Gouveia Dal Pino, D.A. Falceta-Gonçalves, M.S. Nakwacki, G. Kowal, Features of collisionless turbulence in the intracluster medium from simulated Faraday rotation maps—II. The effects of instabilities feedback. *Mon. Not. R. Astron. Soc.* **465**, 4866–4871 (2017). <https://doi.org/10.1093/mnras/stw3050>. arXiv:1611.10183
- C.L. Sarazin, X-ray emission from clusters of galaxies. *Rev. Mod. Phys.* **58**, 1 (1986)
- C.L. Sarazin, *X-Ray Emission from Clusters of Galaxies* (Cambridge University Press, Cambridge, 1988)
- C.L. Sarazin, The physics of cluster mergers, in *Merging Processes in Galaxy Clusters*. ASSL, vol. 272 (2002), pp. 1–38
- A.A. Schekochihin, S.C. Cowley, *Turbulence and Magnetic Fields in Astrophysical Plasmas* (Springer, Berlin, 2007), p. 85
- A. Schekochihin, S. Cowley, J. Maron, L. Malyshkin, Structure of small-scale magnetic fields in the kinematic dynamo theory. *Phys. Rev. E* **65**(1), 016305 (2002a). <https://doi.org/10.1103/PhysRevE.65.016305>. arXiv:astro-ph/0105322
- A.A. Schekochihin, J.L. Maron, S.C. Cowley, J.C. McWilliams, The small-scale structure of magnetohydrodynamic turbulence with large magnetic Prandtl numbers. *Astrophys. J.* **576**, 806–813 (2002b). <https://doi.org/10.1086/341814>. arXiv:astro-ph/0203219
- A.A. Schekochihin, S.C. Cowley, S.F. Taylor, J.L. Maron, J.C. McWilliams, Simulations of the small-scale turbulent dynamo. *Astrophys. J.* **612**, 276–307 (2004). <https://doi.org/10.1086/422547>. arXiv:astro-ph/0312046
- A. Schekochihin, S. Cowley, R. Kulsrud, G. Hammett, P. Sharma, Magnetised plasma turbulence in clusters of galaxies, in *The Magnetized Plasma in Galaxy Evolution*, ed. by K.T. Chyzy, K. Otmianowska-Mazur, M. Soida, R.J. Dettmar (2005a), pp. 86–92. arXiv:astro-ph/0411781
- A.A. Schekochihin, S.C. Cowley, R.M. Kulsrud, G.W. Hammett, P. Sharma, Plasma instabilities and magnetic field growth in clusters of galaxies. *Astrophys. J.* **629**, 139–142 (2005b). <https://doi.org/10.1086/431202>. arXiv:astro-ph/0501362
- A.A. Schekochihin, S.C. Cowley, R.M. Kulsrud, M.S. Rosin, T. Heinemann, Nonlinear growth of firehose and mirror fluctuations in astrophysical plasmas. *Phys. Rev. Lett.* **100**(8), 081301 (2008). <https://doi.org/10.1103/PhysRevLett.100.081301>. arXiv:0709.3828
- A.A. Schekochihin, S.C. Cowley, W. Dorland, G.W. Hammett, G.G. Howes, E. Quataert, T. Tatsuno, Astrophysical gyrokinetics: kinetic and fluid turbulent cascades in magnetized weakly collisional plasmas. *Astrophys. J. Suppl. Ser.* **182**, 310–377 (2009). <https://doi.org/10.1088/0067-0049/182/1/310>. arXiv:0704.0044
- S. Schindler, E. Mueller, Simulations of the evolution of galaxy clusters. 11. Dynamics of the intra-cluster gas. *Astron. Astrophys.* **272**, 137 (1993)
- D.R.G. Schleicher, R. Banerjee, S. Sur, T.G. Arshakian, R.S. Klessen, R. Beck, M. Spaans, Small-scale dynamo action during the formation of the first stars and galaxies. I. The ideal MHD limit. *Astron. Astrophys.* **522**, A115 (2010). <https://doi.org/10.1051/0004-6361/201015184>. arXiv:1003.1135
- D.R.G. Schleicher, J. Schober, C. Federrath, S. Bovino, W. Schmidt, The small-scale dynamo: breaking universality at high Mach numbers. *New J. Phys.* **15**(2), 023017 (2013). <https://doi.org/10.1088/1367-2630/15/2/023017>. arXiv:1301.4371
- R. Schlickeiser, *Cosmic Ray Astrophysics* (2002)
- R. Schlickeiser, D. Ibscher, M. Supsar, Plasma effects on fast pair beams in cosmic voids. *Astrophys. J.* **758**, 102 (2012). <https://doi.org/10.1088/0004-637X/758/2/102>



- W. Schmidt, C. Byrohl, J.F. Engels, C. Behrens, J.C. Niemeyer, Viscosity, pressure and support of the gas in simulations of merging cool-core clusters. *Mon. Not. R. Astron. Soc.* **470**, 142–156 (2017). <https://doi.org/10.1093/mnras/stx1274>. arXiv:1705.06933
- J. Schober, D.R.G. Schleicher, R.S. Klessen, Magnetic field amplification in young galaxies. *Astron. Astrophys.* **560**, A87 (2013). <https://doi.org/10.1051/0004-6361/201322185>. arXiv:1310.0853
- P. Schuecker, A. Finoguenov, F. Miniati, H. Böhringer, U.G. Briel, Probing turbulence in the Coma galaxy cluster. *Astron. Astrophys.* **426**, 387–397 (2004). <https://doi.org/10.1051/0004-6361:20041039>. arXiv:astro-ph/0404132
- V.B. Semikoz, D. Sokoloff, Magnetic helicity and cosmological magnetic field. *Astron. Astrophys.* **433**, L53–L56 (2005). <https://doi.org/10.1051/0004-6361:200500094>. arXiv:astro-ph/0411496
- L.D. Shaw, D. Nagai, S. Bhattacharya, E.T. Lau, Impact of cluster physics on the Sunyaev-Zel'dovich power spectrum. *Astrophys. J.* **725**, 1452–1465 (2010). <https://doi.org/10.1088/0004-637X/725/2/1452>. arXiv:1006.1945
- X. Shi, E. Komatsu, Analytical model for non-thermal pressure in galaxy clusters. *Mon. Not. R. Astron. Soc.* **442**, 521–532 (2014). <https://doi.org/10.1093/mnras/stu858>. arXiv:1401.7657
- X. Shi, E. Komatsu, K. Nelson, D. Nagai, Analytical model for non-thermal pressure in galaxy clusters—II. Comparison with cosmological hydrodynamics simulation. *Mon. Not. R. Astron. Soc.* **448**, 1020–1029 (2015). <https://doi.org/10.1093/mnras/stv036>. arXiv:1408.3832
- X. Shi, E. Komatsu, D. Nagai, E.T. Lau, Analytical model for non-thermal pressure in galaxy clusters—III. Removing the hydrostatic mass bias. *Mon. Not. R. Astron. Soc.* **455**, 2936–2944 (2016). <https://doi.org/10.1093/mnras/stv2504>. arXiv:1507.04338
- T.W. Shimwell, H.J.A. Röttgering, P.N. Best, W.L. Williams, T.J. Dijkema, F. de Gasperin, M.J. Hardcastle, G.H. Heald, D.N. Hoang, A. Horneffer, H. Intema, E.K. Mahony, S. Mandal, A.P. Mechev, L. Morabito, J.B.R. Oonk, D. Rafferty, E. Retana-Montenegro, J. Sabater, C. Tasse, R.J. van Weeren, M. Brügger, G. Brunetti, K.T. Chyzy, J.E. Conway, M. Haverkorn, N. Jackson, M.J. Jarvis, J.P. McKean, G.K. Miley, R. Morganti, G.J. White, M.W. Wise, I.M. van Bemmel, R. Beck, M. Brienza, A. Bonafede, G. Calistro Rivera, R. Cassano, A.O. Clarke, D. Cseh, A. Deller, A. Drabant, W. van Driel, D. Engels, H. Falcke, C. Ferrari, S. Fröhlich, M.A. Garrett, J.J. Harwood, V. Heesen, M. Hoeft, C. Horellou, F.P. Israel, A.D. Kapińska, M. Kunert-Bajraszewska, D.J. McKay, N.R. Mohan, E. Orrù, R.F. Pizzo, I. Prandoni, D.J. Schwarz, A. Shulevski, M. Sipior, D.J.B. Smith, S.S. Sridhar, M. Steinmetz, A. Stroe, E. Varenius, P.P. van der Werf, J.A. Zensus, J.T.L. Zwart, The LOFAR Two-metre Sky Survey. I. Survey description and preliminary data release. *Astron. Astrophys.* **598**, A104 (2017). <https://doi.org/10.1051/0004-6361/201629313>. arXiv:1611.02700
- S.W. Skillman, H. Xu, E.J. Hallman, B.W. O'Shea, J.O. Burns, H. Li, D.C. Collins, M.L. Norman, Cosmological magnetohydrodynamic simulations of galaxy cluster radio relics: insights and warnings for observations. *Astrophys. J.* **765**, 21 (2013). <https://doi.org/10.1088/0004-637X/765/1/21>. arXiv:1211.3122
- P. Slane, S.H. Lee, D.C. Ellison, D.J. Patnaude, J.P. Hughes, K.A. Eriksen, D. Castro, S. Nagataki, A CR-hydro-NEI model of the structure and broadband emission from Tycho's supernova remnant. *Astrophys. J.* **783**, 33 (2014). <https://doi.org/10.1088/0004-637X/783/1/33>
- L. Spitzer, *Physics of Fully Ionized Gases* (1956)
- V. Springel, *E pur si muove*: Galilean-invariant cosmological hydrodynamical simulations on a moving mesh. *Mon. Not. R. Astron. Soc.* **401**, 791–851 (2010). <https://doi.org/10.1111/j.1365-2966.2009.15715.x>. arXiv:0901.4107
- F.A. Stasyszyn, D. Elstner, A vector potential implementation for smoothed particle magnetohydrodynamics. *J. Comput. Phys.* **282**, 148–156 (2015). <https://doi.org/10.1016/j.jcp.2014.11.011>. arXiv:1411.3290
- U.P. Steinwandel, M.C. Beck, A. Arth, K. Dolag, B.P. Moster, P. Nielaba, Magnetic buoyancy in simulated galactic discs with a realistic circum galactic medium. arXiv:1808.09975 (2018)
- A. Stroe, R.J. van Weeren, H.T. Intema, H.J.A. Röttgering, M. Brügger, M. Hoeft, Discovery of spectral curvature in the shock downstream region: CIZA J2242.8+5301. *Astron. Astrophys.* **555**, A110 (2013). <https://doi.org/10.1051/0004-6361/201321267>. arXiv:1305.0005
- A. Stroe, J.J. Harwood, M.J. Hardcastle, H.J.A. Röttgering, Spectral age modelling of the 'Sausage' cluster radio relic. *Mon. Not. R. Astron. Soc.* **445**, 1213–1222 (2014). <https://doi.org/10.1093/mnras/stu1839>. arXiv:1409.1579
- Y. Su, R.P. Kraft, E. Roediger, P. Nulsen, W.R. Forman, E. Churazov, S.W. Randall, C. Jones, M.E. Machacek, Deep Chandra observations of NGC 1404: cluster plasma physics revealed by an infalling early-type galaxy. *Astrophys. J.* **834**, 74 (2017). <https://doi.org/10.3847/1538-4357/834/1/74>. arXiv:1612.00535
- K. Subramanian, The origin, evolution and signatures of primordial magnetic fields. *Rep. Prog. Phys.* **79**(7), 076901 (2016). <https://doi.org/10.1088/0034-4885/79/7/076901>. arXiv:1504.02311
- K. Subramanian, A. Shukurov, N.E.L. Haugen, Evolving turbulence and magnetic fields in galaxy clusters. *Mon. Not. R. Astron. Soc.* **366**, 1437–1454 (2006). <https://doi.org/10.1111/j.1365-2966.2006.09918.x>. arXiv:astro-ph/0505144

- S. Sur, C. Federrath, D.R.G. Schleicher, R. Banerjee, R.S. Klessen, Magnetic field amplification during gravitational collapse—influence of turbulence, rotation and gravitational compression. *Mon. Not. R. Astron. Soc.* **423**, 3148–3162 (2012). <https://doi.org/10.1111/j.1365-2966.2012.21100.x>. arXiv:1202.3206
- M. Takizawa, Hydrodynamic simulations of a moving substructure in a cluster of galaxies: cold fronts and turbulence generation. *Astrophys. J.* **629**, 791–796 (2005). <https://doi.org/10.1086/431927>. arXiv:astro-ph/0505274
- R. Taylor, I. Agudo, T. Akahori, R. Beck, B. Gaensler, G. Heald, M. Johnston-Hollitt, M. Langer, L. Rudnick, A. Scaife, D. Schleicher, J. Stil, D. Ryu, SKA deep polarization and cosmic magnetism, in *Advancing Astrophysics with the Square Kilometre Array (AASKA14)* (2015), p. 113. arXiv:1501.02298
- P.A. Thomas, H.M.P. Couchman, Simulating the formation of a cluster of galaxies. *Mon. Not. R. Astron. Soc.* **257**, 11–31 (1992). <https://doi.org/10.1093/mnras/257.1.11>
- T.S. Tricco, D.J. Price, C. Federrath, A comparison between grid and particle methods on the small-scale dynamo in magnetized supersonic turbulence. *Mon. Not. R. Astron. Soc.* **461**, 1260–1275 (2016). <https://doi.org/10.1093/mnras/stw1280>. arXiv:1605.08662
- P. Trivedi, K. Subramanian, T.R. Seshadri, Primordial magnetic field limits from the CMB trispectrum: scalar modes and constraints. *Phys. Rev. Lett.* **89**(4), 043523 (2014). <https://doi.org/10.1103/PhysRevD.89.043523>. arXiv:1312.5308
- M.S. Turner, L.M. Widrow, Inflation-produced, large-scale magnetic fields. *Phys. Rev. Lett.* **37**, 2743–2754 (1988). <https://doi.org/10.1103/PhysRevD.37.2743>
- V. Vacca, M. Murgia, F. Govoni, L. Ferretti, G. Giovannini, R.A. Perley, G.B. Taylor, The intracluster magnetic field power spectrum in A2199. *Astron. Astrophys.* **540**, A38 (2012). <https://doi.org/10.1051/0004-6361/201116622>. arXiv:1201.4119
- V. Vacca, N. Oppermann, T. Enßlin, J. Jasche, M. Selig, M. Greiner, H. Junklewitz, M. Reinecke, M. Brüggen, E. Carretti, L. Ferretti, C. Ferrari, C.A. Hales, C. Horellou, S. Ideguchi, M. Johnston-Hollitt, R.F. Pizzo, H. Röttgering, T.W. Shimwell, K. Takahashi, Using rotation measure grids to detect cosmological magnetic fields: a Bayesian approach. *Astron. Astrophys.* **591**, A13 (2016). <https://doi.org/10.1051/0004-6361/201527291>. arXiv:1509.00747
- S.I. Vaĭnshtein, Y.B. Zel'dovich, Origin of magnetic fields in astrophysics (turbulent “dynamo” mechanisms). *Sov. Phys. Usp.* **15**, 159–172 (1972). <https://doi.org/10.1070/PU1972v015n02ABEH004960>
- R. Valdarnini, The impact of numerical viscosity in SPH simulations of galaxy clusters. *Astron. Astrophys.* **526**, A158 (2011). <https://doi.org/10.1051/0004-6361/201015340>. arXiv:1010.3378
- F. Vazza, G. Tormen, R. Cassano, G. Brunetti, K. Dolag, Turbulent velocity fields in smoothed particle hydrodynamics simulated galaxy clusters: scaling laws for the turbulent energy. *Mon. Not. R. Astron. Soc.* **369**, L14–L18 (2006). <https://doi.org/10.1111/j.1745-3933.2006.00164.x>. arXiv:astro-ph/0602247
- F. Vazza, G. Brunetti, A. Kritsuk, R. Wagner, C. Gheller, M. Norman, Turbulent motions and shocks waves in galaxy clusters simulated with adaptive mesh refinement. *Astron. Astrophys.* **504**, 33–43 (2009). <https://doi.org/10.1051/0004-6361/200912535>. arXiv:0905.3169
- F. Vazza, G. Brunetti, C. Gheller, R. Brunino, M. Brüggen, Massive and refined. II. The statistical properties of turbulent motions in massive galaxy clusters with high spatial resolution. *Astron. Astrophys.* **529**, A17 (2011). <https://doi.org/10.1051/0004-6361/201016015>. arXiv:1010.5950
- F. Vazza, E. Roediger, M. Brueggen, Turbulence in the ICM from mergers, cool-core sloshing and jets: results from a new multi-scale filtering approach. arXiv:1202.5882 (2012)
- F. Vazza, M. Brüggen, C. Gheller, P. Wang, On the amplification of magnetic fields in cosmic filaments and galaxy clusters. *Mon. Not. R. Astron. Soc.* **445**, 3706–3722 (2014). <https://doi.org/10.1093/mnras/stu1896>. arXiv:1409.2640
- F. Vazza, M. Brüggen, C. Gheller, S. Hackstein, D. Wittor, P.M. Hinz, Simulations of extragalactic magnetic fields and of their observables. *Class. Quantum Gravity* **34**(23), 234001 (2017a). <https://doi.org/10.1088/1361-6382/aa8e60>. arXiv:1711.02669
- F. Vazza, T.W. Jones, M. Brüggen, G. Brunetti, C. Gheller, D. Porter, D. Ryu, Turbulence and vorticity in Galaxy clusters generated by structure formation. *Mon. Not. R. Astron. Soc.* **464**, 210–230 (2017b). <https://doi.org/10.1093/mnras/stw2351>. arXiv:1609.03558
- F. Vazza, G. Brunetti, M. Brüggen, A. Bonafede, Resolved magnetic dynamo action in the simulated intracluster medium. *Mon. Not. R. Astron. Soc.* **474**, 1672–1687 (2018). <https://doi.org/10.1093/mnras/stx2830>. arXiv:1711.02673
- A.A. Vikhlinin, M.L. Markevitch, A cold front in the galaxy cluster A3667: hydrodynamics, heat conduction and magnetic field in the intergalactic medium. *Astron. Lett.* **28**, 495–508 (2002). <https://doi.org/10.1134/1.1499173>. arXiv:astro-ph/0209551
- A. Vikhlinin, M. Markevitch, S.S. Murray, Chandra estimate of the magnetic field strength near the cold front in A3667. *Astrophys. J. Lett.* **549**, L47–L50 (2001). <https://doi.org/10.1086/319126>. arXiv:astro-ph/0008499

- A. Vladimirov, D.C. Ellison, A. Bykov, Nonlinear diffusive shock acceleration with magnetic field amplification. *Astrophys. J.* **652**, 1246–1258 (2006). <https://doi.org/10.1086/508154>. arXiv:astro-ph/0606433
- C. Vogt, T.A. Enßlin, Measuring the cluster magnetic field power spectra from Faraday rotation maps of Abell 400, Abell 2634 and Hydra A. *Astron. Astrophys.* **412**, 373–385 (2003)
- G.M. Voit, Tracing cosmic evolution with clusters of galaxies. *Rev. Mod. Phys.* **77**, 207–258 (2005). <https://doi.org/10.1103/RevModPhys.77.207>. arXiv:astro-ph/0410173
- H.J. Völk, A.M. Atoyan, Early starbursts and magnetic field generation in galaxy clusters. *Astrophys. J.* **541**, 88–94 (2000)
- K. Waagan, C. Federrath, C. Klingenberg, A robust numerical scheme for highly compressible magnetohydrodynamics: Nonlinear stability, implementation and tests. *J. Comput. Phys.* **230**, 3331–3351 (2011). <https://doi.org/10.1016/j.jcp.2011.01.026>. arXiv:1101.3007
- S.A. Walker, J. Hlavacek-Larrondo, M. Gendron-Marsolais, A.C. Fabian, H. Intema, J.S. Sanders, J.T. Bamford, R. van Weeren, Is there a giant Kelvin-Helmholtz instability in the sloshing cold front of the Perseus cluster? *Mon. Not. R. Astron. Soc.* **468**, 2506–2516 (2017). <https://doi.org/10.1093/mnras/stx640>. arXiv:1705.00011
- N. Werner, J.A. ZuHone, I. Zhuravleva, Y. Ichinohe, A. Simionescu, S.W. Allen, M. Markevitch, A.C. Fabian, U. Keshet, E. Roediger, M. Ruszkowski, J.S. Sanders, Deep Chandra observation and numerical studies of the nearest cluster cold front in the sky. *Mon. Not. R. Astron. Soc.* **455**, 846–858 (2016). <https://doi.org/10.1093/mnras/stv2358>. arXiv:1506.06429
- L.M. Widrow, D. Ryu, D.R.G. Schleicher, K. Subramanian, C.G. Tsagas, R.A. Treumann, The first magnetic fields. *Space Sci. Rev.* **166**, 37–70 (2012). <https://doi.org/10.1007/s11214-011-9833-5>. arXiv:1109.4052
- J. Wiener, S.P. Oh, F. Guo, Cosmic ray streaming in clusters of galaxies. *Mon. Not. R. Astron. Soc.* **434**, 2209–2228 (2013). <https://doi.org/10.1093/mnras/stt1163>. arXiv:1303.4746
- J. Wiener, E.G. Zweibel, S.P. Oh, High  $\beta$  effects on cosmic ray streaming in galaxy clusters. *Mon. Not. R. Astron. Soc.* **473**, 3095–3103 (2018). <https://doi.org/10.1093/mnras/stx2603>. arXiv:1706.08525
- M.A.G. Willson, Radio observations of the cluster of galaxies in Coma Berenices—the 5C4 survey. *Mon. Not. R. Astron. Soc.* **151**, 1–44 (1970). <https://doi.org/10.1093/mnras/151.1.1>
- D. Wittor, T. Jones, F. Vazza, M. Brüggén, Evolution of vorticity and enstrophy in the intracluster medium. *Mon. Not. R. Astron. Soc.* **471**, 3212–3225 (2017a). <https://doi.org/10.1093/mnras/stx1769>. arXiv:1706.02315
- D. Wittor, F. Vazza, M. Brüggén, Testing cosmic ray acceleration with radio relics: a high-resolution study using MHD and tracers. *Mon. Not. R. Astron. Soc.* **464**, 4448–4462 (2017b). <https://doi.org/10.1093/mnras/stw2631>. arXiv:1610.05305
- H. Xu, H. Li, D.C. Collins, S. Li, M.L. Norman, Turbulence and dynamo in galaxy cluster medium: implications on the origin of cluster magnetic fields. *Astrophys. J. Lett.* **698**, L14–L17 (2009). <https://doi.org/10.1088/0004-637X/698/1/L14>. arXiv:0905.2196
- H. Xu, H. Li, D.C. Collins, S. Li, M.L. Norman, Evolution and distribution of magnetic fields from active galactic nuclei in galaxy clusters. II. The effects of cluster size and dynamical state. *Astrophys. J.* **739**, 77 (2011). <https://doi.org/10.1088/0004-637X/739/2/77>. arXiv:1107.2599
- V. Yakhot, K.R. Sreenivasan, Anomalous scaling of structure functions and dynamic constraints on turbulence simulations. *J. Stat. Phys.* **121**, 823–841 (2005). <https://doi.org/10.1007/s10955-005-8666-6>. arXiv:nlin/0506038
- F. Zandanel, S. Ando, Constraints on diffuse gamma-ray emission from structure formation processes in the Coma cluster. *Mon. Not. R. Astron. Soc.* **440**, 663–671 (2014). <https://doi.org/10.1093/mnras/stu324>. arXiv:1312.1493
- Y.B. Zel'dovich, The hypothesis of cosmological magnetic inhomogeneity. *Sov. Astron.* **13**, 608 (1970)
- I.B. Zeldovich, A.A. Ruzmaikin, D.D. Sokolov, *Magnetic Fields in Astrophysics: The Fluid Mechanics of Astrophysics and Geophysics*, vol. 3 (Gordon & Breach, New York, 1983)
- I. Zhuravleva, E. Churazov, A.A. Schekochihin, S.W. Allen, P. Arévalo, A.C. Fabian, W.R. Forman, J.S. Sanders, A. Simionescu, R. Sunyaev, A. Vikhlinin, N. Werner, Turbulent heating in galaxy clusters brightest in X-rays. *Nature* **515**, 85–87 (2014). <https://doi.org/10.1038/nature13830>. arXiv:1410.6485
- E. Zinger, A. Dekel, Y. Birnboim, D. Nagai, E. Lau, A.V. Kravtsov, Cold fronts and shocks formed by gas streams in galaxy clusters. *Mon. Not. R. Astron. Soc.* **476**, 56–70 (2018). <https://doi.org/10.1093/mnras/sty136>. arXiv:1609.05308
- V.N. Zirakashvili, V.S. Ptuskin, Diffusive shock acceleration with magnetic amplification by nonresonant streaming instability in supernova remnants. *Astrophys. J.* **678**, 939–949 (2008). <https://doi.org/10.1086/529580>. arXiv:0801.4488
- J.A. Zuhone, E. Roediger, Cold fronts: probes of plasma astrophysics in galaxy clusters. *J. Plasma Phys.* **82**(3), 535820301 (2016). <https://doi.org/10.1017/S0022377816000544>. arXiv:1603.08882

- J.A. ZuHone, M. Markevitch, D. Lee, Sloshing of the magnetized cool gas in the cores of galaxy clusters. *Astrophys. J.* **743**, 16 (2011). <https://doi.org/10.1088/0004-637X/743/1/16>. [arXiv:1108.4427](https://arxiv.org/abs/1108.4427)
- J.A. ZuHone, M. Markevitch, G. Brunetti, S. Giacintucci, Turbulence and radio mini-halos in the sloshing cores of galaxy clusters. *Astrophys. J.* **762**, 78 (2013). <https://doi.org/10.1088/0004-637X/762/2/78>. [arXiv:1203.2994](https://arxiv.org/abs/1203.2994)
- J.A. ZuHone, G. Brunetti, S. Giacintucci, M. Markevitch, Testing secondary models for the origin of radio mini-halos in galaxy clusters. *Astrophys. J.* **801**, 146 (2015a). <https://doi.org/10.1088/0004-637X/801/2/146>. [arXiv:1403.6743](https://arxiv.org/abs/1403.6743)
- J.A. ZuHone, M.W. Kunz, M. Markevitch, J.M. Stone, V. Biffi, The effect of anisotropic viscosity on cold fronts in galaxy clusters. *Astrophys. J.* **798**, 90 (2015b). <https://doi.org/10.1088/0004-637X/798/2/90>. [arXiv:1406.4031](https://arxiv.org/abs/1406.4031)