



Correction to: Tracking the Limnoecological History of Lake Hiidenvesi (Southern Finland) Using the Paleolimnological Approach

Tomi P. Luoto · Marttiina V. Rantala · Mira H. Tammelin

Published online: 5 September 2019
© Springer Nature Switzerland AG 2019

Correction to: Water Air Soil Pollut December 2017, 228:461
<https://doi.org/10.1007/s11270-017-3622-z>

Erroneous information considering Chironomidae and Chaoboridae accumulation was given in Figure 4 published in Luoto et al. (2017). Therefore, it cannot be stated as in the results (page 5) that the chironomid flux increased towards the present. This data is not interpreted or discussed in the manuscript any further, but as a result from the incorrect accumulation unit, the midge accumulation data cannot be used for reference in other research. The volumetric midge sample size given in the methods (page 3) is the

maximum sample size used. Also the midge percentage information in Figure 4 is incorrect. However, the correct percentages were otherwise used throughout the manuscript, including the midge-based reconstructions.

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

The online version of the original article can be found at
<https://doi.org/10.1007/s11270-017-3622-z>

T. P. Luoto (✉)
Department of Environmental Sciences, University of Helsinki,
Niemenkatu 73, FI-15140 Lahti, Finland
e-mail: tomi.luoto@helsinki.fi

M. V. Rantala
Department of Geosciences and Geography, University of
Helsinki, P.O. Box 64, FI-00014 Helsinki, Finland

M. H. Tammelin
Department of Geography and Geology, University of Turku,
FI-20014 Turku, Finland