



Correction to: Comparative cone-beam computed tomography evaluation of the osseous morphology of the temporomandibular joint in temporomandibular dysfunction patients and asymptomatic individuals

Yasin Yasa¹ · Hayati Murat Akgül²

Published online: 13 March 2018

© Japanese Society for Oral and Maxillofacial Radiology and Springer Nature Singapore Pte Ltd. 2018

Correction to: Oral Radiol (2018) 34:31–39
<https://doi.org/10.1007/s11282-017-0279-7>

In the original publication of the article in “Abstract”, the sentence that reads as “The present analyses suggest that a steeper articular eminence inclination may be risk factor” should read as “As a result of our analysis, we concluded that a low eminence angle may be risk factor”

In Conclusion the sentence that reads as “the articular eminence inclination was steeper in the TMD group than in the asymptomatic group in the present study” should read as “The articular eminence inclination was steeper in the asymptomatic group than in the TMD group”.

The original article can be found online at <https://doi.org/10.1007/s11282-017-0279-7>.

✉ Yasin Yasa
yasayasin@yahoo.com

¹ Department of Oral and Maxillofacial Radiology, Faculty of Dentistry, Ordu University, Kumbaşı Mahallesi, Güzelyalı Mh. 94 Sk. No: 2, Altınordu, 52100 Ordu, Turkey

² Department of Oral and Maxillofacial Radiology, Faculty of Dentistry, Ataturk University, Yakutiye, 25240 Erzurum, Turkey