



Correction to: Next Generation Sequencing Bulk Segregant Analysis of Potato Support that Differential Flux into the Cholesterol and Stigmasterol Metabolite Pools Is Important for Steroidal Glycoalkaloid Content

Kacper Piotr Kaminski¹ · Kirsten Kørup² ·
Mathias Neumann Andersen² · Mads Sønderkær¹ ·
Mette Sondrup Andersen¹ · Hanne Grethe Kirk³ ·
Kåre Lehmann Nielsen¹



Published online: 29 September 2018
© European Association for Potato Research 2018

Correction to: Potato Research (2016) 59:81–97
<https://doi.org/10.1007/s11540-015-9314-4>

The original version of this article unfortunately contained a mistake. The Electronic Supplementary Materials (ESMs) were not captured in typesetting. The two perl scripts BAM2coverage.pl and CalculateSkewness.pl are now given in the ESM link provided below.

The online version of the original article can be found at <https://doi.org/10.1007/s11540-015-9314-4>

Electronic supplementary material The online version of this article (<https://doi.org/10.1007/s11540-018-9402-3>) contains supplementary material, which is available to authorized users.

✉ Kacper Piotr Kaminski
kac@bio.aau.dk

Kirsten Kørup
kirstenkoerup@agro.au.dk

Mathias Neumann Andersen
mathiasn.andersen@agro.au.dk

Mads Sønderkær
mson@bio.aau.dk

Mette Sondrup Andersen
msa@bio.aau.dk

Hanne Grethe Kirk
hgk@lkfvandel.dk

Kåre Lehmann Nielsen
kln@bio.aau.dk

¹ Department of Chemistry and Bioscience, Aalborg University, Fredrik Bajers Vej 7H, 9220 Aalborg, Denmark

² Department of Agroecology, Faculty of Science and Technology, Aarhus University, Blichers Allé 20, 8830 Tjele, Denmark

³ Danish Potato Breeding Foundation, Grindstedvej 55, 7184 Vandel, Denmark