



Correction to: Learning curves of two different techniques for the intra-articular injection of the knee joint under fluoroscopic guidance

Paolo Simoni^{1,3} · Olivier Malaise³ · Mounia El Hachemi² · Angelo Tromba² · Grammatina Boitsios¹

Published online: 17 February 2018
© Italian Society of Medical Radiology 2018

Correction to: La radiologia medica
<https://doi.org/10.1007/s11547-017-0847-2>

The published version of this article contained a mistake. In Table 1 of the original article, the numbers of successful injections and failed injections were not correct. The correct Table 1 should read as given below:

The original article can be found online at <https://doi.org/10.1007/s11547-017-0847-2>.

✉ Paolo Simoni
paoloemiliosimoni@gmail.com

- ¹ “Reine Fabiola” Children’s University Hospital, Université Libre de Bruxelles (ULB), 15, Avenue Jean Joseph Crocq, 1020 Brussels, Belgium
- ² Radiology Department, CHU de Liège, CHU du Sart Tilman, Bât. 35, 4000 Liège, Belgium
- ³ Rheumatology Department, CHU de Liège, CHU du Sart Tilman, Bât. 35, 4000 Liège, Belgium

Table 1 Comparison between supine and lateral techniques

Variable	Overall (<i>n</i> = 177)	Supine technique (<i>n</i> = 101)	Lateral technique (<i>n</i> = 76)	<i>p</i> value
Operator				0.92
Operator 1	56 (31.6)	33 (32.7)	23 (30.3)	
Operator 2	63 (35.6)	36 (35.6)	27 (35.5)	
Operator 3	58 (32.8)	32 (31.7)	26 (34.2)	
Sex				
Female	72 (40.7)	40 (39.6)	32 (42.1)	0.74
Male	105 (59.3)	61 (60.4)	44 (57.9)	
Age (years)	45.2 ± 15.0	45.6 ± 14.7	44.6 ± 15.5	0.65
Weight (kg)	77.9 ± 14.8	78.5 ± 13.6	76.9 ± 16.4	0.48
Height (cm)	172.1 ± 9.76	172.5 ± 9.06	171.5 ± 10.7	0.50
BMI (kg/m ²)	26.3 ± 4.48	26.4 ± 4.10	26.1 ± 4.96	0.69
Injection				0.027
Successful	151 (85.3)	81 (80.2)	70 (92.1)	
Failed	26 (14.7)	20 (19.8)	6 (7.9)	
NSAID				0.50
Yes	150 (84.8)	84 (83.2)	66 (86.8)	
No	27 (15.3)	17 (16.8)	10 (13.2)	
Previous trauma or surgery				0.081
No	109 (61.9)	57 (56.4)	52 (69.3)	
Yes	67 (38.1)	44 (43.6)	23 (30.7)	
VAS before injection	2.01 ± 2.55	2.23 ± 2.55	1.71 ± 2.54	0.18
VAS injection	4.10 ± 2.83	4.36 ± 2.68	3.76 ± 3.00	0.16
VAS difference	2.10 ± 2.63	2.13 ± 2.31	2.05 ± 3.03	0.85
(<i>p</i> value)*	(< 0.0001)	(< 0.0001)	(< 0.0001)	
Fluoroscopy (s)	7.06 ± 4.88	7.58 ± 5.26	6.37 ± 4.25	0.10
Extravasation				<0.0001
No	95 (53.7)	34 (33.7)	61 (80.3)	
Yes	82 (46.3)	67 (66.3)	15 (19.7)	

Statistically significant values (*p* < 0.05) are in bold

* Student's *t* test for paired samples