



Correction to: Composition of unfermented, unroasted, roasted cocoa beans and cocoa shells from Peninsular Malaysia

Baizura Aya Putri Agus¹ · Nurul Nadzirah Mohamad¹ · Norhayati Hussain^{1,2}

Published online: 14 September 2018
© Springer Science+Business Media, LLC, part of Springer Nature 2018

Correction to:
Journal of Food Measurement and Characterization
<https://doi.org/10.1007/s11694-018-9875-4>


The original version of this article unfortunately contained an error in Fig. 1. Corrected version of Fig. 1 is given below.

In Results and Discussion, Fat Content subsection, the sentence “Fat content of unroasted cocoa beans (42.65%) was significantly ($p < 0.05$) higher compared to roasted

cocoa beans (30.00%), unroasted cocoa shells (4.87%), roasted cocoa shells (3.09%) and unfermented cocoa beans (0.32%).” should read as below:

Fat content of unroasted cocoa beans (42.65%) was significantly ($p < 0.05$) higher compared to unfermented cocoa beans (41.51%), roasted cocoa beans (30.00%), unroasted cocoa shells (4.87%) and roasted cocoa shells (3.09%).

Fig. 1 Fat content of Peninsular Malaysia unfermented, unroasted, roasted cocoa beans and cocoa shells. Values are expressed as mean \pm standard deviation ($n = 4$). Means with different letters were significantly different at level of $p < 0.05$



The original article can be found online at <https://doi.org/10.1007/s11694-018-9875-4>.

✉ Norhayati Hussain
aryatihussain@upm.edu.my

¹ Department of Food Technology, Faculty of Food Science and Technology, Universiti Putra Malaysia, Serdang, Selangor, Malaysia

² Halal Products Research Institute, Universiti Putra Malaysia, Putra Infoport, 43400 Serdang, Selangor, Malaysia