

Erratum to: Primary PCI in dabigatran-treated patient: is transradial approach and bivalirudin infusion a safe and effective therapeutic option?

Francesco Summaria · Enrico Romagnoli ·
Marina Mustilli · Roberto Patrizi · Adolfo Pagnanelli

Published online: 6 February 2014
© SIMI 2014

Erratum to: Intern Emerg Med
DOI 10.1007/s11739-013-1037-4

The authors would like to correct the editing error in the original publication of Letter to the Editor. The value found in the second column, first paragraph, 0.5 mg/kg should be corrected to 0.75 mg/kg.

The line should read as:

Considering the evidence of a high intracoronary thrombotic burden, an i.v. bolus of bivalirudin (0.75 mg/kg) followed by infusion (1.75 mg/kg/h) was started and a manual thrombus aspiration was performed (Fig. 1b).

The online version of the original article can be found under doi:[10.1007/s11739-013-1037-4](https://doi.org/10.1007/s11739-013-1037-4).

F. Summaria (✉) · E. Romagnoli · R. Patrizi
Interventional Cardiology, Cath Lab Policlinico Casilino,
Via Casilina, 1049, 00168 Rome, Italy
e-mail: f.summaria@gmail.com

M. Mustilli
CCU S. Pertini Hospital, Rome, Italy

A. Pagnanelli
Emergency Department Policlinico Casilino, Rome, Italy