

Erratum to: Mechanism of Phospholipid Hydrolysis for Oyster *Crassostrea plicatula* Phospholipids During Storage Using Shotgun Lipidomics

Qinsheng Chen¹ · Xincen Wang² · Peixu Cong¹ · Yanjun Liu¹ · Yuming Wang¹ · Jie Xu¹ · Changhu Xue¹

Published online: 13 November 2017
© AOCS 2017

Erratum to: Lipids <https://doi.org/10.1007/s11745-017-4305-7>

The authors would like to correct two figures from the original version of their article. The corrections to Figs. 4 and 7 are as follows:

1. Fig. 4

The three CH₂ groups between O and NH₂ have been corrected to show two CH₂ groups. For EPA, the vertical dashed line from the O=C–O bond has been moved to the O–CH bond.

The original article can be found online at <https://doi.org/10.1007/s11745-017-4305-7>.

✉ Jie Xu
xujie9@ouc.edu.cn

Qinsheng Chen
chenqinsheng2017@hotmail.com

Xincen Wang
294580785@qq.com

Peixu Cong
cpx69@139.com

Yanjun Liu
fs-lauyang@hotmail.com

Yuming Wang
yumingwang@ouc.edu.cn

Changhu Xue
xuech@ouc.edu.cn

¹ College of Food Science and Engineering, Ocean University of China, No. 5, Yu Shan Road, Qingdao 266003, Shandong Province, China

² College of Tea and Food Science, Wuyi University, No. 16, Wu Yi Avenue, Wuyishan 354300, Fujian Province, China




Fig. 4 –ESI/MS² product ion spectrum of a precursor ion at m/z 750.6 PakEtn (18:0/20:5) from the oyster *Crassostrea plicatula*

2. Fig. 7

The first diglyceride produced by PtdCho should have contained an oxygen atom, thus the vertical dashed line from the CH₂–O bond has been moved to the O–P bond. The same is true for PtdOH, so the vertical dashed line from the P–O bond has been moved to the O–CH₂ bond. For Lyso-PtdCho (16:0), the vertical dashed line from the O–CH bond has been moved to the O=C–O bond. In addition, the carbon number of EPA has been corrected to 20, instead of 22, near the CH₂–C=O bond.

Fig. 7 Hydrolyzation mechanism of PtdCho (16:0/20:5) by the corresponding enzymes during storage at 4 °C

