

Erratum to: What Demographic and Clinical Characteristics Correlate With Expectations With Trapeziometacarpal Arthritis?

Lana Kang MD, MSc, Joseph Nguyen MPH, Sohaib Z. Hashmi MD,
Steve K. Lee MD, Andrew J. Weiland MD, Carol A. Mancuso MD

Published online: 5 July 2017

© The Association of Bone and Joint Surgeons® 2017

Erratum to: Clin Orthop Relat Res DOI 10.1007/s11999-017-5359-9

The authors of the published study, “What Demographic and Clinical Characteristics Correlate With Expectations With Trapeziometacarpal Arthritis?” would like to correct the following errors found in the Patients and Methods section of their paper.

In the Patients and Methods section, the sentence describing the number of patients who chose injection and the number of patients who chose surgery is incorrect. The sentence should read: “Seventy-three patients chose injection (82%) and 16 patients (18%) chose surgery.”

Second, in the Patients and Methods section, the sentence describing the percentage of patients who listed their dominant hand side as the affected side is incorrect. The sentence should read: “Sixty-four percent of the patients had their dominant side as the affected side.”

Third, in Table 1, under the heading “History of depression or anxiety”, the number of patients who reported a history of depression or anxiety is incorrect. Eight patients (10%), rather than 73, reported depression or anxiety, and 73 patients (90%), rather than 8, reported no history of depression or anxiety.

The authors apologize for these errors.

The online version of the original article can be found under
doi:[10.1007/s11999-017-5359-9](https://doi.org/10.1007/s11999-017-5359-9).

S. Z. Hashmi
Department of Orthopaedic Surgery, Northwestern University,
Chicago, IL, USA

L. Kang (✉), J. Nguyen, S. K. Lee, A. J. Weiland,
C. A. Mancuso
Hospital for Special Surgery, 535 E 70th Street, New York, NY
10021, USA
e-mail: kangl@hss.edu