



Correction to: Effects of a combined exercise plus diet program on cardiorespiratory fitness of breast cancer patients

K. Okumatsu¹ · T. Tsujimoto² · K. Wakaba¹ · A. Seki³ · R. Kotake³ · T. Yamauchi⁴ · S. Hirayama⁵ · H. Kobayashi⁶ · H. Yamauchi³ · K. Tanaka⁷

Published online: 17 September 2018
© The Japanese Breast Cancer Society 2018

Correction to: Breast Cancer

<https://doi.org/10.1007/s12282-018-0889-x>

In the original publication of this article, Table 1 was published incorrectly. The correct Table 1 is given in the following page.

The original article can be found online at <https://doi.org/10.1007/s12282-018-0889-x>.

✉ K. Okumatsu
rits.1159@gmail.com

¹ Graduate School of Comprehensive Human Sciences, Faculty of Health and Sports Sciences, University of Tsukuba, 1-1-1 Tennodai, Ibaraki 305-8574, Japan

² Faculty of Human Sciences, Shimane University, 1060 Nishikawatsucho, Matsue, Shimane 690-8504, Japan

³ Department of Breast Surgical Oncology, St. Luke's International Hospital, 9-1 Akashicho, Chou-ku, Tokyo 104-8560, Japan

⁴ Division of Medical Oncology, Department of Internal Medicine, St. Luke's International Hospital, 9-1 Akashicho, Chou-ku, Tokyo 104-8560, Japan

⁵ Department of Management and Planning, Central Sports Co., Ltd, 1-21-2 Shinkawa, Chuo-ku, Tokyo 104-8255, Japan

⁶ Department of General Medicine, Mito Medical Center, Tsukuba University Hospital, 3-2-7 Miyamachi, Mito, Ibaraki 310-0015, Japan

⁷ Faculty of Health and Sports Sciences, University of Tsukuba, 1-1-1 Tennodai, Tsukuba, Ibaraki 305-8574, Japan

Table 1 Baseline characteristics of study participants

Characteristics	Intervention group (n=21)	Usual-care group (n=11)	P
Age, years (mean±SD)	51±6	53±5	0.14
Race, no. (%)			
Asian (Japanese)	21 (100)	11 (100)	
Education, no. (%)			0.04
High school	6 (29)	0	
Some college	12 (57)	5 (45)	
≥ College	3 (14)	6 (55)	
Time since cancer diagnosis, years (mean±SD)	4±2	3±1	0.14
Disease stage, no. (%)			0.71
0	1 (5)	0 (0)	
I	10 (48)	8 (73)	
II	7 (33)	2 (18)	
III	3 (14)	1 (9)	
Postmenopausal at baseline, (%)	7 (33)	4 (11)	0.62
Cancer treatment, no. (%)			
Surgery	21 (100)	11 (100)	–
Chemotherapy	8 (38)	3 (27)	0.31
Radiation	13 (62)	8 (73)	0.66
Tamoxifen	14 (67)	7 (64)	0.14
Aromatase inhibitor	7 (33)	4 (36)	–
Smoking habit, no. (%)			0.68
Never	16 (76)	8 (73)	
Former	4 (19)	3 (27)	
Current	1 (5)	0 (0)	

SD standard deviation