

CORRECTION

## Correction to: Multicenter, Randomized, Controlled Study Comparing Tafluprost/Timolol Fixed Combination with Latanoprost/Timolol Fixed Combination in Primary Open-Angle Glaucoma and Ocular Hypertension

Katsuyoshi Suzuki · Naomi Otsuka · Hiroko Hizaki · Masayo Hashimoto ·

Yasuaki Kuwayama · On behalf of the Tafluprost/Timolol Versus Latanoprost/Timolol (TTVLT) Study Group

Published online: January 8, 2019  
© Springer Healthcare Ltd., part of Springer Nature 2019

Correction to: Adv Ther (2018) 35:796–808  
<https://doi.org/10.1007/s12325-018-0718-9>

In the original publication, the range to derive the *P* values is incorrectly represented in Tables 2 and 3. The corrected tables are provided below.

---

The original article can be found online at <https://doi.org/10.1007/s12325-018-0718-9>.

K. Suzuki (✉)  
Suzuki Eye Clinic, Yamaguchi, Japan  
e-mail: suzuki\_eye@grace.ocn.ne.jp

N. Otsuka · H. Hizaki · M. Hashimoto  
Santen Pharmaceutical Co., Ltd., Osaka, Japan

Y. Kuwayama  
Fukushima Eye Clinic, Osaka, Japan

**Table 2** Mean ( $\pm$  standard deviation) superficial punctate keratopathy scores (range 0–3)

	TAF/TIM (n=58)	LAT/TIM (n=50) <sup>a</sup>	p-value <sup>b</sup>
Total SPK score			
Week 0	0.7 (0.9)		
Week 6	0.5 (0.8)	p=0.007 <sup>c</sup>	
Week 12	0.4 (0.7)		
Change from BL at Week 12	-0.2 (0.7)	-0.2 (1.1)	0.240
Superior SPK score			
Week 0	0.1 (0.2)		
Week 6	0.0 (0.1)	p=0.500 <sup>c</sup>	
Week 12	0.0 (0.1)		
Change from BL at Week 12	0.0 (0.2)	0.0 (0.3)	0.461
Mid SPK score			
Week 0	0.2 (0.5)		
Week 6	0.2 (0.4)	p=0.549 <sup>c</sup>	
Week 12	0.2 (0.4)		
Change from BL at Week 12	-0.1 (0.4)	-0.2 (0.7)	0.416
Inferior SPK score			
Week 0	0.4 (0.6)		
Week 6	0.3 (0.6)	p=0.035 <sup>c</sup>	
Week 12	0.3 (0.5)		
Change from BL at Week 12	-0.2 (0.5)	0.0 (0.6)	0.085

Bold indicates significant values

BL baseline, LAT latanoprost, SPK superficial punctate keratopathy, TAF tafluprost, TIM timolol

<sup>a</sup> Data missing for one patient in the LAT/TIM group

<sup>b</sup> Calculated using Wilcoxon two-sample test

<sup>c</sup> Calculated using Wilcoxon one-sample test

**Table 3** Mean ( $\pm$  standard deviation) tear break-up times and hyperemia scores (range 0–3)

	TAF/TIM (n=58)	LAT/TIM (n=50) <sup>a</sup>	p-value
TBUT, sec			
Week 0	4.13 (1.85)	4.63 (2.49)	0.241 <sup>b</sup>
Week 6	4.27 (2.04)	4.32 (2.79)	p=0.398 <sup>b</sup>
Week 12	4.32 (2.13)	4.34 (2.67)	–
Change from BL at Week 12	0.19 (2.09)	-0.28 (2.36)	0.274 <sup>b</sup>
Hyperemia score			
Week 0	0.3 (0.5)	0.3 (0.6)	0.722 <sup>c</sup>
Week 6	0.3 (0.5)	0.3 (0.5)	p=1.000 <sup>d</sup>
Week 12	0.4 (0.5)	0.3 (0.5)	–
Change from BL at Week 12	0.1 (0.3)	0.0 (0.3)	0.445 <sup>c</sup>

BL baseline, LAT latanoprost, TAF tafluprost, TBUT tear breakup time, TIM timolol

<sup>a</sup> Data missing for one patient in the LAT/TIM group

<sup>b</sup> Calculated using *t* test

<sup>c</sup> Calculated using Wilcoxon two-sample test

<sup>d</sup> Calculated using Wilcoxon one-sample test